



**Fast computation of  
aerosol optical  
properties**

B. Aouizerats et al.

This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

# Development of an online radiative module for the computation of aerosol optical properties in 3-D atmospheric models: validation during the EUCAARI campaign

**B. Aouizerats<sup>1</sup>, O. Thouron<sup>1</sup>, P. Tulet<sup>1,2</sup>, M. Mallet<sup>3</sup>, L. Gomes<sup>1</sup>, and J. S. Henzing<sup>4</sup>**

<sup>1</sup>CNRM/GAME, Météo-France/CNRS-INSU, URA1357, 42 av G. Coriolis, 31057 Toulouse, France

<sup>2</sup>LACy, Université de La Réunion, 15 av René Cassin, 97715 Saint-Denis, France

<sup>3</sup>LA, Université de Toulouse, 14 av Ed Belin, 31400 Toulouse, France

<sup>4</sup>Netherlands Organisation for Applied Scientific Research, TNO, 80015 Utrecht, The Netherlands

Received: 30 April 2010 – Accepted: 19 May 2010 – Published: 1 June 2010

Correspondence to: B. Aouizerats (benjamin.aouizerats@cnrm.meteo.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Obtaining a good description of aerosol optical properties for a physically and chemically complex evolving aerosol is computationally very expensive at present. The goal of this work is to propose a new numerical module computing the optical properties for complex aerosol particles at low numerical cost so that it can be implemented in atmospheric models. This method aims to compute the optical properties online as a function of a given complex refractive index deduced from the aerosol chemical composition and the size parameters corresponding to the particles.

The construction of look-up tables from the imaginary and the real part of the complex refractive index and size parameters will also be explained. This approach is validated for observations acquired during the EUCAARI campaign on the Cabauw tower during May 2008 and its computing cost is also estimated.

These comparisons show that the module manages to reproduce the scattering and absorbing behaviour of the aerosol during most of the fifteen-day period of observation with a very cheap computationally cost.

## 1 Introduction

While the greenhouse effect on global warming is quite well understood and leads to a quantification of global temperature increases, the effects of aerosol particles on the radiative budget of the atmosphere are still modelled only roughly (Forster et al., 2007). The direct, indirect and semi-direct effects of aerosols may have opposite impacts on the Earth's radiative budget, as noted in the third and fourth reports by the Intergovernmental Panel on Climate Change (IPCC, 2007), and there are still large uncertainties of radiative forcing by anthropogenic aerosols.

Modelling the aerosol particles' size distribution, chemical composition, optical properties, and impacts on radiative forcing is therefore a major objective in the understanding and quantification of the effects of aerosols on the atmosphere radiative

**GMDD**

3, 735–768, 2010

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



budget. Atmospheric research models such as meteorological forecasting models and regional-global climate models, generally use climatology or parametrization of the aerosol optical properties in order to quantify their impacts on radiations (Kinne et al., 2006; Solmon et al., 2008).

5 Studying aerosols with meso- to fine-scale models may imply high variability of the physical and chemical properties, such as the description of size distributions and chemical compositions of urban polluted aerosols (Costabile et al., 2009; Raut and Chazette, 2008). In such cases, the temporal and spatial variability of aerosol size and chemical description is so high that online coupling with a radiative module is the  
10 only way to obtain a good quantification of the aerosol optical properties, radiative direct forcing and feedback. The current methods used to quantify the optical properties are not accurate enough to take the high variability of aerosol composition and size distribution into account.

At those resolutions, it is imperative to consider the ageing of the aerosol and so, the  
15 evolution of the size distribution and the chemical composition of the aerosol particles. For example, the absorptivity of primary absorbing carbonaceous particles is due to the coating of carbonaceous aerosols by secondary hydrophilic particles (Jacobson, 2000; Mikhailov et al., 2006). In parallel, the increase of the particle size also impacts the scattering and absorbing power of the aerosol (Seinfeld and Pandis, 1997). In  
20 that sense, it is necessary to have a good description of both chemical and physical aspects. Modelling the chemical evolution of aerosols and computing the aerosol optical properties as prognostic variables is numerically very expensive. Because of this high cost, which results from an iterative computation of particle optical properties, the computation of aerosol optical properties as a function of its size distribution and chemical  
25 composition is never performed within meso-scale atmospheric models nor within weather models. In several models, the impacts of aerosols on radiation is parameterized either by climatology tables or as a function of the main compounds concentration deduced from the literature (Kinne et al., 2006). In Grini et al. (2006), the effect of dust particles on radiation is studied with an evolving aerosol size distributions, but the

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



aerosol is supposed to be only one kind of chemical composition. We propose a more complex approach, with a number of aerosol compounds taken into account higher than in usual methods (Lesins et al., 2002).

In this context, the goal of the present work is to set up a computationally cheap module depending on a complex physical and chemical aerosol description to compute the optical properties at six short wavelengths.

The EUCAARI field experiment (Kulmala et al., 2009) which took place during May 2008 near Cabauw (Netherlands) provided us with a complete data set of aerosol measurements for the validation of the work. The Cabauw tower was fully equipped with a set of instruments measuring various aerosol properties. The location of the Cabauw site between the North Sea and the industrialized area of Rotterdam allowed different aerosol types, from polluted to maritime air masses, to be observed.

The first section will describe the bases used to compute the aerosol optical properties, and highlight the sensitivity of those parameters as a function of both physical and chemical aerosol description. Then the building of look-up tables in order to save time during the computation will be explained, and the resulting optical properties will be evaluated in regard of Mie computations. Finally, the comparisons between observations acquired during the EUCAARI campaign at the Cabauw tower and computations from the module will be presented.

## 2 Computation of optical parameters

In order to compute the aerosol optical properties, we applied the Mie theory (Mie, 1908) assuming that the aerosol was made up of spherical particles. Relative to other theories that exist to compute the optical properties of a particle, the Mie theory remains the method that is the most commonly used at present. For a given complex refractive index  $K = k_r + i \cdot k_i$  and a given effective radius  $r$  at a wavelength  $\lambda$ , the Mie theory computes the extinction, absorption and scattering, efficiency of the particle which are noted  $Q_{\text{ext,abs,sca}}(r, \lambda)$ , respectively, and the Legendre polynomials of the phase function.

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Because of the high cost of the explicit fluxes computation, the radiative models commonly use the following three optical parameters: the Extinction coefficient or  $b$ , the Single Scattering Albedo (ratio of scattering to extinction) or SSA, and the asymmetry factor,  $g$ , giving the radiative direction information. Considering the aerosol size distribution  $n(r)$ , the optical parameters can be computed for a given radius  $r$  as:

$$b(\lambda) = \int_0^{+\infty} Q_{\text{ext}}(r, \lambda) \pi r^2 n(r) dr$$

$$\text{SSA}(\lambda) = \frac{\int_0^{+\infty} Q_{\text{sca}}(r, \lambda) \pi r^2 n(r) dr}{\int_0^{+\infty} Q_{\text{ext}}(r, \lambda) \pi r^2 n(r) dr}$$

$$g(\lambda) = \frac{\int_0^{+\infty} r^2 n(r) Q_{\text{sca}}(r, \lambda) g'(r, \lambda) dr}{\int_0^{+\infty} r^2 n(r) Q_{\text{sca}}(r, \lambda) dr}$$

One of the first steps to apply the Mie theory to an aerosol particle with a complex chemical composition, is to define a refractive index for the whole aerosol.

It is important to consider an evolving refractive index for the aerosol. Indeed, the main objective of this study is to perform an online computation of the aerosol optical properties considering an aerosol evolving through its size distribution and its chemical composition. The purpose of this study is to set up a module for a large set of aerosol compositions. As discussed in Chylek et al. (2000), the Maxwell-Garnett mixing rule suits the best for situations with many insoluble particles suspended in solution. Yet, the small scale studies about aerosol particles and with a high spatial variability usually take place in urban areas, which precisely show those conditions.

In order to define a refractive index corresponding to an aerosol particle composed of different chemical components, the Maxwell-Garnett equation (Maxwell-Garnett, 1904) as defined in Tombette et al. (2008) allows us to link the chemical composition of the aerosol to a refractive index and then to take the particle size distribution into account.

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.1 The Maxwell-Garnet equation

This approach considers the aerosol as being made up of an inclusion and an extrusion.

The inclusion is composed of the primary and solid parts of the aerosol, whereas the extrusion is composed of the secondary and liquid parts of the aerosol. Then the effective refraction index is:  $k_{\text{aer}} = e^2$

with  $e = e_2 \frac{\epsilon_1 + 2\epsilon_2 + 2f(\epsilon_1 - \epsilon_2)}{\epsilon_1 + 2\epsilon_2 - f(\epsilon_1 - \epsilon_2)}$  where

- $\epsilon_i$  are the complex effective dielectric (square root of the refractive index) constants in which subscripts 1 and 2 stand for the inclusion and the extrusion.
- $f$  is the volumic fraction of inclusion.

In the computation it is assumed that aerosol particles are only composed of primary Organic Carbon (OCp), Black Carbon (BC), Dust, nitrates ( $\text{NO}_3^-$ ), sulphates ( $\text{SO}_4^{2-}$ ), ammonium ( $\text{NH}_4^+$ ), water ( $\text{H}_2\text{O}$ ) and Secondary Organic Aerosols (SOA).

The inclusion or core of the aerosol is then composed of the OCp, the BC and the dust whereas the extrusion or shell is composed of  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{NH}_4^+$ ,  $\text{H}_2\text{O}$  and SOA.

The refractive index for each component considered is as defined by Krekov (1993) and Tulet et al. (2008) and shown in Table 1.

## 2.2 Sensitivity of optical parameters

Here, we conducted various sensitivity tests based on variations of the aerosol size distribution (Sect. 2.3.1) and chemical composition (Sect. 2.3.2). Although these results are well known by the radiation community, the goal of this section is to highlight the non-linearity of the aerosol optical properties evolution and the necessity of the

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



proposed parametrization. In order to show the importance of considering a consistent size distribution and chemical composition for the aerosol, Mie computations were performed with different refractive indexes and different size distributions.

### 2.2.1 Sensitivity to size distribution

5 The size distributions were represented by a lognormal function and described by a median radius and a geometric standard deviation. The effective radius, representing the predominant radius with respect to radiation, is expressed as  $r_{\text{eff}} = r_v^3 / r_s^2$  with  $r_v$ ,  $r_s$  standing respectively for the volume mean radius and surface mean radius depending on the geometric standard deviation.

10 Figure 1 shows that for a given refractive index of 1.55–0.1i, the increase of the median radius and the geometric standard deviation has a direct impact on the mass extinction efficiency distribution. For the three geometric standard deviations considered: 1.55, 1.75 and 1.95, the evolution of the mass extinction efficiency is non-linear with respect to both the median radius and the effective radius, and can show a difference of 40%. The maximum value for each geometric standard deviation considered stands for a different median radius and can reach  $3.6 \text{ m}^2 \text{ g}^{-1}$  for a geometric standard deviation of 1.55 whereas it reaches only  $2.5 \text{ m}^2 \text{ g}^{-1}$  for a geometric standard deviation of 1.95.

15 Because aerosol size distributions are usually expressed according to the median radius and also because the non-linearity of the parameters stand as a function of both effective and median radius, the results will be shown as a function of the median radius in the rest of the study, for ease of understanding. Figure 2 shows the extinction efficiency  $Q_{\text{ext}}$ , drawn in black and the three size distributions corresponding to the same median radius of  $0.045 \mu\text{m}$  and the three geometric standard deviation are overlaid in red blue and green. The mass extinction efficiency results from the integration of the extinction efficiency multiplied by the size distribution (and a factor of  $\pi r^2$ ). Figure 2 shows that the non-linearity of the extinction efficiency versus the median or effective radius mainly comes from the high fluctuations of the extinction efficiency up to  $2 \mu\text{m}$ .

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2.2 Sensitivity to chemical composition

The influence of the chemical composition of the aerosol is shown in Fig. 3. The aerosol is considered to have a fixed median radius of  $0.15\ \mu\text{m}$ , a fixed geometric standard deviation of 1.6, and a fixed total mass concentration of  $10\ \mu\text{g m}^{-3}$ .

We calculated the aerosol SSA for different refractive indexes by using the Maxwell-Garnett equation (see Sect. 2.2), considering an aerosol composed of 20 to 80% of core (primary aerosol), and 80 to 20% of a shell of secondary species composed with the following mass ratio:  $0.75\ \text{NH}_4^+$ ,  $1\ \text{H}_2\text{O}$ ,  $1.35\ \text{SO}_4^{2-}$ ,  $0.9\ \text{NO}_3^-$  and 1 SOA to ensure the electroneutrality of the secondary solution.

The resulting SSA is computed for 3 different core compositions depending of the BC/OCp ratio, and for a core proportion varying from 20 to 80% of the total aerosol mass in steps of 5%. For example, an aerosol composed of 20% primary species with  $\text{BC}=0.25\ \text{OCp}$  inside the core, gives a SSA computed at 550 nm of 0.87 whereas for an aerosol composed of 80% primary species with the same OCp/BC ratio, the SSA value at the same wavelength is 0.64.

It is noteworthy that, for the same primary mass, the core composition is also very important for the computation of the SSA. For a primary mass of 50%, and for as much BC as OCp, the SSA computed reaches 0.62 whereas, with  $\text{BC}=1/4\ \text{OCp}$  the SSA value is 0.77.

This is consistent with our expectations because a low SSA means that the extinction efficiency is mainly caused by absorption, and the SSA has lower values when the black carbon concentration, which is a primary and highly absorbing component, increases.

The same phenomenon is observed at the six wavelengths considered but is not shown here.

**GMDD**

3, 735–768, 2010

### Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3 Methodology for the building of the look-up tables

An analytical solution was used, employing a look-up table of aerosol optical properties and a mathematical analytical function approximating the Mie computation.

In order to minimize the number of stored terms, the construction of the look-up tables was adjusted in two different ways described thereafter.

#### 3.1 Description of the polynomial interpolation method

The first way to minimize the number of stored terms was by approximating the optical parameters with a double 5th degree polynomial interpolation according to the median radius. Considering the median radius as a stored input parameter would be consistent for no more than a hundred cases as pointed out by Grini et al. (2006). To avoid this constraint, the computation and storage of the fifth degree polynomial coefficients that best fitted the optical parameters evolving according to the median radius was taken. In this way, the input terms of the look-up tables are the complex refractive index and the geometric standard deviation. However, in several cases as shown in Fig. 4, the shape of the Mie resulting parameter evolution was poorly reproduced by the best fitted mono polynomial interpolation. The mean relative errors were around 200% and showed a difference of  $1.5 \text{ m}^2 \text{ g}^{-1}$  at the maximum of mass extinction efficiency. For this reason the interpolation of the curve was divided into two different curves defined by the evolution from the first radius considered to radius corresponding to the maximum of the curve  $r_{\text{cut}}$ , and from  $r_{\text{cut}}$  to the last radius considered. Then as shown on Fig. 5, for each case considered, the two fifth degree polynomial coefficient and the  $r_{\text{cut}}$  were stored. The total number of stored terms was 13, which is around an eighth of the 100 stored for a discrete radius computation.

GMDD

3, 735–768, 2010

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To summarize, the polynomial coefficients stored corresponding to the polynomial  $P(r)$  follow the equation:

$$P(r) = \begin{cases} a_{10} + a_{11}r + a_{12}r^2 + a_{13}r^3 + a_{14}r^4 + a_{15}r^5 & \text{if } r < r_{\text{cut}} \\ a_{20} + a_{21}r + a_{22}r^2 + a_{23}r^3 + a_{24}r^4 + a_{25}r^5 & \text{if } r > r_{\text{cut}} \end{cases}$$

where  $r_{\text{cut}}$  is defined as the median radius corresponding to the maximum value of the optical parameter, and the 12  $a_{i,j}$ , with  $i = 1, 2$ ,  $j = 0$  to 5 are stored in the look-up table. The coefficients  $a_{i,j}$  were computed by a least square approach for the fitting of the two parts of the curve.

### 3.2 Range of parameters chosen

The input terms of the look-up tables are then the imaginary and real part of the complex refractive index of the aerosol, and the geometric standard deviation of the size distribution. The second way to minimize the stored terms was by choosing of input terms in the look-up tables so as to obtain pseudo linearity between adjacent pairs of resulting optical properties.

Tests were performed to select a minimum amount of stored terms (not shown here), and six  $k_i$ , eight  $k_r$  and eight  $\sigma$  were chosen because of the pseudo linearity between pairs of terms. This selection method allowed us to interpolate between the results of each optical property corresponding to the two stored values. The stored terms of the input parameters as well as the 6 wavelengths are reproduced in Table 2. As an example, if the value of the geometric standard deviation is 1.52, the corresponding aerosol optical properties are the weighted interpolations between the optical properties corresponding to the stored geometric standard deviations of 1.45 and 1.65.

As in most cases, if we position ourselves between two stored  $k_i$ ,  $k_r$  and  $\sigma$  we compute the eight polynomials corresponding for the eight different combinations and we interpolate with the appropriate weight for obtaining the weighted optical properties. Although the computation is performed at 6 wavelengths which may be considered

## GMDD

3, 735–768, 2010

### Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



as input parameters, there is no possible interpolation between the wavelengths as performed for the other input parameters.

### 3.3 Evaluation of the method in regard of direct Mie computations

In order to evaluate the dual polynomial interpolation method, a comparison between the resulting and the direct Mie computed aerosol optical properties was performed. This comparison was done at the 6 previously explicated  $\lambda$ , for 100 considered values of median radii, and for  $20k_r$ ,  $20k_i$  and  $18\sigma$  equally spaced in order not to be located at the stored values, leading to 4 320 000 comparison points for the 3 optical parameters. Because the computed parameters stands for values of  $k_r$ ,  $k_i$  and  $\sigma$  which require an interpolation by the module (they do not match to the stored values), this comparison allows to evaluate the module in regard of both polynomial interpolation method and choice of input stored terms.

Figure 6 shows the result of the comparison for the three optical properties between the dual polynomial interpolation module and the direct Mie application. On the top (respectively middle and bottom) is represented the mass extinction efficiency, single scattering albedo and asymmetry factor, computed by the dual polynomial interpolation as a function of the same parameter computed from the direct Mie application. Due to the very high number of scatter plots (4 320 000 for each optical parameter), the comparison is shown as a 2-D density function computed over 100 equally splitted bins. For each bin, the 2-D density function is computed as a function of the number of sampling included in the bin. The first coloured contour (dark blue) stands for 99% of the optical parameters computed by the dual polynomial interpolation included within the area. The 1-D density function of the Mie computed optical parameters are represented by the continuous red lines and shows to which optical parameter values do the most cases stand for. The 1-D density function is computed as a function of the total number of sampling.

First, Fig. 6 shows that for the three optical parameter computations, the dual polynomial interpolation method manages to reproduce the values computed by the direct

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## Fast computation of aerosol optical properties

B. Aouizerats et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Mie application. Indeed, almost all of the parameters computed by the module follow the linear curve  $x=y$  represented by a continuous black line. The correlation coefficients value standing for the three optical parameters are respectively 0.9992, 0.9946 and 0.9994. The most large differences stand for extreme cases with very low median radius or very high geometric standard deviation. As an example, the highest values of mass extinction efficiencies which can reach more than  $15 \text{ g m}^{-2}$  stand for very unusual combinations of parameters such as a geometric standard deviation below 1.4, a real part of the refractive index of 1.80. These extreme cases all occur on the first wavelength. We can also notice that for the single scattering albedo, the highest dispersion stand for very low values, meaning that the scattering is mainly absorbing. Moreover, the most computed cases, represented by the continuous red line, occur where the density function is the highest, and at this points there is a good linearity between the two methods.

### 3.4 Note to obtain the look-up tables

The very large number of parameters inside the tables does not allow to write them in the present article. However, the tables are available on request to the corresponding authors (benjamin.aouizerats@cnrm.meteo.fr, pierre.tulet@meteo.fr).

## 4 Validation during the EUCAARI campaign

For the validation of the module, a full set of measured data for the inputs and of the outputs was required. Concerning inputs, a description of the aerosol size distribution giving the median radius and the geometric standard deviation for each mode, associated with the aerosol chemistry giving the aerosol composition was needed. Concerning outputs, the measurements of the aerosol optical properties was also necessary.

During the EUCAARI field experiment, a complete set of instruments were available at the Cabauw tower. Between 15 and 29 May 2008, all the selected instruments requested to test this module were operational.

## 4.1 Description of observations

### 4.1.1 Period and instruments

The data were processed to give one average value over 30 min for each parameter during the 15 days.

- The aerosol size distribution was deduced after merging the SMPS (model TSI 3034) and APS (model TSI 3321) size spectral observations: the SMPS measures the aerosol size distribution between 10 and 500 nm, and the APS between 500 nm and 20  $\mu\text{m}$ . The measurements were considered to have been made in dry conditions ( $\text{RH} < 50\%$ ).

Then, a lognormal fit that best approached the observed distribution was found. At each time step, a least square approach gave the median radius, the geometric standard deviation and the mass concentration for each aerosol mode observed by both instruments. To link the number of aerosol directly measured to the mass of aerosol, the hypothesis of spherical particles with a density of  $2.5 \text{ g cm}^{-3}$  was done.

- The chemical composition of the aerosol was deduced from the observations of an aerosol mass spectrometer (AMS, cTOF type) and a multi angle absorption photometer (MAAP model 5012). The AMS gave us the mass concentration of particulate organic matter (POM),  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ ,  $\text{SO}_4^{2-}$  at each time step and the black carbon (BC) concentration was deduced from the MAAP measurements. The water concentration was deduced from the thermodynamical equilibrium EQSAM (Metzger et al., 2002) on dry conditions.

- The scattering coefficient at 550 nm was measured by a nephelometer (model TSI 3563). A truncation correction is performed on the nephelometer data. The absorption coefficient was measured by the MAAP at 670 nm and deduced at

**GMDD**

3, 735–768, 2010

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



550 nm by measuring the Angström coefficient with an aethalometer. These measurements were also made in dry conditions.

A summary of the instruments and the associated aerosol properties is presented in Table 3.

#### 4.1.2 Methodology and hypotheses

Some assumptions were made from the observations. The choice was to split the POM between primary organic carbon and secondary organic carbon with a 60–40% distribution. A second hypothesis was to split the aerosol compounds equally between modes. We also multiplied the composition concentrations observed by the AMS by a factor deduced from the ratio of masses, itself deduced from the fit of the SMPS+APS size spectrum and the total of masses observed by the AMS and the MAAP. It was noticed that the AMS had a cut-off diameter of 500 nm. Then, a checking was also processed in order to see whether the masses observed by the AMS+MAAP were the same as those observed by the SMPS+APS integrated from 0.01 to 0.5  $\mu\text{m}$ .

#### 4.1.3 First results from instruments

The 15-day period of the study showed different aerosol concentration level due to different regimes. As presented in Fig. 7, the aerosol composition gives us several items of information: from 15 to 17 May, the aerosol shows a continental composition; from 17 to 21 May, the low level of aerosols is characteristic of a scavenging period with a clean air mass and fresh aerosols; from 21 to 29 May, the aerosol comes again from a continental air flux and the secondary fraction of the aerosol, in particular the inorganic species, also grows in proportion. Finally, the 29 May air flux was different and dusts were present during this period. The total measured mass concentration is also represented in Fig. 7 as a black line. Also one can note that the ratio of primary aerosol to secondary aerosol remains quite stable during the fourteen days of measurements.

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 8 shows the evolution of the aerosol size distribution during the fourteen days of measurement. This figure is the result of the fit applied to the SMPS+APS data and used as inputs in the module.

The Fig. 8 shows the evolution of all the size distribution parameters versus time. The median diameter evolves as the geometric standard deviation. The maximum of mass also evolves consequently as already shown in Fig. 7. Finally, the combination of data measuring size description and chemical composition gives a good representation of the period.

## 4.2 Comparison of computed optical properties and observations

Figure 9 shows the single scattering albedo at 550 nm as calculated from the nephelometer and the MAAP measurements (continuous red line) and computed by the module (dashed blue line).

Figure 10 shows the mass extinction efficiency at 550 nm as calculated from observations (continuous red line) and computed by the module (dashed blue line). There were no instruments able to measure the asymmetry factor during the EUCAARI campaign. Therefore, there is no comparison possible with the computed values.

First, the main trend shows that both mass extinction efficiency and single scattering albedo are correctly reproduced by the module during the 15 days. The correlation coefficient for the 15-day period and for the mass extinction efficiency is 0.94, and 0.89 for the single scattering albedo.

Concerning the single scattering albedo, the module mainly manages to reproduce the highest as the lowest values observed, even with high temporal variability. The main error is the overestimation of the single scattering albedo on 20 May when the observations reach 0.5 whereas the computed value is 0.6.

Concerning the mass extinction efficiency comparison, again there is a good agreement between the observations and the computations. However, it is noticeable that from 20 to 24 May, the computations overestimate the mass extinction efficiency by 10 to 80%. To understand why there is a less good correlation especially between 20 and

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



24 May, Fig. 11 shows the evolution of the difference between the observed and computed mass extinction efficiency (in red) during the 15-day period, and the evolution of the relative error on the integrated mass up to 500 nm observed by the SMPS+APS and by the AMS+MAAP.

The discrepancy between observed and computed mass extinction efficiency between 20 and 24 May, may be directly linked to a major difference of masses calculated from by the two different experimental systems, the SMPS+APS for the aerosol size distribution and AMS+MAAP for the aerosol chemical composition. Between 20 and 24 May, the relative error between both systems on the aerosol integrated mass up to 500 nm can reach more than 100% and the major differences between the observed and computed mass extinction efficiency occur at these same maxima. Thus, the main differences observed may probably come from a discrepancy in the instruments in term of mass measurement, leading to an inconsistency between the inputs and outputs of the module.

## 5 Conclusions

In order to quantify the direct and semi-direct effect of an aerosol, it is necessary to compute the aerosol optical properties in atmospheric models depending on an evolving and complex aerosol particle. To study the aerosol particles interaction with radiation in highly temporal and spatial variable locations, such as urban areas, it is necessary to consider a large number of chemical species.

Nevertheless, the high computing cost of the classically applied Mie theory generally leads to a climatological parametrization of aerosol optical properties in regional-climate models.

The numerical computation was performed for six wavelengths using a vectorial computer (NEC SX8 type), in order to compare the time cost of this module with the other atmospheric processes (turbulence, advection, chemistry, aerosol solver, etc.).

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## Fast computation of aerosol optical properties

B. Aouizerats et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The comparison was made between two standard simulations, with and without computation of the aerosol optical properties. The standard simulation required a cpu time per point per time-step of 98.0  $\mu$ s versus the standard simulation with the aerosol optical properties computation has a cpu time per point per time-step of 99.7  $\mu$ s. These results show that the numerical cost of the module was no more than 1.7% of the standard simulation total cost. As an example, the chemistry solving cost of these same simulations is 35.6%.

We can then consider that the previously described module is numerically very cheap relative to other processes and is consequently affordable for most atmospheric modelling.

This work presents a new, computationally cheap module dedicated to the online computation of optical properties according to the particle chemical composition and size distribution.

To minimize the computing cost, the module is founded on look-up tables built by a dual fifth degree polynomial interpolation of the parameters depending on the median radius of the aerosol size distribution. The parameters are the geometric standard deviation of the lognormal size distribution of the aerosol mode considered, and the imaginary and real part of the complex aerosol refractive index corresponding to a chemical composition deduced from the Maxwell-Garnett equation.

The module was then evaluated by using observations acquired during the EUCAARI campaign (May 2008). The numerical cost was also tested.

The comparisons between optical properties modelled by the module and those acquired at the Cabauw tower during fifteen days of measurements shows a good correlation. Furthermore, the numerical cost of the module is shown to be very low, allowing its implementation in atmospheric models treating aerosol size distribution and chemical composition. This optical module is already coupled with the ORILAM aerosol scheme (Tulet et al., 2005, 2006) implemented into the atmospheric research model Meso-NH (Lafore et al., 1998). This development will be used in future works to investigate feedbacks of polluted aerosols on radiation and urban climate (especially impacts

on radiative heating, development of the urban boundary layer and the urban breeze). Such a model could also be used for studying the effect of urban aerosols on UV radiation and the possible feedbacks on atmospheric photochemistry (such as the ozone production) in urban zones.

5 *Acknowledgements.* This work has been partially funded by European Commission 6th Framework program project EUCAARI, contract no 036833-2 (EUCAARI), and by the French National Research Agency (ANR) under the AEROCLOUD program, contract no 06-BLAN-0209. Astrid Kiendler-Scharr from Research Center Juelich, Germany, is acknowledged for providing AMS data. The Meso-NH team is also acknowledged for its support. The topical editor O. Boucher is  
10 also acknowledged for his constructive remarks improving this manuscript.



The publication of this article is financed by CNRS-INSU.

## References

- 15 Chylek, P., Videen, G., Geldart, W., Dobbie, S., and Tso, W.: Effective medium approximation for heterogeneous particles, in: *Light Scattering by Nonspherical Particles: Theory, Measurements and Geophysical Applications*, 273–308, 2000. 739
- Costabile, F., Birmili, W., Klose, S., Tuch, T., Wehner, B., Wiedensohler, A., Franck, U., König, K., and Sonntag, A.: Spatio-temporal variability and principal components of the  
20 particle number size distribution in an urban atmosphere, *Atmos. Chem. Phys.*, 9, 3163–3195, doi:10.5194/acp-9-3163-2009, 2009. 737
- Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D., Haywood, J., Lean, J., Lowe, D., Myrhe, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and van Dorland, R.:

**GMDD**

3, 735–768, 2010

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changes in atmospheric constituents and in radiative forcing, in: *Climate Change 2007: The Physical Science Basis*, Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, 2007. 736

Grini, A., Tulet, P., and Gomes, L.: Dusty weather forecast using the MesoNH atmospheric model., *J. Geophys. Res.*, 111, D19205, doi:10.1029/2005JD007007, 2006. 737, 743

IPCC (Ed.): *Climate change 2007: The scientific basis*. Contribution of working group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, <http://www.ipcc.ch/> (last access: 27 May 2010), 2007. 736

Jacobson, M.: A physically-based treatment of elemental carbon optics: Implications for global direct forcing of aerosols, *Geophys. Res. Lett.*, 27, 217–220, 2000. 737

Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S. E., Bernsten, T., Berglen, T. F., Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, J., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Herzog, M., Horowitz, L., Isaksen, I., Iversen, T., Kirkevåg, A., Kloster, S., Koch, D., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: An AeroCom initial assessment – optical properties in aerosol component modules of global models, *Atmos. Chem. Phys.*, 6, 1815–1834, doi:10.5194/acp-6-1815-2006, 2006. 737

Krekov, M.: *Aerosols Effects on Climate*, University of Arizona Press, 1993. 740, 755

Kulmala, M., Asmi, A., Lappalainen, H. K., Carslaw, K. S., Pöschl, U., Baltensperger, U., Hov, Ø., Brenquier, J.-L., Pandis, S. N., Facchini, M. C., Hansson, H.-C., Wiedensohler, A., and O'Dowd, C. D.: Introduction: European Integrated Project on Aerosol Cloud Climate and Air Quality interactions (EUCAARI) – integrating aerosol research from nano to global scales, *Atmos. Chem. Phys.*, 9, 2825–2841, doi:10.5194/acp-9-2825-2009, 2009. 738

Lafore, J. P., Stein, J., Asencio, N., Bougeault, P., Ducrocq, V., Duron, J., Fischer, C., Hérelil, P., Mascart, P., Masson, V., Pinty, J. P., Redelsperger, J. L., Richard, E., and Vilà-Guerau de Arellano, J.: The Meso-NH Atmospheric Simulation System. Part I: adiabatic formulation and control simulations, *Ann. Geophys.*, 16, 90–109, 1998, <http://www.ann-geophys.net/16/90/1998/>. 751

Lesins, G., Chylek, P., and Lohmann, U.: A study of internal and external mixing scenarios

- and its effect on aerosol optical properties and direct radiative forcing, *J. Geophys. Res.*, 107(D10),4094, doi:10.1029/2001JD000973, 2002. 738
- Maxwell-Garnett, J.: Colours in metal glasses and in metallic films, *Philos. T. Roy. Soc. A*, 203, 385–420, 1904. 739
- 5 Metzger, S., Dentener, F., Pandis, S., and Lelieveld, J.: Gas/aerosol partitioning: 1. A computationally efficient model, *J. Geophys. Res.*, 107, D16, doi:10.1029/2002GL014769, 2002. 747
- Mie, G.: Beiträge zur Optik trüber Medien, speziell kolloidaler Metallösungen, *Ann. Phys.-Leipzig*, 330, 377–445, 1908. 738
- 10 Mikhailov, E. F., Vlasenko, S. S., Podgorny, I. A., Ramanathan, V., and Corrigan, C. E.: Optical properties of soot water drop agglomerates: An experimental study, *J. Geophys. Res.*, 111, 1815–1834, doi:10.1029/2005JD006389, 2006. 737
- Raut, J.-C. and Chazette, P.: Vertical profiles of urban aerosol complex refractive index in the frame of ESQUIF airborne measurements, *Atmos. Chem. Phys.*, 8, 901–919, doi:10.5194/acp-8-901-2008, 2008. 737
- 15 Seinfeld, J. and Pandis, S.: *Atmospheric Chemistry and Physics*, Wiley interscience pub., 1997. 737
- Solmon, F., Chuang, P., Meskhidze, N., and Chen, Y.: Acidic processing of mineral dust iron by anthropogenic compounds over the North Pacific Ocean, *J. Geophys. Res.*, 114, D02305, doi:10.1029/2008JD010417, 2008. 737
- 20 Tombette, M., Chazette, P., Sportisse, B., and Roustan, Y.: Simulation of aerosol optical properties over Europe with a 3-D size-resolved aerosol model: comparisons with AERONET data, *Atmos. Chem. Phys.*, 8, 7115–7132, doi:10.5194/acp-8-7115-2008, 2008. 739
- Tulet, P., Crassier, V., Cousin, F., Shure, K., and Rosset, R.: ORILAM, a three moment lognormal aerosol scheme for mesoscale atmospheric model. On-line coupling into the MesoNH-C model and validation on the ESCOMPTE campaign, *J. Geophys. Res.*, 110, D18201, doi:10.1029/2004JD005716, 2005. 751
- 25 Tulet, P., Grini, A., Griffin, R., and Petitcol, S.: ORILAM-SOA: a computationally efficient model for predicting secondary organic aerosols in 3D atmospheric models., *J. Geophys. Res.*, 111, D23208, doi:10.1029/2006JD007152, 2006. 751
- Tulet, P., Mallet, M., Pont, V., Pelon, J., and Boone, A.: The 7–13 March 2006 dust storm over West Africa: generation, transport, and vertical stratification, *J. Geophys. Res.*, 113, D00C08, doi:10.1029/2008JD009871, 2008. 740, 755

---

**Fast computation of aerosol optical properties**B. Aouizerats et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Fast computation of aerosol optical properties

B. Aouizerats et al.

**Table 1.** Refractive indexes used for the main aerosol species at six wavelength from Krekov (1993) and from Tulet et al. (2008) for the dust.

Specie/ Wavelength	217.5 nm	345 nm	550 nm	925 nm	2.285 $\mu\text{m}$	3.19 $\mu\text{m}$
BC	1.80–0.74i	1.80–0.74i	1.83–0.74i	1.88–0.69i	1.97–0.68i	2.10–0.72i
OCp	1.45–0.001i	1.45–0.001i	1.45–0.001i	1.46–0.001i	1.49–0.001i	1.42–0.0126i
Du	1.448–0.00292i	1.448–0.00292i	1.478–0.01897i	1.4402–0.00116i	1.41163–0.00106i	1.41163–0.00106i
H <sub>2</sub> O	1.36–3.60E–8i	1.34–3.00E–9i	1.33–1.80E–8i	1.33–5.75E–7i	1.31–1.28E–4i	1.42–2.54E–1i
NO <sub>3</sub> <sup>-</sup>	1.53–5.00E–3i	1.53–5.00E–3i	1.53–6.00E–3i	1.52–1.30E–2i	1.51–1.30E–2i	1.35–1.00E–2i
NH <sub>4</sub> <sup>+</sup>	1.52–5.00E–4i	1.52–5.00E–4i	1.52–5.00E–4i	1.52–5.00E–4i	1.51–5.00E–4i	1.35–1.40E–2i
SO <sub>4</sub> <sup>2-</sup>	1.52–5.00E–4i	1.52–5.00E–4i	1.52–5.00E–4i	1.52–5.00E–4i	1.51–5.00E–4i	1.35–1.40E–2i
SOA	1.45–0.001i	1.45–0.001i	1.45–0.001i	1.46–0.001i	1.49–0.001i	1.42–0.0126i

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 2.** Input stored terms into the look-up tables for the polynomial interpolation.

$k_r$	$k_i$	$\sigma$	$\lambda$
1.45	-0.001	1.05	217.5 nm
1.50	-0.006	1.25	345 nm
1.55	-0.008	1.45	550 nm
1.60	-0.02	1.65	925 nm
1.65	-0.1	1.85	2.285 $\mu\text{m}$
1.70	-0.4	2.05	3.19 $\mu\text{m}$
1.75	x	2.25	x
1.80	x	2.45	x

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table 3.** Summary of aerosol measurements on Cabauw tower during the EUCAARI campaign.

	Size distribution	chemistry	optical properties
SMPS	$D_i, \sigma_i M_i$	x	x
APS	$D_i, \sigma_i M_i$	x	x
AMS	x	$M_{\text{POM}}, M_{\text{NO}_3^-}, M_{\text{SO}_4^{2-}}, M_{\text{NH}_4^+}$ ,	x
MAAP	x	$M_{\text{BC}}$	$b_{\text{abs}}$
Nephelometer	x	x	$b_{\text{scat}}$

**Fast computation of aerosol optical properties**

B. Aouizerats et al.




**Fig. 1.** Mass extinction efficiency as a function of the median radius in solid lines and in function of the effective radius in dashed lines for three geometric standard deviation at the wavelength of 550 nm and a complex refractive index of  $1.55-0.1i$ .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Fast computation of aerosol optical properties**

B. Aouizerats et al.




**Fig. 2.** Extinction efficiency overlaid on the three size distributions corresponding to the same geometric standard deviations as in Fig. 1 and the same median radius of  $0.045 \mu\text{m}$  and a refractive index of  $1.55-0.1i$ .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Fast computation of aerosol optical properties**

B. Aouizerats et al.




**Fig. 3.** Single scattering albedo at the 550 nm wavelength versus percentage of primary aerosol for 3 different core compositions with a median radius of 0.15  $\mu\text{m}$  and a geometric standard deviation of 1.65.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Fast computation of aerosol optical properties

B. Aouizerats et al.



**Fig. 4.** Mass extinction efficiency (left) at 550 nm for given values of  $\sigma$  and  $k_r$ ,  $k_i$  as a function of the radius for the Mie computation versus the mono polynomial interpolation and the relative error (right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Fast computation of aerosol optical properties

B. Aouizerats et al.



**Fig. 5.** Mass extinction efficiency (left) at 550 nm for given values of  $\sigma$  and  $k_r$ ,  $k_i$  as a function of the radius for the Mie computation versus the dual polynomial interpolation and the relative error (right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Fast computation of aerosol optical properties**

B. Aouizerats et al.




**Fig. 6.** Comparison of the optical properties (mass extinction efficiency, single scattering albedo and asymmetry factor) computed by the dual interpolation method (y-axis) and the direct Mie application (x-axis) represented by the 2-D density function in color. Overlaid is the 1-D density function (right scale, continuous red line) of the Mie computed parameters, and the continuous black line represents  $x=y$  regression.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Fast computation of aerosol optical properties

B. Aouizerats et al.




**Fig. 7.** Evolution of aerosol chemical composition from AMS+MAAP and mass integrated from the SMPS and APS in  $\mu\text{g}\cdot\text{m}^{-3}$  during the fourteen days of measurement.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Fast computation of  
aerosol optical  
properties**

B. Aouizerats et al.




**Fig. 8.** Evolution of aerosol size distribution fits obtained from SMPS+APS during the fourteen days of measurement.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Fast computation of  
aerosol optical  
properties**

B. Aouizerats et al.




**Fig. 9.** Evolution of the single scattering albedo at 550 nm measured at Cabauw during EUCAARI by the nephelometer and the MAAP, and computed by the module.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Fast computation of aerosol optical properties

B. Aouizerats et al.



**Fig. 10.** Evolution of the mass extinction efficiency in  $\text{g}\cdot\text{m}^{-2}$  at 550 nm measured at Cabauw during EUCAARI and computed by the module.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 11.** Relative error between the mass measured by the AMS+MAAP and that deduced from the SMPS+APS up to 500 nm (blue line) and difference between mass extinction efficiency observed and computed (red line).

## Fast computation of aerosol optical properties

B. Aouizerats et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

