



The potential of multi-organ-on-chip models for assessment of drug disposition as alternative to animal testing

Damiën van Berlo^{1,3}, Evita van de Steeg²,
Hossein Eslami Amirabadi^{1,2} and Rosalinde Masereeuw¹

Abstract

The development of new medicines suffers from attrition, especially in the development pipeline. Eight out of nine drug candidates entering the clinical testing phase fail, mostly due to poor safety and efficacy. The low predictive value of animal models, used in earlier phases of drug development, for effects in humans poses a major problem. In particular, drug disposition can markedly differentiate in experimental animals versus humans. Meanwhile, classic *in vitro* methods can be used but these models lack the complexity to mimic holistic physiological processes occurring in the human body, especially organ–organ interactions. Therefore, better predictive methods to investigate drug disposition in the preclinical phase are needed, for which recent developments in multiorgan-on-chip methods are very promising. To be able to capture human physiology as good as possible, multiorgan-on-chips should feature 1) human cells endogenously expressing main transporters and metabolizing enzymes; 2) organ models relevant for exposure route; 3) individual organs-on-chip connected in a physiologically relevant manner; 4) a tight cellular barrier between the compartments; 5) organ models properly polarized in 3D; 6) allow for sampling in all major compartments; 7) constructed from materials that do not absorb or adsorb the compound of interest; 8) cells should grow in absence of fetal calf serum and Matrigel; 9) validated with a panel of compounds with known characteristics in humans; 10) an integrated computer model translating concentrations to the human situation. Here, an overview of available systems is presented and the difficult route towards a fully validated system is discussed.

Addresses

¹ Utrecht Institute for Pharmaceutical Sciences, Div. Pharmacology, Utrecht University, Utrecht, the Netherlands

² The Netherlands Organisation for Applied Scientific Research (TNO), Zeist, the Netherlands

³ National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands

Corresponding author: van Berlo, Damiën (damiën.van.berlo@rivm.nl)

Current Opinion in Toxicology 2021, 27:8–17

This review comes from a themed issue on **Translational Toxicology**

Edited by **Roos Masereeuw** and **Damiën van Berlo**

Available online 10 May 2021

For a complete overview see the [Issue](#) and the [Editorial](#)

<https://doi.org/10.1016/j.cotox.2021.05.001>

2468-2020/© 2021 The Authors. Published by Elsevier B.V. This is an

open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords

Multiorgan-on-chip, Advanced *in vitro* models, Drug disposition, Pharmacokinetics, ADME, Alternatives to animal testing.

Efficacy, safety, and drug disposition in humans: lost in translation

For many severe, fatal, and debilitating diseases, treatment options are scarce, insufficient, or do not exist at all. Unfortunately, the drug development process is slow and extremely expensive. Regulatory authorities such as the United States Food and Drug Administration and the European Medicines Agency require the use of experimental animals to test absorption, distribution, metabolism, and excretion (ADME), efficacy and safety. It takes around 12 years to bring a drug to the market with costs estimated to amount to at least hundreds of millions of dollars up to \$ 2.6 billion per drug that reaches the market [1]. The main reason for costs in terms of time and funds is the high attrition rate [2]. Currently, only one out of nine drug candidates (11%) that enters the clinical study phase is successfully introduced into the market. The main factor causing attrition is the lack of efficacy and safety in humans, even though available results from animal tests predict otherwise. The result of this is that effective medication for many severe and debilitating human diseases is currently not available.

It is commonly known that effects in experimental animals exposed to a compound of interest can be very different from effects observed in humans. This has been shown for pharmaceuticals [2–5] and for chemicals, especially for their carcinogenicity [6–9]. One of the main causative factors for the low predictivity observed for safety and efficacy when comparing experimental animals and humans is that drug disposition reflected by ADME processes is very different in experimental animals and humans. In other words, ADME of drugs determines their safety and efficacy profile, therefore predicting these processes (and drug metabolism and pharmacokinetics (DMPK), a related,

highly similar concept used for pharmaceuticals) in the preclinical phase is of the utmost importance. In a review by Shanks *et al.*, the concentration of a specific compound that reaches the blood upon oral exposure is compared between several commonly used experimental animal species and humans. The plots show a shotgun pattern when human data were plotted against animal data [10]; the correlation between animal and human data was very low. A recent systematic review comparing human and animal renal clearance measured for 20 renally excreted drugs showed that rats significantly overestimate human renal clearance [11]. Clearly, results in animals do not always translate well to humans.

Despite this loss in translation, animal tests are obligatory for both pharmaceuticals and chemicals marketing. For instance, pharmacokinetic (PK) studies are routinely performed in animals in the drug discovery phase. Typically, these studies involve intravenous and oral administration of a clinically relevant formulation in rats followed by a thorough study of the disposition of a potentially new drug within the organism [12].

Another factor, that is quickly becoming more relevant, is the societal pressure to develop alternatives to animal tests because of ethical considerations. The most recent manifestation is that the US EPA has published a directive in September 2019 in which is stated that funding for animal testing will be reduced by 30% in 2025 and will be abolished completely by 2035.

Classic cell culture usually involves the culture of a commercially available cell line as a monolayer on a 2D surface (culture dish, flask, or wells plate). Medium is applied statically on top of the cells so that they receive nutrients, and periodic replacement is needed to provide fresh nutrients when the medium has been exhausted. Cells and immortalized cell lines are often derived from animal sources (e.g. MDCKII, LLC-PK1 cells) or human tumors (e.g. Caco-2, HT-29 cells). A notable shortcoming of such cell lines for investigation of ADME-DMPK is that they often lack metabolic competence; this means that toxicity can be both overestimated and underestimated because potentially toxic metabolites might not be formed, while a toxic parent compound may not be metabolized into nontoxic metabolites (e.g., Ref. [13]). Such simple systems do provide relevant information about basic features of the PK profile (e.g. intestinal absorption, hepatic metabolic rate, or transporter interactions) or *in vitro* toxicity profile of parent compounds, but not much more than that. Clearly, if we want to investigate more complex processes in humans, we need a higher level of complexity.

At the moment, fully validated alternative *in vitro* models to test ADME-DMPK are not yet available or at least not yet ready for implementation. But there have

been exciting developments in the *in vitro* field in the last decade. Landmark studies on organoid- and organ-on-chip technology have shown the great potential of these technologies to replace the animal as the preferred model to predict the effects on humans. Examples include 1) the improved prediction of treatment efficacy for cystic fibrosis patients is made possible by culturing intestinal organoids [14], 2) the development of organ-on-chip systems in which cells are grown on tubular structures, which improves maturation and differentiation into multiple cell types, allowing for the investigation of the transport of substances across epithelial cell layers from a microfluidic flow mimicking vascular flow [15–17] and 3) the development of hepatic spheroids that express bile acid transporters [18]. Moreover, connecting multiple individual organ-on-chip systems can generate a system with which ADME-DMPK of pharmaceuticals, chemicals, or nutrients can be assessed.

Using multiorgan-on-chip technology to assess ADME-DMPK *in vitro*

Organ-on-chip revolves around the culture of one or more cell systems in a small chamber or bioreactor with a laminar flow, providing nutrients and appropriate levels of shear stress. An organ-on-chip system constitutes the following elements: 1) microfluidic chip, a credit card size device containing microchannels for medium flow and microchambers for cell culturing; 2) compartment to culture cells (commercial cell line, primary cells, stem cell-derived cells or organoid) or tissue segments/slices; 3) microfluidic flow through one or multiple microchannels/tubing providing culture medium and a test compound of interest and may additionally constitute: 4) optionally: scaffold or 3D gel on which cells are grown to simulate physiological structures or create barrier models; 5) Optionally: biosensors to measure endpoints such as shear stress, metabolism, barrier integrity and/or viability, or bio-actuators that stimulate relevant physical stimuli.

Using organ-on-chip, cells can be kept in culture for much longer before they dedifferentiate, undergo apoptosis or become senescent, opening the door to *in vitro* investigation of long-term exposures, in contrast to the acute exposures that are investigated with classic *in vitro* methods. Also because microfluidic flows are laminar, they only mix by diffusion, which means that biochemical gradients can be achieved and controlled more accurately permitting better control of the tissue microenvironment. Another advantage of shear stress induced by microfluidic flow on the cells is the fact that increased cellular maturation as well as differentiation into multiple cell types (especially for adult stem cells and induced pluripotent cell types) present in the organ of origin can be achieved, for example, for kidney [19] and intestine [16,20]. For ADME-DMPK specifically, a very relevant characteristic of organ-on-chip technology

is that multiple organ models in one or more chips can be interconnected. When each of those chips features a barrier system, the uptake, transport, and metabolism of a compound through the system can be monitored. Such a system can be based on a 2D membrane or, depending on the physiological structure of organs, on a hollow fiber membrane mimicking functional parts of organs such as the kidney (tubules [21]), gut (intestinal tube [16]), or liver (bile duct [15]). Different components would be present on either side of the membrane/hollow fiber, and transepithelial transport of compounds can be investigated by this approach.

Of course, the selection of the appropriate individual organ modules in multiorgan-on-chip models highly depends on the mimicked exposure route. Examples of individual organ-on-chip systems that could be integrated in a multiorgan-on-a-chip system for ADME-DMPK assessment are summarized in Table 1. Figure 1 shows how such individual organ-on-chip modules could be connected to physiologically mimic the oral, intravenous, dermal, and inhalatory exposure routes. Here, it should be noted that different setups are possible and can have their value for specific purposes. To be able to completely replace the animal test for ADME-DMPK assessment of a novel compound, it will be necessary to integrate organ-on-chip modules for most relevant organs (or even multiple modules per organ for different organ parts, for example, the proximal


tubule and the glomerulus for the kidney or the alveolar and bronchial regions for the lung). Sometimes very specific questions related to ADME-DMPK need to be answered, for which a more limited number of organ-on-chip modules would be appropriate. It can be important to investigate the interaction between two organ systems, for instance. Importantly, an ADME-OoC system should only be as complex as necessary because, inevitable, there are drawbacks to a more complex system (it certainly is more technologically challenging, but also problematic for validation of such a system, for achieving sufficient cellular function and viability in all integrated cell systems and so forth).

From a technical point of view, research groups that work on multiorgan chips have adopted three strategies to connect individual organ models in one system, as shown in Table 2. The first is to design one chip with different organs that are connected to each other by a simple gravity-driven flow. This is to decrease the complexity of the operation and, therefore, allow increasing the throughput of the experiments. Because of the simpler layout, these systems often benefit from more inert materials to pharmaceutical compounds, avoiding drug absorption and adsorption. The drawback of such systems is that the simplicity of the design reduces the complexity of the microenvironment in the integrated organs. Prof. Michael Shuler and Prof. James Hickman (co-founders Hesperos) are among the pioneers of pumpless multiorgan chips [22]. The second strategy keeps the different organ models in one chip but adds on-chip, e.g. TissUse [23] and CN BIO [24], or off-chip, e.g. Prof. Leclerc's lab [25], peristaltic pumps to the system. This method enables researchers to use more physiologically relevant models, especially barrier models, in their multiorgan chips. This, consequently, comes with a price of more difficult operation and lower throughput than the first strategy. These two first strategies give us the opportunity to use standard static organ models, such as Transwell inserts, in a dynamic microenvironment of a microfluidic system. Although limited, sensors can also be integrated in these systems. The third strategy goes one step further in technical aspects and adds more physiological relevance and sensor technologies. The idea here is to model each organ in separate chips and connect them with tubing to each other to create a modular and independent multiorgan model [26]. This physiological complexity creates technical complexities which makes the operation of such systems more difficult and thus their throughput lower. Up to now, the physiological complexity of the multiorgan chips seems to be proportional to their technical complexity and inversely proportional to the throughput of these systems. More automation, standardization, and high-throughput readout technologies can help improve the relationship between complexity and throughput [27,28].

Table 1 Examples of single organ-on-chip models needed to create an ADME-DMPK-OoC model for each of the exposure routes.

Exposure route	OoC systems needed	Examples
Dermal	Skin-OoC	[34,35];
	Heart-OoC	[36,37];
	Liver-OoC	[38,39];
	Kidney-OoC	[40,41];
	Bile duct-OoC	[42];
	Intestine-OoC	[16,43];
Intravenous	Heart-OoC	See dermal exposure route
	Liver-OoC	See dermal exposure route
	Kidney-OoC	See dermal exposure route
	Bile duct-OoC	See dermal exposure route
	Intestine-OoC	See dermal exposure route
Oral	Stomach-OoC	[44];
	Intestine-OoC	See dermal exposure route
	Liver-OoC	See dermal exposure route
	Kidney-OoC	See dermal exposure route
	Bile duct-OoC	See dermal exposure route
Inhalatory	Lung-OoC	[45–47];
	Heart-OoC	See dermal exposure route
	Liver-OoC	See dermal exposure route
	Kidney-OoC	See dermal exposure route
	Bile duct-OoC	See dermal exposure route
	Intestine-OoC	See dermal exposure route

Figure 1



Main exposure routes (black text) for pharmaceuticals and chemicals and their disposition in relevant organ systems (red text) to be evaluated by a potential multiorgan-on-chip system; created with [BioRender.com](https://www.biorender.com).

The road ahead for drug disposition on a chip

Multiorgan-on-chip technology is still in its infancy, and many challenges need to be overcome before a mature, fully validated ADME-chip system is ready to be implemented for regulatory testing. A number of technical challenges that need to be overcome include the following:

- Connection of the different individual chip modules is a challenge; this should be done in a physiologically relevant manner and the individual systems should receive the appropriate amount of shear stress in order for the cells to properly differentiate and mature.
- It can be difficult to keep such a complex system bubble-free and sterile to prevent infection with pathogens when handling it; for some organ-on-chip systems, this is known to occur when changing medium or treating cells. Automation/robotization of the system can be a great asset in this regard as well as the integration of microsensors.
- Development of a universal medium that does not only keep all different cells alive, but also keeps them functional can be challenging. Due to the laminar flow in a microfluidic system, it might be possible to let various medium types flow through the system; otherwise, separated and isolated medium flows should be integrated into the system. When using

organoid-based models, general organoid medium can work quite well (e.g., Refs. [29,30]).

- Also, to be a truly animal-free test method, it will be necessary to avoid the use of FCS and Matrigel.
- The choice of material for 3D printing of the chips and support structures is important: while PDMS is convenient and practical to work with, it is also known to be able to bind compounds and can release them in an unpredictable manner, which greatly limits the control one has over the intended compound concentration at a cellular site.

Clearly, the more complex an *in vitro* system is, the more difficult and time-consuming it will be to validate it [31,32]. Standardization of all the different components (cell system, 3D-printed chip, scaffold, medium, microfluidic flows, endpoints measured via biosensors etc.) is needed; each individual cell system should be tested for viability, functionality, and barrier function. Moreover, each individual system should still function properly after connecting them to form a predictive drug disposition system allowing study of ADME processes. Standardization of such a system, which is needed to ensure that reproducible results can be achieved, requires integration of microsensors to monitor relevant model parameters and could benefit greatly from extensive automatization/robotization.

Table 2 Multi-OoC systems in the scientific literature applicable for ADME-DMPK assessment.

	Group	Organs	Standardization	Technical complexity	Throughput/ automation	Material	Complexity of microenvironment	Sensors	
ultiorgan system in a single chip with gravity-driven flow	Shuler and Hickman lab (Hesperos)	- Liver and intestine [48,49] - Liver and bone marrow [22,50] - Liver, tumor and marrow [50] - Liver and heart [51] - Liver, heart, muscle and neurons [52] - Liver, fat, kidney, bone marrow and intestine [53]	++	+	+++	++	+	++	
	Sung lab (Hongik University)	- Liver, intestine [54–57] - Liver, tumor [58]	+	++	++	+	+	+	
	Inspiero (Akura™ Flow)	- Liver and tumor [59–61]	++	++	+++	+++	+	+	
Multiorgan system in a single chip with peristaltic flow	TissUse (Humimic)	- Liver and neurospheres [62] - Liver and skin [63–65] - Liver and intestine [63] - Liver and lung [66] - Liver and testis [67] - Liver and kidney [68] - Tumor and skin [65]	++	++	++	++	++	+	
	Griffith lab (MIT & CN BIO)	- Liver and intestine [24,69,70] - Liver, intestine, lung, heart muscle, brain, pancreas, skin and kidney [71]	++	++	++	+++	++	+	
	Leclerc lab (CNRS Paris)	- Liver, intestine [25,72] - Liver and pancreas [73] - Liver and kidney [74]	++	++	++	+	+	++	
	Multiorgan system with multiple chips	Ingber lab (Wyss Institute)	- Gut, liver, kidney, heart, lung, skin, brain, vasculature [26]	+	+	+	+	+++	+
		Shrike (Harvard) & Khademhosseini (UCLA) lab	- Liver and heart [75] - Liver, lung and heart [76]	+	+	+	+	++	+++

A novel *in vitro* model is validated using a fit-for-purpose approach (i.e. the model works for a specific purpose intended). When an ADME-chip is to be used to assess pharmaceuticals in a contract research environment, it should usually comply with the good laboratory practice (GLP) standard which has significant implications; validated and highly standardized protocols are required for cell culture and all used equipment. Implementation in the R&D lab of a specific pharmaceutical company depends on the needs and preferences of that specific company, in terms of regulations little is imposed by the authorities. For regulatory testing of chemicals and food components however, strict regulations exist: an *in vitro* assay should be submitted to the European Union Reference Laboratory for alternatives to animal testing (EURL-ECVAM) so it can be validated by a test laboratory accredited by the European Union Network of Laboratories for the Validation of Alternative Methods (EU-NETVAL). Extensive guidance on how to develop a novel *in vitro* assay is given in the guidance document on Good In vitro Method Practices (GIVIMP), which has been drafted by the OECD [33]. Validation by EURL-ECVAM can be a lengthy procedure and it is important that a system is finalized at some point, because the introduction of small changes may implicate that the system needs to be re-validated. This is highly challenging in a field that is evolving so rapidly; the decision when to start a laborious and lengthy validation of a multiorgan-on-chip system is a tough call to make.

To test predictivity of drug disposition in a multi-organ-on-chip system, its performance should be validated against both human (gold standard, but

availability can be limited) and animal (when human data are not available or not useful) ADME-PK data. Measurement of the concentration of parent compound and their metabolites in each organ compartment (e.g. blood and urine for the kidney, blood and bile duct for the liver, blood and gut lumen for the gut) using quantitative methods such as liquid chromatography-tandem mass spectrometry would be recommended. An integrated physiologically based PK model should be developed to translate the concentrations and volumes in the different compartments of the multiorgan-on-chip (e.g. bile, blood, urine, feces) to their human equivalent. Preferably, compounds should be included in the test battery that are well predicted by current methods (*in vitro* and *in vivo*) as well as compounds for which current systems fail to predict the human ADME properties. Examples of such compounds are well-known drugs such as anti-pyrene and verapamil (high permeability), theophylline and atenolol (low permeability), high organ extraction compounds such as propranolol and morphine), and substances with more complex ADME characteristics include substrates for drug transporter proteins and/or metabolizing enzymes (diclofenac, digoxin, atorvastatin).

This will facilitate thorough assessment of the performance of the model in relation to animal experimentation. Important to note is that human data are the gold standard here, not animal data; a model that would excellently reproduce data obtained in experimental animals would still be of limited value when attempting to predict effects in humans.

Box 1. Recommendations/requirements for an *in vitro* system designed to mimic drug disposition/ADME.

- 1) Cells should be of human origin and should express functional transporter systems and metabolizing enzymes at physiological levels;
- 2) Selected chip modules mimicking specific organ functionalities models should be incorporated for the organs relevant for the exposure route of interest;
- 3) These models should be connected in a physiologically relevant way;
- 4) Cells should form a tight barrier between the different compartments relevant for ADME/PK that are mimicked by the system, such as blood, urine, feces, bile etc.
- 5) Organ systems should be properly 3D-polarized (*i.e.* transport should occur from apex to base or the other way around, just like in the actual organ);
- 6) Preferably, the system should allow for longer term cell culture, so ADME might be investigated upon a repeated dose regimen;
- 7) The system should allow for sampling in all the major compartments that are mimicked (e.g. blood, urine, bile, airway lumen or feces depending on the route of exposure), allowing for monitoring of transport through the system and metabolism of the compound of interest;
- 8) The system should be constructed from materials that do not absorb/adsorb the compound of interest;
- 9) Cells should grow in absence of FCS and Matrigel for the system to be truly animal-free and thus a real alternative to the animal test;
- 10) The system should be validated for a carefully selected panel of compounds with known ADME-DMPK in humans, using appropriate positive and negative controls;
- 11) Integrated physiologically based pharmacokinetic (PBPK) computer model translating *in vitro* findings to the human situation, that is, by upscaling the volumes in the individual organ-on-chip model compartments (blood, urine, bile feces etc.) to the volumes of respective compartments in people so that concentrations in humans can be predicted;
- 12) The system should incorporate a number of microsensors so that model parameters can be closely monitored; preferably, it should be automated/robotized as much as possible to facilitate standardization, which is difficult for such a complex system.

Conclusions and future perspective

Groundbreaking work has been performed in which steps have been taken towards the creation of multi-organ-on-chip models. In **Box 1**, the main characteristics for a multiorgan-on-chip system to provide a true alternative to an animal ADME-DMPK study are summarized. At present, none of the created models fulfill all of the criteria mentioned. When striving to create a model capable of improving on (and replacing) animal tests, close collaboration is needed between a great number of disciplines, that is, experts in cell biology, microfluidics, 3D printing, microelectronics, physiologically based PK modeling, artificial intelligence/machine learning and robotization should or could be linked up. Moreover, connections to R&D labs from the pharmaceutical, chemical and food industry as well as regulating agencies will be crucial to ensure that a system can be readily implemented and complies with regulatory, quality-related and practical requirements.

The complexity of such collaboration between stakeholders from diverse backgrounds requires setting up connective structures (*i.e.* hubs) from where stakeholder communication and project management is coordinated and where a database is present with information on experts in relevant fields. Independence of such structures/hubs would positively affect collaboration.

It is clear that the road ahead is long and is filled with plenty of obstacles, but the rewards upon achieving the goal (a fully developed and validated multiorgan-on-chip model) are great: faster and more cost-effective drug development resulting in better availability of treatment for all kinds of diseases, better testing of chemicals and food components and replacement of ethically sensitive animal tests by *in vitro* methods.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This collaborative effort is co-funded by the PPP Allowance made available by Health~Holland, Top Sector Life Sciences & Health, the Netherlands (grant no. LSHM20045-HSGF) to stimulate public-private partnerships.

References

Papers of particular interest, published within the period of review, have been highlighted as:

- * of special interest
- ** of outstanding interest

1. DiMasi JA, Grabowski HG, Hansen RW: **Innovation in the pharmaceutical industry: new estimates of R&D costs.** *J Health Econ* 2016, **47**. <https://doi.org/10.1016/j.jhealeco.2016.01.012>.
2. Arrowsmith J, Miller P: **Trial watch: phase II and phase III attrition rates 2011-2012.** *Nat Rev Drug Discov* 2013, **12**. <https://doi.org/10.1038/nrd4090>.
3. Heywood R: **Clinical toxicity – could it have been predicted? Post-marketing experience.** In *C. Work. Ser. Anim. Toxic. Stud. – Their Relev. Man.* Edited by Lumley CE, Walker SR, London, UK: Quay Publishing; 1989:57–67.
4. Spriet-Pourra C AM: *Drug withdrawal from sale.* 1994.
5. van Meer PJK, Kooijman M, Gispens-de Wied CC, Moors EHM, Schellekens H: **The ability of animal studies to detect serious post marketing adverse events is limited.** *Regul Toxicol Pharmacol* 2012, **64**:345–349. <https://doi.org/10.1016/j.yrtph.2012.09.002>.
6. Knight A, Bailey J, Balcombe J: **Animal carcinogenicity studies: 1. Poor human predictivity.** *ATLA Altern Lab Anim* 2006, **34**. <https://doi.org/10.1177/026119290603400117>.
7. Haseman JK: **Using the NTP database to assess the value of rodent carcinogenicity studies for determining human cancer risk.** *Drug Metab Rev* 2000, **32**:169–186. <https://doi.org/10.1081/DMR-100100570>.
8. Gold LS, Slone TH, Ames BN: **What do animal cancer tests tell us about human cancer risk?: Overview of analyses of the carcinogenic potency database.** *Drug Metab Rev* 1998, **30**: 359–404. <https://doi.org/10.3109/03602539808996318>.
9. Adler S, Basketter D, Creton S, Pelkonen O, Van Benthem J, Zuang V, Andersen KE, Angers-Loustau A, Aptula A, Bal-Price A, Benfenati E, Bernauer U, Bessens J, Bois FY, Boobis A, Brandon E, Bremer S, Broschard T, Casati S, Coecke S, Corvi R, Cronin M, Daston G, Dekant W, Felter S, Grignard E, Gundert-Remy U, Heinonen T, Kimber I, Kleinjans J, Komulainen H, Kreiling R, Kreysa J, Leite SB, Loizou G, Maxwell G, Mazzatorta P, Munn S, Pfuhrer S, Phrakonkham P, Piersma A, Poth A, Prieto P, Repetto G, Rogiers V, Schoeters G, Schwarz M, Serafimova R, Tähti H, Testai E, Van Delft J, Van Loveren H, Vinken M, Worth A, Zaldivar JM: **Alternative (non-animal) methods for cosmetics testing: current status and future prospects-2010.** *Arch Toxicol* 2011, **85**. <https://doi.org/10.1007/s00204-011-0693-2>.
10. Shanks N, Greek R, Greek J: **Are animal models predictive for humans? *Philos Ethics, Humanit Med* 2009, **4**:2. <https://doi.org/10.1186/1747-5341-4-2>.**
In this review, the authors show that the bioavailability of a drug (*i.e.*, the concentration that reaches the blood) is very different in humans compared to the most commonly used species in pre-clinical animal experiments, *i.e.* dogs, primates and rodents.
11. Jansen K, Pou Casellas C, Groenink L, Wever KE, Masereeuw R: **Humans are animals, but are animals human enough? A systematic review and meta-analysis on interspecies differences in renal drug clearance.** *Drug Discov Today* 2020, **25**. <https://doi.org/10.1016/j.drudis.2020.01.018>.
- Species differences in transporter expression or substrate affinity might lead to prediction errors and limit the translatability of animal data. In an attempt to quantify differences in renal drug clearance between animals and humans, a systematic review and meta-analysis was conducted using published clearance data for a diverse set of renally excreted drugs. Identified differences were related to the drug excretion profile (filtration vs active secretion) and physicochemical drug properties (*e.g.*, physiological charge and molecular weight). The authors demonstrated that rats, despite being the most commonly used model, are an inadequate species for preclinical renal drug clearance testing.
12. Webborn PJ: **The role of pharmacokinetic studies in drug discovery: where are we now, how did we get here and where are we going?** *Future Med Chem* 2014, **6**:1233–1235. <https://doi.org/10.4155/fmc.14.76>.
13. Gomez-Lechon M, Donato M, Lahoz A, Castell J: **Cell lines: a tool for in vitro drug metabolism studies.** *Curr Drug Metabol* 2008, **9**:1–11. <https://doi.org/10.2174/138920008783331086>.
- The authors show that the predictive potential of advanced *in vitro* methods can be superior to animal tests. By studying patient-specific mutations in the cystic fibrosis transmembrane conductance regulator the authors show that the drug responses observed in mini-guts or rectal organoids can be used to predict which patients may be potential responders to the drug. Such organoids can be implemented in multi-

organ-on-chip systems, allowing for standardized and high-throughput analysis.

14. Dekkers JF, Berkers G, Kruisselbrink E, Vonk A, De Jonge HR, Janssens HM, Bronsveld I, Van De Graaf EA, Nieuwenhuis EES, Houwen RHJ, Vleggaar FP, Escher JC, De Rijke YB, Majoor CJ, Heijerman HGM, De Winter-De Groot KM, Clevers H, Van Der Ent CK, Beekman JM: **Characterizing responses to CFTR-modulating drugs using rectal organoids derived from subjects with cystic fibrosis.** *Sci Transl Med* 2016, **8**. <https://doi.org/10.1126/scitranslmed.aad8278>.

A state-of-the-art bile duct model is described in this paper. Liver stem cells from mice were differentiated into cholangiocyte-like cells that were grown on hollow fiber membranes. This model can be integrated in a multi-organ-on-chip approach used to assess ADME-DMPK.

15. Chen C, Jochems PGM, Salz L, Schneeberger K, Penning LC, Van De Graaf SFJ, Beuers U, Clevers H, Geijssen N, Masereeuw R, Spee B: **Bioengineered bile ducts recapitulate key cholangiocyte functions.** *Biofabrication* 2018, **10**. <https://doi.org/10.1088/1758-5090/aac8fd>.

The well-known and well characterized Caco-2 cell line was grown on an ECM-coated hollow fiber membrane under physiological shear stress. This approach resulted in improved phenotype, i.e. a more varied cellular population closer to the cellular variety observed in the human intestine.

16. Jochems PGM, van Bergenhenegouwen J, van Genderen AM, Eis ST, Wilod Versprille LJF, Wichers HJ, Jeurink PV, Garszen J, Masereeuw R: **Development and validation of bioengineered intestinal tubules for translational research aimed at safety and efficacy testing of drugs and nutrients.** *Toxicol Vitro* 2019, **60**. <https://doi.org/10.1016/j.tiv.2019.04.019>.

Development of a high-throughput microfluidic in vitro kidney platform that can be used for drug screening to obtain information about kidney toxicity in an early stage of pharmaceutical development. The authors show that drug-induced kidney injury can be detected robustly by assessing a combination of cell viability, LDH release (i.e., cell membrane integrity) and miRNA release.

17. Vormann MK, Vriend J, Lanz HL, Gijzen L, van den Heuvel A, Hutter S, Joore J, Trietsch SJ, Stuut C, Nieskens TGT, Peters JGP, Ramp D, Caj M, Russel FGM, Jacobsen B, Roth A, Lu S, Polli JW, Naidoo AA, Vulto P, Masereeuw R, Wilmer MJ, Suter-Dick L: **Implementation of a human renal proximal tubule on a chip for nephrotoxicity and drug interaction studies.** *J Pharmaceut Sci* 2021. <https://doi.org/10.1016/j.xphs.2021.01.028>.

A novel model to assess drug-induced cholestasis is described, based on the development of 3D spheroid models from either primary human hepatocytes or HepaRG cells. With this model, it is possible to identify drugs that specifically exhibit cholestatic toxicity early on.

18. Hendriks DFG, Puigvert LF, Messner S, Mortiz W, Ingelman-Sundberg M: **Hepatic 3D spheroid models for the detection and study of compounds with cholestatic liability.** *Sci Rep* 2016, **6**. <https://doi.org/10.1038/srep35434>.
19. Homan KA, Gupta N, Kroll KT, Kolesky DB, Skylar-Scott M, Miyoshi T, Mau D, Valerius MT, Ferrante T, Bonventre JV, Lewis JA, Morizane R: **Flow-enhanced vascularization and maturation of kidney organoids in vitro.** *Nat Methods* 2019, **16**. <https://doi.org/10.1038/s41592-019-0325-y>.
20. Kasendra M, Luc R, Yin J, Manatakis DV, Kulkarni G, Lucchesi C, Sliz J, Apostolou A, Sunuwar L, Obrigewitch J, Jang K-J, Hamilton GA, Donowitz M, Karalis K: **Duodenum Intestine-Chip for preclinical drug assessment in a human relevant model.** *eLife* 2020, **9**:1–23. <https://doi.org/10.7554/eLife.50135>.
21. Jansen J, Fedecostante M, Wilmer MJ, Peters JG, Kreuser UM, Van Den Broek PH, Mensink RA, Boltje TJ, Stamatialis D, Wetzels JF, Van Den Heuvel LP, Hoenderop JG, Masereeuw R: **Bioengineered kidney tubules efficiently excrete uremic toxins.** *Sci Rep* 2016, **6**. <https://doi.org/10.1038/srep26715>.

The authors were able to integrate microelectrode arrays in a pumpless multi-layer multi-organ chip that features several human cell and human cancer cell types. This allows for the investigation of on-target and off-target (e.g., depending on drug metabolism) effects.

22. McAleer CW, Long CJ, Elbrecht D, Sasserath T, Bridges LR, Rumsey JW, Martin C, Schnepper M, Wang Y, Schuler F, Roth AB, Funk C, Shuler ML, Hickman JJ: **Multi-organ system for the evaluation of efficacy and off-target toxicity of**

anticancer therapeutics. *Sci Transl Med* 2019, **11**. <https://doi.org/10.1126/scitranslmed.aav1386>.

23. Kühnl J, Tao TP, Brandmair K, Gerlach S, Rings T, Müller-Vieira U, Przbilla J, Genies C, Jaques-Jamin C, Scheppy A, Marx U, Hewitt NJ, Maschmeyer I: **Characterization of application scenario-dependent pharmacokinetics and pharmacodynamic properties of permethrin and hyperforin in a dynamic skin and liver multi-organ-chip model.** *Toxicology* 2021, **448**. <https://doi.org/10.1016/j.tox.2020.152637>.

24. Tsamandouras N, Chen WLK, Edington CD, Stokes CL, Griffith LG, Cirit M: **Integrated gut and liver microphysiological systems for quantitative in vitro pharmacokinetic studies.** *AAAPS J* 2017, **19**:1499–1512. <https://doi.org/10.1208/s12248-017-0122-4>.

25. Bricks T, Hamon J, Fleury MJ, Jellali R, Merlier F, Herpe YE, Seyer A, Regimbeau J-M, Bois F, Leclerc E: **Investigation of omeprazole and phenacetin first-pass metabolism in humans using a microscale bioreactor and pharmacokinetic models.** *Biopharm Drug Dispos* 2015, **36**:275–293. <https://doi.org/10.1002/bdd.1940>.

An automated robotic system which takes care of automatic operation, dosing and sampling. The system maintained the viability and functionality of eight vascularized organ chips for up to 3 weeks. The automated culture system allows for imaging of cells in the organ chips and repeated sampling of separate compartments. This greatly improves throughput and standardization, and avoids the problems of manual handling in terms of infection with fungi, bacteria or viruses as well as the introduction of bubbles.

26. Novak R, Ingram M, Marquez S, Das D, Delahanty A, Herland A, Maoz BM, Jeanty SSF, Somayaji MR, Burt M, Calamari E, Chalkiadaki A, Cho A, Choe Y, Chou DB, Cronce M, Dauth S, Divic T, Fernandez-Alcon J, Ferrante T, Ferrier J, FitzGerald EA, Fleming R, Jalili-Firoozinezhad S, Grevesse T, Goss JA, Hamkins-Indik T, Henry O, Hinojosa C, Huffstater T, Jang K-J, Kujala V, Leng L, Mannix R, Milton Y, Nawroth J, Nestor BA, Ng CF, O'Connor B, Park T-E, Sanchez H, Sliz J, Sontheimer-Phelps A, Swenor B, Thompson G, Touloumes GJ, Tranchemontagne Z, Wen N, Yadi M, Bahinski A, Hamilton GA, Levner D, Levy O, Przekwas A, Prantl-Baun R, Parker KK, Ingber DE: **Robotic fluidic coupling and interrogation of multiple vascularized organ chips.** *Nat Biomed Eng* 2020. <https://doi.org/10.1038/s41551-019-0497-x>.

27. Chandrasekaran A, Abduljawad M, Moraes C: **Have microfluidics delivered for drug discovery?** *Expet Opin Drug Discov* 2016, **11**:745–748. <https://doi.org/10.1080/17460441.2016.1193485>.

28. Cui P, Wang S: **Application of microfluidic chip technology in pharmaceutical analysis: a review.** *J Pharm Anal* 2019, **9**: 238–247. <https://doi.org/10.1016/j.jpaha.2018.12.001>.

29. Schneeberger K, Sánchez-Romero N, Ye S, van Steenbeek FG, Oosterhoff LA, Pla Palacin I, Chen C, van Wolferen ME, van Tienderen G, Lieshout R, Colemonts-Vroninks H, Schene I, Hoekstra R, Verstegen MMA, van der Laan LJW, Penning LC, Fuchs SA, Clevers H, De Kock J, Baptista PM, Spee B: **Large-scale production of LGR5-positive bipotential human liver stem cells.** *Hepatology* 2020, **72**. <https://doi.org/10.1002/hep.31037>.

30. Kaushik G, Ponnusamy MP, Batra SK: **Concise review: current status of three-dimensional organoids as preclinical models.** *Stem Cell* 2018, **36**. <https://doi.org/10.1002/stem.2852>.

31. Marx U: **Biology-inspired microphysiological systems to advance medicines for patient benefit and animal welfare.** *ALTEX* 2020, **37**:1–30. <https://doi.org/10.14573/altex.2001241>.

32. Rebelo SP, Dehne EM, Brito C, Horland R, Alves PM, Marx U: **Validation of bioreactor and human-on-a-chip devices for chemical safety assessment.** *Adv Exp Med Biol* 2016. https://doi.org/10.1007/978-3-319-33826-2_12.

33. *Guidance document on good in vitro method practices (GIVIMP).* OECD; 2018. <https://doi.org/10.1787/9789264304796-en>.

34. Sriram G, Alberti M, Dancik Y, Wu B, Wu R, Feng Z, Ramasamy S, Bigliardi PL, Bigliardi-Qi M, Wang Z: **Full-thickness human skin-on-chip with enhanced epidermal morphogenesis and barrier function.** *Mater Today* 2018, **21**:326–340. <https://doi.org/10.1016/j.mattod.2017.11.002>.

35. Lukács B, Bajza Á, Kocsis D, Csorba A, Antal I, Iván K, Laki AJ, Erdő F: **Skin-on-a-chip device for ex vivo monitoring of transdermal delivery of drugs—design, fabrication, and testing.** *Pharmaceutics* 2019, **11**. <https://doi.org/10.3390/pharmaceutics11090445>.
36. Sakamiya M, Fang Y, Mo X, Shen J, Zhang T: **A heart-on-a-chip platform for online monitoring of contractile behavior via digital image processing and piezoelectric sensing technique.** *Med Eng Phys* 2020, **75**. <https://doi.org/10.1016/j.medengphy.2019.10.001>.
37. Marsano A, Conficconi C, Lemme M, Occhetta P, Gaudiello E, Votta E, Cerino G, Redaelli A, Rasponi M: **Beating heart on a chip: a novel microfluidic platform to generate functional 3D cardiac microtissues.** *Lab Chip* 2016, **16**. <https://doi.org/10.1039/c5lc01356a>.
38. Wang Y, Wang H, Deng P, Chen W, Guo Y, Tao T, Qin J: **In situ differentiation and generation of functional liver organoids from human iPSCs in a 3D perfusable chip system.** *Lab Chip* 2018, **18**. <https://doi.org/10.1039/c8lc00869h>.
39. Lee H, Chae S, Kim JY, Han W, Kim J, Choi Y, Cho DW: **Cell-printed 3D liver-on-a-chip possessing a liver microenvironment and biliary system.** *Biofabrication* 2019, **11**. <https://doi.org/10.1088/1758-5090/aaf9fa>.
40. Vriend J, Peters JGP, Nieskens TTG, Škovroňová R, Blaimschein N, Schmidts M, Roepman R, Schirris TJJ, Russel FGM, Masereeuw R, Wilmer MJ: **Flow stimulates drug transport in a human kidney proximal tubule-on-a-chip independent of primary cilia.** *Biochim Biophys Acta - Gen Subj* 2020, **1864**. <https://doi.org/10.1016/j.bbagen.2019.129433>.
41. Vriend J, Nieskens TTG, Vormann MK, van den Berge BT, van den Heuvel A, Russel FGM, Suter-Dick L, Lanz HL, Vulto P, Masereeuw R, Wilmer MJ: **Screening of drug-transporter interactions in a 3D microfluidic renal proximal tubule on a chip.** *AAPS J* 2018, **20**. <https://doi.org/10.1208/s12248-018-0247-0>.
42. Du Y, Khandekar G, Llewellyn J, Polachek W, Chen CS, Wells RG: **A bile duct-on-a-chip with organ-level functions.** *Hepatology* 2020, **71**. <https://doi.org/10.1002/hep.30918>.
43. Langerak N, Ahmed HMM, Li Y, Middel IR, Eslami Amirabadi H, Malda J, Masereeuw R, van Rooij R: **A theoretical and experimental study to optimize cell differentiation in a novel intestinal chip.** *Front Bioeng Biotechnol* 2020, **8**. <https://doi.org/10.3389/fbioe.2020.00763>.
44. Lee KK, McCauley HA, Broda TR, Kofron MJ, Wells JM, Hong CI: **Human stomach-on-a-chip with luminal flow and peristaltic-like motility.** *Lab Chip* 2018, **18**. <https://doi.org/10.1039/c8lc00910d>.
45. Huh D, Matthews BD, Mammoto A, Montoya-Zavala M, Hsin HY, Ingber DE: **Reconstituting organ-level lung functions on a chip.** *Science* 2010, **328**:1662–1668. <https://doi.org/10.1126/science.1188302>.
46. Stucki JD, Hobi N, Galimov A, Stucki AO, Schneider-Daum N, Lehr CM, Huwer H, Frick M, Funke-Chambour M, Geiser T, Guenat OT: **Medium throughput breathing human primary cell alveolus-on-chip model.** *Sci Rep* 2018, **8**. <https://doi.org/10.1038/s41598-018-32523-x>.
47. Thacker VV, Dhar N, Sharma K, Barrile R, Karalis K, McKinney JD: **A lung-on-chip model of early m. Tuberculosis infection reveals an essential role for alveolar epithelial cells in controlling bacterial growth.** *eLife* 2020, **9**. <https://doi.org/10.7554/eLife.59961>.
48. Esch MB, Ueno H, Applegate DR, Shuler ML: **Modular, pumpless body-on-a-chip platform for the co-culture of GI tract epithelium and 3D primary liver tissue.** *Lab Chip* 2016, **16**:2719–2729. <https://doi.org/10.1039/c6lc00461j>.
49. Esch MB, Mahler GJ, Stokol T, Shuler ML: **Body-on-a-chip simulation with gastrointestinal tract and liver tissues suggests that ingested nanoparticles have the potential to cause liver injury.** *Lab Chip* 2014, **14**:3081–3092. <https://doi.org/10.1039/c4lc00371c>.
50. Sung JH, Kam C, Shuler ML: **A microfluidic device for a pharmacokinetic-pharmacodynamic (PK-PD) model on a chip.** *Lab Chip* 2010, **10**:446–455. <https://doi.org/10.1039/b917763a>.
51. Oleaga C, Riu A, Rothmund S, Lavado A, McAleer CW, Long CJ, Persaud K, Narasimhan NS, Tran M, Roles J, Carmona-Moran CA, Sasserath T, Elbrecht DH, Kumanchik L, Bridges LR, Martin C, Schnepfer MT, Ekman G, Jackson M, Wang YI, Note R, Langer J, Teissier S, Hickman JJ: **Investigation of the effect of hepatic metabolism on off-target cardiotoxicity in a multi-organ human-on-a-chip system.** *Biomaterials* 2018, **182**:176–190. <https://doi.org/10.1016/j.biomaterials.2018.07.062>.
52. Oleaga C, Bernabini C, Smith AST, Srinivasan B, Jackson M, McLamb W, Platt V, Bridges R, Cai Y, Santhanam N, Berry B, Najjar S, Akanda N, Guo X, Martin C, Ekman G, Esch MB, Langer J, Ouedraogo G, Cotovio J, Breton L, Shuler ML, Hickman JJ: **Multi-Organ toxicity demonstration in a functional human in vitro system composed of four organs.** *Sci Rep* 2016, **6**:1–17. <https://doi.org/10.1038/srep20030>.
53. Mahler GJ, Esch MB, Glahn RP, Shuler ML: **Characterization of a gastrointestinal tract microscale cell culture analog used to predict drug toxicity.** *Biotechnol Bioeng* 2009, **104**:193–205. <https://doi.org/10.1002/bit.22366>.
54. won Jeon J, Choi N, Lee SH, Sung JH: **Three-tissue micro-physiological system for studying inflammatory responses in gut-liver Axis.** *Biomed Microdevices* 2020, **22**:1–11. <https://doi.org/10.1007/s10544-020-00519-y>.
55. Lee DW, Ha SK, Choi I, Sung JH: **3D gut-liver chip with a PK model for prediction of first-pass metabolism.** *Biomed Microdevices* 2017, **19**. <https://doi.org/10.1007/s10544-017-0242-8>.
56. Choe A, Ha SK, Choi I, Choi N, Sung JH: **Microfluidic Gut-liver chip for reproducing the first pass metabolism.** *Biomed Microdevices* 2017, **19**:1–11. <https://doi.org/10.1007/s10544-016-0143-2>.
57. Lee SY, Sung JH: **Gut–liver on a chip toward an in vitro model of hepatic steatosis.** *Biotechnol Bioeng* 2018, **115**:2817–2827. <https://doi.org/10.1002/bit.26793>.
58. Lee H, Kim DS, Ha SK, Choi I, Lee JM, Sung JH: **A pumpless multi-organ-on-a-chip (MOC) combined with a pharmacokinetic–pharmacodynamic (PK–PD) model.** *Biotechnol Bioeng* 2017, **114**:432–443. <https://doi.org/10.1002/bit.26087>.
59. Kim JY, Fluri DA, Kelm JM, Hierlemann A, Frey O: **96-Well format-based microfluidic platform for parallel interconnection of multiple multicellular spheroids.** *J Lab Autom* 2015, **20**:274–282. <https://doi.org/10.1177/2211068214564056>.
60. Kim JY, Fluri DA, Marchan R, Boonen K, Mohanty S, Singh P, Hammad S, Landuyt B, Hengstler JG, Kelm JM, Hierlemann A, Frey O: **3D spherical microtissues and microfluidic technology for multi-tissue experiments and analysis.** *J Biotechnol* 2015, **205**:24–35. <https://doi.org/10.1016/j.jbiotec.2015.01.003>.
61. Frey O, Misun PM, Fluri DA, Hengstler JG, Hierlemann A: **Reconfigurable microfluidic hanging drop network for multi-tissue interaction and analysis.** *Nat Commun* 2014, **5**. <https://doi.org/10.1038/ncomms5250>.
62. Materne EM, Ramme AP, Terrasso AP, Serra M, Alves PM, Brito C, Sakharov DA, Tonevitsky AG, Lauster R, Marx U: **A multi-organ chip co-culture of neurospheres and liver equivalents for long-term substance testing.** *J Biotechnol* 2015, **205**:36–46. <https://doi.org/10.1016/j.jbiotec.2015.02.002>.
63. Maschmeyer I, Hasenberg T, Jaenicke A, Lindner M, Lorenz AK, Zech J, Garbe LA, Sonntag F, Hayden P, Ayehunie S, Lauster R, Marx U, Materne EM: **Chip-based human liver-intestine and liver-skin co-cultures – a first step toward systemic repeated dose substance testing in vitro.** *Eur J Pharm Biopharm* 2015, **95**:77–87. <https://doi.org/10.1016/j.ejpb.2015.03.002>.
64. Wagner I, Materne EM, Brincker S, Süßbier U, Frädrich C, Busek M, Sonntag F, Sakharov DA, Trushkin EV, Tonevitsky AG, Lauster R, Marx U: **A dynamic multi-organ-chip for long-term cultivation and substance testing proven by 3D human liver and skin tissue co-culture.** *Lab Chip* 2013, **13**:3538–3547. <https://doi.org/10.1039/c3lc50234a>.

65. Hübner J, Raschke M, Rüttschle I, Gräßle S, Hasenberg T, Schirrmann K, Lorenz A, Schnurre S, Lauster R, Maschmeyer I, Steger-Hartmann T, Marx U: **Simultaneous evaluation of anti-EGFR-induced tumour and adverse skin effects in a microfluidic human 3D co-culture model.** *Sci Rep* 2018, **8**:1–12. <https://doi.org/10.1038/s41598-018-33462-3>.
66. Schimek K, Frentzel S, Luettich K, Bovard D, Rüttschle I, Boden L, Rambo F, Erfurth H, Dehne EM, Winter A, Marx U, Hoeng J: **Human multi-organ chip co-culture of bronchial lung culture and liver spheroids for substance exposure studies.** *Sci Rep* 2020, **10**:1–13. <https://doi.org/10.1038/s41598-020-64219-6>.
67. Baert Y, Ruetschle I, Cools W, Oehme A, Lorenz A, Marx U, Goossens E, Maschmeyer I: **A multi-organ-chip co-culture of liver and testis equivalents: a first step toward a systemic male reprotoxicity model.** *Hum Reprod* 2020, **35**:1029–1044. <https://doi.org/10.1093/humrep/deaa057>.
- The authors have developed and optimized a multi-organ-on-chip system with liver spheroids and a renal proximal tubule barrier system for long term cell culture. A 14-day repeated dose experiment with cyclosporine with or without rifampicin was performed, showing the potential of such system to go beyond the acute exposure regimens that are typically assessed in vitro.
68. Lin N, Zhou X, Geng X, Drewell C, Hübner J, Li Z, Zhang Y, Xue M, Marx U, Li B: **Repeated dose multi-drug testing using a microfluidic chip-based coculture of human liver and kidney proximal tubules equivalents.** *Sci Rep* 2020, **10**:1–15. <https://doi.org/10.1038/s41598-020-65817-0>.
69. Chen WLK, Edington C, Suter E, Yu J, Velazquez JJ, Velazquez JG, Shockley M, Large EM, Venkataramanan R, Hughes DJ, Stokes CL, Trumper DL, Carrier RL, Cirit M, Griffith LG, Lauffenburger DA: **Integrated gut/liver micro-physiological systems elucidates inflammatory inter-tissue crosstalk.** *Biotechnol Bioeng* 2017, **114**:2648–2659. <https://doi.org/10.1002/bit.26370>.
- In this paper, an ulcerative colitis multi-organ-on-chip system is presented consisting of gut, liver and circulating Treg and Th17 cells. Incorporation of an immune component in an in vitro model is a valuable asset for investigation of the effect of drugs on immunity and immune-mediated effects. This open top microfluidic system allows for easy operation and benefits from low adsorption of drugs in the system.
70. Trapecar M, Communal C, Velazquez J, Maass CA, Huang YJ, Schneider K, Wright CW, Butty V, Eng G, Yilmaz O, Trumper D, Griffith LG: **Gut-liver physiometrics reveal paradoxical modulation of IBD-related inflammation by short-chain fatty acids.** *Cell Syst* 2020, **10**. <https://doi.org/10.1016/j.cels.2020.02.008>. 223–239.e9.
71. Edington CD, Chen WLK, Geishecker E, Kassis T, Soenksen LR, Bhushan BM, Freake D, Kirschner J, Maass C, Tsamandouras N, Valdez J, Cook CD, Parent T, Snyder S, Yu J, Suter E, Shockley M, Velazquez J, Velazquez JJ, Stockdale L, Papps JP, Lee I, Vann N, Gamboa M, Labarge ME, Zhong Z, Wang X, Boyer LA, Lauffenburger DA, Carrier RL, Communal C, Tannenbaum SR, Stokes CL, Hughes DJ, Rohatgi G, Trumper DL, Cirit M, Griffith LG: **Interconnected micro-physiological systems for quantitative biology and pharmacology studies.** *Sci Rep* 2018, **8**:1–18. <https://doi.org/10.1038/s41598-018-22749-0>.
72. Prot JM, Maciel L, Bricks T, Merlier F, Cotton J, Paullier P, Bois FY, Leclerc E: **First pass intestinal and liver metabolism of paracetamol in a microfluidic platform coupled with a mathematical modeling as a means of evaluating ADME processes in humans.** *Biotechnol Bioeng* 2014, **111**:2027–2040. <https://doi.org/10.1002/bit.25232>.
73. Essaouiba A, Okitsu T, Kinoshita R, Jellali R, Shinohara M, Danoy M, Legallais C, Sakai Y, Leclerc E: **Development of a pancreas-liver organ-on-chip coculture model for organ-to-organ interaction studies.** *Biochem Eng J* 2020, **164**. <https://doi.org/10.1016/j.bej.2020.107783>.
74. Jellali R, Paullier P, Fleury MJ, Leclerc E: **Liver and kidney cells cultures in a new perfluoropolyether biochip.** *Sensor Actuator B Chem* 2016, **229**:396–407. <https://doi.org/10.1016/j.snb.2016.01.141>.
75. Zhang YS, Aleman J, Shin SR, Kilic T, Kim D, Shaegh SAM, Massa S, Riahi R, Chae S, Hu N, Avci H, Zhang W, Silvestri A, Nezhad AS, Manbohi A, De Ferrari F, Polini A, Calzone G, Shaikh N, Alerasool P, Budina E, Kang J, Bhise N, Ribas J, Pourmand A, Skardal A, Shupe T, Bishop CE, Dokmeci MR, Atala A, Khademhosseini A: **Multisensor-integrated organs-on-chips platform for automated and continual in situ monitoring of organoid behaviors.** *Proc Natl Acad Sci USA* 2017, **114**:E2293–E2302. <https://doi.org/10.1073/pnas.1612906114>.
76. Skardal A, Murphy SV, Devarasetty M, Mead I, Kang HW, Seol YJ, Zhang YS, Shin SR, Zhao L, Aleman J, Hall AR, Shupe TD, Kleensang A, Dokmeci MR, Jin Lee S, Jackson JD, Yoo JJ, Hartung T, Khademhosseini A, Soker S, Bishop CE, Atala A: **Multi-tissue interactions in an integrated three-tissue organ-on-a-chip platform.** *Sci Rep* 2017, **7**:1–16. <https://doi.org/10.1038/s41598-017-08879-x>.