

TNO Public) TNO 2025 R11935 24 September 2025

Author(s) Johan H. ten Veen

Jeroen Schokker

with support from the scientific board of

Geopark Peelhorst & Maasvallei

Classification report TNO Public
Title TNO Public
Report text TNO Public
Appendices TNO Public

Number of pages 219 (excl. front and back cover)

Number of appendices 2

Sponsor Gemeente Boekel

Project name Begrenzing en beschrijving Geopark Peelhorst/Maasvallei –

Unesco

Project number 060.63548

All rights reserved

No part of this publication may be reproduced and/or published by print, photoprint, microfilm or any other means without the previous written consent of TNO.

© 2025 TNO

Cover: The Peel Boundary Fault with disturbed sediment layers in a research trench near Bakel. (photo: VU – Ronald van Balen)

Contents

1	Introduction	
1.1	Reader's Guide	
2	Name and identity of the aUGGp	5
3 3.1	Location and delineation	
4	Surface area	10
5 5.1 5.2	Physical and geographical characteristics	11
6	Geosites	
7	General geological description	37
Key re	ferences	56
Signat	ure	59
Apper Apper Apper		62 214

Disclaimer

This report includes sections that were originally written in Dutch. This in particular pertains to the Geosite descriptions. For the Dutch-to-English translations the assistance of artificial intelligence (Microsoft Copilot) was used occasionally. While the AI tool was used to enhance efficiency and consistency in translation, all outputs were reviewed by the authors for accuracy and contextual appropriateness. Readers should note that AI-generated translations may not fully capture idiomatic expressions or nuanced meanings inherent in the original Dutch text. The authors bear full responsibility for the final content and interpretations presented.

) TNO Public 1/219

) TNO Public) TNO 2025 R11935

) TNO Public 2/219

1 Introduction

In October 2026, the aspiring Geopark Peelhorst and Maasvallei (PHMV) intends to submit an application to obtain the UNESCO Global Geopark status. For this purpose, it is essential that a final boundary for the Geopark area is determined, with at least 40 Geosites included. This must be scientifically substantiated and described. The Steering Committee for PHMV has requested TNO – Geological Survey of the Netherlands (TNO-GDN) to draft both the delineation and description for the Geopark and to characterize 40 selected geosites. A key requirement is that the scientific description sufficiently underpins the selection of the aspiring UNESCO Global Geopark (aUGGp) area in terms of its international geological significance. This report outlines the choices regarding the boundary, the area's size, and the number of Geosites included within. Furthermore, a comparison is made with Geoparks holding the UNESCO status within a radius of approximately 100 km.

1.1 Reader's Guide

In line with the assignment, this geological description of the Geopark Peelhorst & Maasvallei initially draws upon available scientific literature and reports, such as the Landscape Biography and Field Guide "Breuken in het Land van Peel en Maas". These sources already describe visible and accessible locations within the Peelhorst & Maasvallei region. The descriptions provided in these sources have been reviewed to ensure they are sufficiently up-to-date and comprehensive, reflecting the latest scientific insights. Where necessary, they have been updated.

For illustrations, existing publications have been used, alongside material available from TNO-GDN (including GIS map data), illustrative material from the provinces of Noord-Brabant and Limburg and local municipalities, as well as freely available sources elsewhere.

The structure of the report partially follows the UNESCO guidelines for preparing the Application Dossier (or bid book, see Figure 1). The first section of the report provides the "General information" (section 2A of the application dossier) and describes the "Name and Identity", "location and delineation", the "surface area", and the "physical and geographical characteristics" of the aUGGP. Key in this description are the geological elements and a plethora of manifestation that, together, shape the storylines of the aUGGp. The second part complies with section 2C (Main geological highlights) and presents a list of the 40 geosites selected. Note that the full description of the geosites is presented as appendix (cf. UNESCO quidelines). The third part of this report will support section 2E of the application dossier (Verification of UGGP criteria) and provides a "General geological description". Some repetition occurs between this part and the first part on "physical and geographical characteristics", the context of both descriptions is different, however. The required bibliography (section 3.6) is largely based on a version compiled by the Scientific Board of the aUGGP and supplemented where relevant, but at the moment not yet complete. By following this structure, the report supports the application dossier, but is not meant to be the complete application text as it does not address educational, geo-touristic, and administrative aspects.

) TNO Public 3/219

With the above in mind, it is important to mention that this report is a data source from which information can be selected for preparation of the application dossier (bid book). In support of that process, the scientific content and terminology presented herein can used directly. It is acknowledged though that language, style, and formulations can subject to adjustments during the various editorial stages of the bid book production.

Some figures in this document are presented as "draft" version. For the time being these should suffice to illustrate the geological description. Later these will be finalized and integrated in the application dossier as part of a follow-up project.

Application Dossier 1. Letter of Intrest 2. Application Dossier (in english). 2.A General information. 2.B Document checklist. 2.C Location of the area 2.D Main geological highlight and other elements. 2.E Verification of UNESCo Global Geoparks Criteria. 2.F Intrest and Arguments for becoming a UNESCO Global Geopark 3. List of Mandatory Annexes 3.1 Self-evaluation document. 3.2 An additional and separate copy of Section E1.1 "Geological heritage and Conservation". 3.3 An explicit endorsement of any relevant local and regional authorities and a letter of support from the National Commission for UNESCO or the government body in chrge of relations with UNESCO. 3.4 A large scale of map. 3.5 One-page geological and geographic summary. 3.6 Complete bilbliography of the area in Earth Sciecnes highlighting international publications.

Figure 1 – Guidelines for preparation of the AUGGp's application dossier. The highlighted items are (in part) presented in this report.

) TNO Public 4/219

2 Name and identity of the aUGGp

Sustainable Harmony: of Faults, Humans, and Rivers

The Geopark Peelhorst and Maasvallei (hereinafter aUGGp) encompasses a unique area that stands out by the intrinsic relationship between faults, waters, rivers, and men, ever since the first inhabitance of the area, several millennia ago. The area connects the provinces of Limburg and North Brabant through a complex fault system that extends from the surface to several kilometres deep. The area considered is within the influence of the Peelrand Fault and the Tegelen Fault and is named after 1) a region uplifted and tilted by fault activity, called the **Peelhorst**, and 2) the **Maasvallei** that crosses the Peelhorst and has been shaped by the continuously changing course of the Meuse river that is closely tied to tectonic movements (Figure 2). Although there have rarely been large and disastrous earthquakes, the aUGGp exhibits many phenomena linked to the presence and activity of faults. These faults, along with the horsts and grabens they delimit, have significantly shaped the landscape.




Figure 2 – DRAFT: Artist impression of the geological elements that shape the aUGGP Peelhorst & Maasvallei (shaded area)

The Geopark features several **geological elements** with rare hydrological **phenomena** and significant ecological diversity that are directly associated with faults and fault movements. Often linked to the hydrological situation and the different elevations and soil types are diverse habitats with diverse **flora and fauna**, including special vegetation types and various species of birds, mammals, reptiles, amphibians, and butterflies. This **ecological diversity and the unicity of the ecosystems** sets the Geopark apart from other tectonically elevated areas in the Lower Rhine Graben.

Both geology, (geo)hydrology, peat formation and ecology have led to a unique relationship between humans and nature, which is exemplified by a varied cultural history including a long period of adaptation, a period of landscape exhaust that culminated during the agricultural industrialization and a recent period of sustainable restoration of the landscapes. The

) TNO Public 5/219

geological elements of the aUGGp contain numerous **geosites** that tell the stories of controlling geological forces and their ever-changing interaction with humans, showcasing the importance of mankind as geological factor in the natural environment as well. The aspiring Geopark should raise visitors' awareness of geology and its relationship with nature and culture and invites them to 'explore and experience" the various **manifestations** related to this relationship. The Geopark Peelhorst and Maasvallei will do so by communicating five inspiring **storylines** that make the geology of the faults, the ensuing unique landscapes and the cultural adaptation and reclamation visible and tangible.

In recent decades, scientific research focused on locating and characterizing the faults and their multivariate influence. Since then, governments and various area partners, residents, businesses, along with researchers, were able to enhance the visibility of the faults. Many hiking and cycling routes have been set out, and information panels have been placed. To make the scientific research accessible, a landscape biography and a practical field guide have been published.

The story of the geopark inspires schools, nature education centers, and organizations such as IVN and local history circles to further disseminate the articulated knowledge. It encourages the tourism sector to offer an interesting stay in Brabant and Limburg, and agricultural entrepreneurs to explore other revenue models and the development of regional products. It will also serve as a solid foundation for the (to be developed) environmental vision and supports the uniqueness of the area. For years, initiatives have been developed by governments, residents, businesses, nature organizations, etc. to further develop the identity of the region as a whole.

Overall, the **Peelhorst & Maasvallei** aspiring geopark has a unique combination of geological, hydrological, ecological, and cultural-historical features that make it outstandingly significant among other tectonically active horst and graben structures in Europe and forms a valuable area for scientific study and conservation. It provides the best way to experience faults by exploring their many spheres of influence. Making the invisible visible, now and in the future.

) TNO Public 6/219

3 Location and delineation

The aspiring Geopark includes both the Peelhorst and the Maasvallei and is outlined by the Peel Boundary Fault Zone and the River Meuse (Maas) which are intersecting at both the northwestern and southeastern extremities of the aUGGp (Figure 3). The aspiring Geopark is situated in the European Cenozoic Rift System (ECRIS), a large horst and graben system that runs across Western Europe. This Graben system includes the Lower Rhine Graben of which the Roer Valley Rift System (RVRS) in its northwestern reaches is the main tectonic feature. Within the RVRS, the Peelhorst, also known as the Peel Block, is a geologically significant horst located in the southeastern part of the Netherlands, including the eastern part of the provinces of Brabant and the northern part of Limburg. To the west it is bordered by the active Peel Boundary Fault Zone (Peelrand breukzone) that marks the transition to the Roer Valley Graben and is part of the aUGGp as well (Figure 2). The Tegelen Fault delineates the Peelhorst to the east and marks the transition to the Venlo Block (graben). The River Meuse transects the Peelhorst at a right angle and formed the "Maasvallei." North of where it crosses the Tegelen Fault, the river changes its coarse parallel to the trend of the Venlo Block and gradually shifted northwards. Both the river incision and its northward shift are strongly influenced by tectonic block uplift and -tilting.




Figure 3 – A) Location of the aUGGP Peelhorst & Maasvallei (green droplet) in relation to other (109) UNESCO Global Geoparks in 28 European countries (source: https://www.europeangeoparks.org). B) Position of the aUGGP within the European Cenozoic Rift System (ECRIS) and relative to nearby Geoparks in the Netherlands, Belgium, and Germany. 2 = Vulkaneifel Geopark; 5 = 51 = De Hondsrug Geopark, 68 = Famenne-Ardenne Geopark (BE), 91 = Geopark Mellerdall; 105 = Schelde Delta Geopark (NL-BE).

) TNO Public 7/219

3.1 Relationship with surrounding Geoparks

To date, two UNESCO Global Geoparks are present within the Netherlands and one within the bordering country Belgium, i.e. Scheldedelta UGGp, Hondsrug UGGp, and Famenne Ardenne UGGp (Figure 1). The latter, however, is more than 100 km apart

Geopark Scheldedelta in the southwest of the Netherlands is a Netherlands-Belgium crossborder geopark that represents a dynamic estuarine landscape that, over 50 million years, was shaped by subsidence, uplift, sea-level changes, and tidal forces that resulted in sediment-starved lowlands and preserved tidal landscapes. The unicity of the UGGP lies in the more recent interaction between these natural processes, climate change, coastal dynamics, and human impact as a geological force (dikes, waterworks, land reclamation).

Geopark Hondsrug it themed around Ice-age geology and cultural history, with a strong focus on glacial geomorphology and human adaptation. The geopark represents a glacial landscape formed during the Saalian glaciation (~150,000 years ago). It features linear till ridges and ice-age landforms, such as megaflutes, pingo remnants, erratics (boulders from Scandinavia). The UGGp embodies rich archaeological heritage (e.g., hunebeds, prehistoric settlements).

The Famenne-Ardenne Geopark focuses on karst geology, paleoenvironments, and biodiversity, with strong ties to scientific research and tourism. The geopark is characterized by karst landscapes in the Calestienne limestone belt and contains geological formations dating back to the Paleozoic era. The area boasts an exceptional geological heritage, with more karst phenomena (caves, swallow holes, underground rivers) to admire than anywhere else in Europe.

The aUGGp Peelhorst & Maasvallei is themed around tectonic activity, paleoseismology, and fault-induced hydrology. The area is formed by a tectonic fault zone (Peel Boundary Fault Zone) separating the elevated Peelhorst from the subsiding Roer Valley Graben. It is an active fault system with ongoing vertical movement (~5 cm per 1000 years), historical earthquakes (e.g., Roermond 1992) and related tilting of the Peelhorst that significantly influenced the course of the River Meuse. The uniquely manifested geological and hydrological features include "wijstgronden" (iron-rich seepage zones caused by groundwater forced to the surface due to impermeable fault zones), bog iron banks and rust-coloured water streams.

A comparison of the main geological themes and features of the aUGGp's and surrounding UGGp's (presented in Table 1) learns that each of them exhibits unique geological features and that the cultural history, related to geology and landscape evolution, consequently is different. The evaluation highlights the Peelhorst & Maasvallei aUGGP as possessing a notably high degree of uniqueness.

) TNO Public 8/219

Table 1 - Result of a comparison of the aUUGP with surrounding UGGP's $\,$

	Doolhorst 9 Manualloi alloco	Scholdodola III GD	In ademia III Co	Comound Ardonno 11000
Geological landscape	• Faults & Rivers	Stuarine & Coastal	• Glacial	• Karst
Rivers	• Maas	Schelde River, Streams developed perpendicular to the escarpment are stream valleys and sand drift areas	• Drentse Aa, Hunze	 Lesse, Lomme, Ourthe rivers formed incised valleys perpendicular to tectonic ridges
Faults	Peelrandbreuk	n.a.	n.a.	Variscan faults and folds
Morphology	Contrast in elevation between uplified Peethorst and Roer Valley Graben across the Peet Boundary Fauttone. River terraces where noter Mass transects uplifted Peethorst	Contrast in elevation between estuarine and fluvial landscapes across the Brabantse Wal escarpment		 Atternating bands of limestone ridges and shale depressions (Appalachian-style relief)
Hydrology	• Wijst (iron-rich seapage)	 Mixing of salt and freshwater in tidal estuary ebb and flow 	Stream valleys caved by methwater from retreating glaciers Water-filled pingo remnants Wet/dry zones related to clay and sand reidged	Karstic hydrology Groundwater emergence at boundary time aquifers and impermeable shales
Biodiversity	• Hoogveen, wist (vet) • Stuffzand (dry)	Wetlands, salt marshes, peat meadows, intertidal zones, and floodplains, strongly influenced by the transition between saltwater and freshwater environments	Related to hydrological contrast in stream valleys	Limstone graslands (grazing and haymaking)
Climate control	 Meuse river behavior and course Drought (present) 	• Rising sea level (present)	Glaciations (past) Droughts (present) Biodiversity shifts (present)	Kloods (present)
Human impact	Peat extraction and landreclamation affected hydrology significantly. Former hydrological situation are restored at multiple locations	• Sediment management • Dykes • Changing estuary morphology	• Drainage	• Landscape attering through quarrying
Human adaptation	 Settlement locations Agriculture adapted to the reversed hydrology 	 Adaptive water management Resilient infrastructure. 	• Settlement	• Settlement
Land reclamation	 Ditches, canals, and pumping stations used to regulate water levels and reclaim land for peat extraction, drainage, and agricultural development. 	• Drainage • Urban development	• Drainage • Cultivation	 Not a dominant theme; some natural depressions and valleys were reclaimed for productive use
Earliest inhabitants	 The edges of the Peelhorst were first inhabited during the late Bronze Age and early Iron Age (1500 BC) 	 Early hunter-gatherers and farmers settled in the delta form the Mesolithicum (-9500 BC)onward 	 Neanderthal hunters settled in the region over 120,000 years ago, shortly after the formation of the Hondsrug glacial ridge 	Paleolithicum (-35,000 BC)
Economics	Peat extraction Inon bog Sand, gravel exploitation	Historic towns like Antwerp, Bergen op Zoom, and Reinnerswaal became centers of trade and culture due to their strategic locations along waterways frishery Townstran - Tourism	• Peat extraction	• Limestone exploitation
Tourism & Heritage	Walking and cycling routes with fault and wijst themes Watermills Wijst areas Historical settlements Od agricultural zones History of peat extraction River valley nature parks	Historic constructions (medieval dykes and sluices) Historic towns Traditional polder landscapes	Hunebeds (prehistoric tombs) Medieval Carl Tracks Historic villages and churches Walking and cycling routes across ridges and valleys.	Gaves attracting hundreds of thousands of visitors annually Beer

Some topical features occur in other Geoparks as well, though in a different geological setting

Unique to the aUGGP

) TNO Public 9/219

4 Surface area

The aUGGp Peelhortst & Maasvallei covers an area of 1.600 km², comprises 14 municipalities and counts more than 500.000 inhabitants. Spread over the area of the aspiring Geopark are 40 identified sites of geological scientific interest (Figure 4).




Figure 4 – The delineation of the aUGGP Peelhorst & Maasvallei with its geosites.

) TNO Public 10/219

5 Physical and geographical characteristics

The Peelhorst & Maasvallei encompasses an area with an extensive range of manifestations, such as terrain steps, significant differences in hydrology, shifting rivers, and large inaccessible peatlands that all have their bearing on unique flora and fauna and cultural development. For centuries, mankind adapted to this landscape, but the connection of all these phenomena was only poorly understood until 100 years ago. Coal exploration in the Peel during the last century made the subsurface more comprehensible through deep borehole- and later also seismic data. Although the search for coal did not lead to actual exploitation, it made a significant contribution to the geological knowledge of the subsurface. For instance, numerous smaller faults were found in addition to the major faults. Researchers still gratefully use the data collected at that time but also stress that more research is still needed to locate underexplored faults and to elucidate their effect on landscape development as that is not yet fully understood. The main potential of the aUGGp lies in its further utilization as freshwater reservoir during times of climate change, ensuring sufficient water for drinking, agriculture, and nature, making the application of geological knowledge a socially relevant issue for the 21st century and beyond.

5.1 Geological Elements

The Peel Boundary Fault Zone and the Tegelen Fault are the key geological elements of the aUGGp that, through their activity, demonstrate that the geological history of Europe is not limited to large mountain complexes like the Alps, Ardennes, or Eifel. The latter stand out through by presence of all types of hard rocks and prominent tectonics feature that can be observed directly in the field. In the relatively flat Netherlands, the thick pile of unconsolidated sediments (soft rocks) that was deposited over the last ~50 Myr in the delta toward the North Sea, turns out to be fragmented into smaller fault blocks delineated by main faults such as the Peelboundary and the Tegelen Fault zones. These faults, together with the intervening fault blocks and the interactions with the Meuse river, make up the five **geological elements** within the aUGGp, each of which is characterized by a unique combination of geological, hydrological, ecological, and cultural historical phenomena.

5.1.1 Peel Boundary Fault Zone (1)

The Peel Boundary Fault zone is an active fault zone that separates the higher-lying Peelhorst from the sinking Roer Valley Graben. As with many other faults, it is not a single straight line but rather represents a zone with fault segments that at some places merge and at other diverge, leaving an anastomosing fault pattern. Although less visible in the landscape, many of the surficial faults (indicated in red in Figure 18) merge at depths greater than 1 km, resulting in flower-like structures in cross-sectional view. Considering all this, it is difficult to delineate a single Peel Boundary Fault and it is better to refer to it as a zone. The fault branches that are significant as far as manifestation are concerned, are named individually (for example the Milheeze fault), but still belong to the same complex fault zone.

) TNO Public 11/219




Figure 5 – DRAFT: The five geological element that are the building blocks of the aUGGp. The striped parts of the elements fall outside the aUGGP area. Geological element 6 (Meinweg) is not part of the aUGGP, which is motivated below.

) TNO Public 12/219

5.1.2 Peelhorst (2)

The Peelhorst is a prominent **geological element** in the aUPPg, characterized by its elevated position relative to the surrounding areas. It is bordered by the Peel Boundary Fault (Peelrandbreuk) and the Tegelen Fault. The faults mark the transition to the subsiding and lower-lying Roer Valley Graben and Venlo Block in the west and east, respectively. The Peelhorst features a unique landscape which is distinct from lower-lying rift grabens and is characterized by a variety of hydrogeologic manifestations and ecological habitats. On the Peelhorst, fine and impermeable, shallow marine, and nearshore sediments from the Miocene (Breda Subgroup) are directly overlain by coarse Maas sediments deposited in a time the river occupied a more westerly position. Because the Peelhorst was continuously uplifted, the Miocene deposist are now close to the surface and form a natural barrier for water to penetrate. Consequently, the Peelhorst turned into a "bathtub" that is typified by poor drainage and high groundwater levels. Conditions that around 4000 BC eventually led to the formation of vast peatlands that used to cover large part of the Peelhorst. The diverse hydrological conditions support a wide range of flora and fauna. The region is home to various species of birds, mammals, reptiles, amphibians, and butterflies, making it an important ecological area. Efforts are ongoing to restore the original peatland ecosystems in the Deurnese & Mariapeel and the Groote Peel. Both are Ramsar international Wetlands and European Natura 2000-regions because of the unique interactions of (geo)hydrological system and the ensuing biosphere.

5.1.3 Tegelen Fault (3)

The Tegelen Fault delimits the Peelhorst to the east. Although terrain steps are not observed along this fault, it is considered to be an active and important fault for several reasons. Foremost, the Tegelen Fault plays a significant role in blocking the transport of water towards the east, i.e. east of the watershed on the Peelhorst. This is exemplified by many spots of ironrich seepages and numerous streams that run towards the Maas and emerge in the vicinity of the faults. Natural earthquakes have not been prominently associated with the Tegelen Fault. Recent seismic activity linked to this fault was induced by human activities. The felt earthquake of magnitude 1.7 near the Californië geothermal field in 2018 was a result of geothermal operations, specifically the injection of cold water into the fault zone. Additionally, various hiking trails have been established along the Tegelen Fault, such as those offered by the Breukenland Web-App. This app provides an adventurous journey along the Peelrand Fault and Tegelen Fault, where you can listen to fascinating stories about earthquakes, underground waterfalls, and other geological phenomena along the way.

5.1.4 Maas Valley (4)

Whereas the Meuse flows over the Campine Block, the Roer Valley Graben, and the Peelhorst in an orientation more or less perpendicular to the orientation of the tectonic blocks, this changes east of the Tegelen Fault. Here, the Meuse shifts in a north-westerly direction, more or less parallel to the orientation of the Venlo Block, an area that is delineated by the Tegelen fault to the west and the Vierssen Fault to the east.

In this geological element, the river Meuse shows another way of adapting to tectonic influence. Since the last ice age, the Meuse has shifted in a northeasterly direction on the Venlo Block, which is evident in the preferential preservation of river terraces on the western bank of the Meuse. This shift is the result of the tilting of the Venlo Block in a northeasterly direction, a phenomenon that also occurs with the Peelhorst. During the Holocene, there have

) TNO Public 13/219

been an above-average number of shifts in the river channels of the Rhine and Meuse near the Peel Boundary and Tegelen Fault Zones (*Stouthamer and Berendsen, 2001*). Stream valleys developed especially after the Younger Dryas and are representative of the tilt-induced drainage system that sources at the Tegelen Fault and stream into the Maas. The Maas Valley nicely illustrates that faults have great influence on the direction of stream courses. The longer segments of the past and present river courses parallel the main faults, whereas meanders occur as overstep zones from one fault to another.

5.1.5 Maas terraces (5)

Approximately 300,000 years ago, the River Meuse was still located south of the Peel Boundary Fault, in the Roer Valley Graben. Due to lateral displacement towards the northeast, it occupied its current location in the Venlo Slenk for the last 150,000 years. The Maas terraces developed when the River Meuse started its northern course and cut through the uplifting Peelhorst, locally forming up to 15 m high steep cliffs. Consequently, Meuse deposits are found in the subsurface of the entire Geopark. The intersection of the Maas terraces and the Peel Boundary Fault nicely illustrates how faults in soft and unconsolidated sediments have great influence on the direction of streams that, counter intuitively, flow against the northerly current direction of the Meuse. Overall, this reach-to-reach spatial variance in river terrace preservation and morphology can be ascribed to tectonically driven variations in river gradient and subsurface lithology, and to river-driven throughput of sediment supply.

) TNO Public 14/219

De Meinweg (6) and Eastern Maas terraces - A motivation for exclusion

For the scientific substantiation for the application of UNESCO status, 40 characteristic places (geosites) within the aUGGP must be designated. As this is a relatively limited number, for the selection, a balance is sought between sites of international, national, or regional significance. Furthermore, the uniqueness of a geosite is considered so that the aUGGP has as much variation in manifestations and experience as possible. Both within and outside the aUGGP, there are countless "parels" (gems) that also fall within the theme of the geopark and are well accessible to tourists and recreational visitors and often serve educational purposes. For example, there are about 20 beautiful wijst areas in the Geopark area. Not all these sites are included in the application but are well documented and also included on the gems map (which is a living document). The aUUGP "Peelhorst & Maasvallei" tells the story of the interaction of faults, rivers, and humans and is delineated and defined by the PBFZ and the Maas. By using these two geologically important boundaries, some valuable areas fall outside the geopark. These are the Meinweg area (municipality of Roerdalen) and the eastern Maas bank (municipalities of Beesel and Swalmen).

<u>Meinweg</u>

National Park De Meinweg lies east of Roermond and is part of the Limburg Maas terraces with a unique terrace landscape. The Meinweg area extends far into Germany (up to Wassenberg), and the border with Germany does not form a geological boundary. The terraces in the area are connected by three steep transitions with a total height difference of 50 meters. These terraces mainly represent high terraces formed by the Rhine, in conjunction with the river Roer and displacement of the PBFZ.



Figure 6 – Landscape form east of the Maas and across the border with Germany.

Perpendicular to these terraces run several brook valleys and rivers. In between are ponds, extensive forests and heathlands, estates, and a varied agricultural landscape. The Limburg Maas terraces not only have rich nature but also a fascinating cultural history. Agriculture has played an important role in this from an early stage. The area has been used communally by surrounding villages for centuries. Today, it is a place where recreation, agriculture, habitation, and nature management go hand in hand.

) TNO Public 15/219

Nevertheless, the Meinweg area is not included in the aUGGP for the following reasons (listed in decreasing order of relevance):

- 1. The geological story is more complex than that of Peelhorst and Maasvallei alone.
- 2. The area extends far into Germany.
- 3. Many of the manifestations related to geology are not unique compared to the aUGGP.
- 4. There are currently no national or international participation partners within the Meinweg area.

Eastern Meuse terraces

The part of the eastern Meuse terraces that is part of the municipality of Roermond has a special event in recent history, namely the Roermond earthquake with the epicentre at Herkenbosch. It is worth considering adding this area to the aUGGP because here the direct tectonic influence of the faults on humans was felt and seen. On the other hand, the event is mainly anecdotal, and much stronger earthquakes regularly occur worldwide. Although the earthquake was also felt within the geopark area, the story of the geopark is not so much about earthquakes but rather about faults and their influence on the landscape and how humans deal with them. Adding this area to the base area makes the Geopark story less coherent and also difficult to frame geologically.

The part of the eastern Maas bank that is part of the municipalities of Beesel and Swalmen is characterized by steep terraces and brooks that drain into the Meuse. There are also some fault-related manifestations in this area. At Turfbroek in Asenray (Limburg), the location of the Peel Boundary Fault can be determined quite accurately based on the boundary between green and yellow grass. The presence of the fault also affects the roof structure of a farm there. Despite repeated repairs, the structure continues to crack and subside.

Boukoul is a good example of a castle farm on the PBFZ with a moat fed by seepage water. The Natura-2000 area Swalmdal is located around the Swalm, a meandering brook on the eastern Maas bank that is deeply incised into the Maas terraces landscape. The brook lies at the transition from the plateau between Maas and Rhine to the Maas valley, this transition roughly coincides with the Netherlands-Germany border. At various places at the foot of the terraces, seepage occurs, and springs emerge.

The eastern Maas bank is not included in the aUGGP for the following reasons (listed in decreasing order of relevance):

- 1. The earthquake story is mainly anecdotal and can also be told/demonstrated within the aUGGP itself.
- 2. Adding this area makes the Geopark story less coherent and also difficult to frame geologically, as the Peelrand fault extends much further into Germany.
- 3. There are currently no national or international participation partners from the municipalities of Roermond, Beesel, and Swalmen.

) TNO Public 16/219

5.2 Storylines

Each of the **geological elements** is characterized by a range of **manifestations** in morphology, soil and/or water conditions, the associated ecology (flora & fauna) and the human interaction that make the long geological history of the area tangible. Six inspiring **storylines** exist at the intersections of the **geological elements** and **manifestations**. The storylines (Figure 7) represent a set of Unique Selling Points compared to UNESCO Global Geoparks worldwide and describe the dynamic, living fault landscape shaped by tectonic activity and its influence on the freshwater system (rivers, streams, groundwater). **Humans** have always played a significant role, either through adaptation or by actively utilizing the opportunities provided by the **geological elements**, thereby also changing them over time. The human impact is visible throughout cultural history and can, at times, be considered a geological factor in its own right.




Figure 7 – DRAFT: Storylines of the aspiring Unesco Geopark Peelhorst & Maasvallei

5.2.1 Active faults

The Peelhorst is characterized by significant fault movements along active fault zones that record historic earthquake activity and have geomorphological scarp expressions that remained, despite erosion and human interference. The resulting **terrain steps**, to date, are still visible at many locations and are usually around 1 m in height. Very rarely, fault displacements are accompanied by light to moderate earthquakes. The terrain steps, together with information about the displaced sediment units at the surface (from trench studies) or below the surface (from boreholes and geophysical surveying such as seismic and ground-penetrating radar studies), and hydrological phenomena such as "wijst," allow to trace the faults through the landscape. The most up-to-date fault pattern is a results of connecting the many "dots" of information, rather than being visible and traceable over lengths of kilometres.

Terrain steps: The total elevation difference at faults grew fairly gradual over the millions years of fault activity. At the fault lines, though, clear terrain steps are visible in several places. The height of these terrain steps varies. In some places, the terrain step is clearly visible. Terrain steps are especially visible where a paved road crosses a fault line. Examples include Scheiweg and Haveltweg (Handel fault), Daalhorst (Gemert-Zuid fault), and Esp (Peelrand

) TNO Public 17/219

fault). If the terrain steps at all the faults are combined, the height difference between the lowest point in the municipality of Gemert-Bakel (12 m above sea level at Esdonk) and the highest point at De Bult in Milheeze (30 m above sea level) is a remarkable 18 m. For reference: the height difference between the North Sea and the municipal boundary is smaller. Terrain steps are most prominent at the 6 active fault lines, but also exist at some inactive fault lines. Due to solifluction phenomena during the ice ages and human interference, many of the original terrain steps may have dissapeared.

Earthquakes: The most famous recent earthquake in the Netherlands is the Roermond earthquake of 1992 (magnitude 5.8), which occurred at a time the Uden earthquake of 1932 (magnitude 5.0) was almost forgotten. The Roermond earthquake remains the strongest earthquake ever recorded in Europe north of the Alps. Therefore, scientists claimed that the Peel Boundary Fault is one of the most important, still active faults in the Lower Rhine region (Ahorner, 1994). He also stated that it is an extraordinarily active fault structure by Central European standards, which undoubtedly must be considered a potential earthquake source. This Roermond earthquake was a significant stimulus for renewed research into fossil earthquakes in the Roer Valley Graben system. Other (historical and prehistoric) earthquakes find their evidence in research trenches over faults that are visible in the landscape as terrain steps.

These trenching studies (e.g. *Van den Berg et al., 2002*; *Van Balen et al., 2019; 2024*) have shown that three to four paleo-earthquakes occurred in the northwestern part of the Peel Boundary Fault Zone. Two of these events took place during the Late Pleniglacial to Late Glacial transition, around 15,000 and 14,000 years ago, with estimated magnitudes of approximately Mw 7. These earthquake were at least 10 times stronger than the Roermond earthquake and involved a fault segment of approximately 35 km. The timing of both events suggest that glacio-isostatic adjustment is the triggering mechanism. A younger earthquake occurred in the Holocene, between approximately 10,000 BP and 800 BP. The recurrence interval of periods with major, surface-rupturing earthquakes is therefore estimated at 10,000 to 15,000 years, depending on location and type of fault activity.


Fluidization of sediment during earthquakes is a well-known phenomenon, caused by ground shaking. This has also been observed in Herkenbosch during the Roermond earthquake of 1992. The presence of fluidized sediment along the fault structure is further evidence that the abrupt fault displacement was the result of an earthquake, rather than a gradual, creeping process.

The most recent recorded earthquake along the Peel Boundary Fault occurred on 9 October 2020 near Sint Odiliënberg, south of Roermond (magnitude 1.2; hypocenter depth 15 km). Just three weeks earlier, two minor quakes (M 1.1 and 2.0) were also recorded in Sint Odiliënberg within the span of an hour (source: https://www.knmi.nl/nederland-nu/seismologie/aardbevingen)

Hydrology: The elevation change across the faults is often accompanied by an abrupt change in the phreatic groundwater levels of up to 1.0 to 2.5 meters due to the impermeability of the fault. This causes iron-rich groundwater seepage at the upthrown side of the fault and the formation of bog iron banks close the fault. The surface water in the surrounding ditches and trenches is often bright orange. Such discoloration is characteristic of wijst, i.e. iron-rich groundwater emerging at fault zones. These hydrological conditions at the fault zones support rare ecological phenomena that are unique in Europe and have been recognized as an internationally significant geological monument.

Due to the seepage and the emergence of numerous streams on the upthrown sides of the fault (the footwall), the downthrown sides of the faultzones (the hanging wall) have been an attractive place for settlement ever since prehistoric times. At present the fault zones are focal points for educational and recreational activities. Various walking and cycling routes, like the

) TNO Public 18/219



"Breuken Beleven" (Fault Experience) route, allow people to explore and learn about the fault's hydrological and ecological features.

Figure 8 – Photograph and drawing of the east wall of the trench across the Peer Boundary Fault near Bakel, showing the fluidized clayey sediment along the fault plane (modified from Van Balen et al. 2019).

An important observation from the trench studies at Bakel (Van Balen et al., 2019) and Uden, Lapperre et al., 2022) is that clayey sediment is the presence of clayey sediment within the fault structure itself (Figure 8), occurring both along the main fault branch and along several smaller branches. Structures within this sediment indicate that it must have reached its current position from deeper layers while still in a liquid state. This is a significant indication that the abrupt fault displacement is the result of an earthquake and not a gradual, creeping process. Moreover, this clayey sediment in the fault plays a key role in water management, as it enhances horizontal impermeability and thus contributes to the marked groundwater jump around the faults in the southern Netherlands.

5.2.2 Changing Rivers and Streams

Processes in the Earth's crust have had many consequences for the course of rivers in the Geopark area. The Peel Boundary Fault zone has played an exceptional role in this, as relatively large and abrupt shifts have occurred in this narrow fault zone. The Meuse river has had to adapt to these changes in the Earth's crust and landscape. By geological timescale standards, these developments have taken place quite recently. Consequently, research into the Roer Valley Graben System and the Peel Boundary Fault zone has provided much information about the interaction between fault movements, climate change (glaciations) and river dynamics. Much of this interaction is still visible today as elevated Maas terraces, abandoned Maas meanders, and river and stream valleys that flow upstream relative to the main drainage network (Figure 9).

) TNO Public 19/219




Figure 9 - Map of the Lower Meuse Valley showing the Weichselian and Holocene terraces and stream valleys. The present-day channel of the Meuse is represented by 0 yr BP. Modified from Woolderink et al., 2020 and Woolderink, 2021.

) TNO Public 20/219

Terraces: As the Ardennes and the Rhenish Massif rose, the river terraces of the Rhine and Maas in Limburg, Belgium, and North Rhine-Westphalia in Germany began to form around 3 million years ago. The terraces were formed as the rivers cut deeper into the landscape, leaving a terrace as a remnant of the old river plain high in the landscape. Due to the continuous subsidence of the Roer Valley Graben system, a substantial sediment package accumulated here, especially in the Roer Valley Graben itself, where the Maas originally flowed. In this graben, and to a lesser extent in the Venlo Graben and on the northern edge of the Kempenhorst, the Rhine and Maas deposited sand and gravel. Around 450,000 years ago, the Rhine shifted to a more northerly position, and the Maas became the main river in the south of the Netherlands. After the Rhine, the Maas also gradually shifted to the northeast over time. This process began with the filling of the Roer Valley Graben, allowing the Maas to shift over the Peelhorst and eventually into the Venlo Graben. The tilting of the Peelhorst towards the northeast caused the Maas to incise into the Peelhorst whereas it lower reaches became gradually shifted eastward, leading to the formation of the northern Maas Valley. By changing the course of the river, the eastward tilt of the Peelhorst influenced local geography and hydrology and became a notable historical feature.

Shifts and tilts due to movements along the Peelrand and Tegelen fault zones have greatly influenced river dynamics. The Maas terraces from the last ice age have been displaced by up to 1.5 meters, and the river course has changed significantly at the fault zones (see Figure 9).

Meanders: The surface movements within the PH&MV area also had a significant impact on the lateral variability of sediment types at or near the subsurface, thereby influencing the course of the Rhine, Meuse, and Roer rivers. Due to the relatively rapid subsidence of the Roer Valley Graben, the Rhine, Meuse, and Roer deposited their coarse, gravelly sediment there for a long time. On the Peelhorst there is a much thinner layer of river sediments; older, fine-grained deposits, such as the Breda Formation, lie near the surface. Thus, sediments of different compositions and ages are found next to each other. This relatively rapid change in the composition of the subsurface around the Peel Boundary Fault has impacted the dynamics of the Meuse at the transition from the Roer Valley Graben to the Peelhorst.

In the Roer Valley Graben, the Meuse meanders through a fairly wide river plain with significant lateral movement. On the Peelhorst, there is a much narrower river plain with an almost straight channel, and the Maas has hardly moved laterally (Figure 9). A key factor in the change in the river's course could be the difference in resistance to erosion. The coarse post Weichselian sediments in the river terraces in the Roer Valley Graben, despite their large grain size, are easier to move due to the lack of cohesion. On the Peelhorst, finer sediments stick together much more, resulting in less material being moved and the river having less opportunity to meander. There is a second reason for the decrease in meandering: As the Peelhorst rises relative to the Roer Valley Graben and the Venlo Graben, the Meuse responds by cutting into the sediment. This leaves less energy for lateral erosion.

The significant impact of faults on river dynamics is also evident from an atypical meander near Roermond, which formed during the beginning of the Allerød (13,900-12,850 years ago). This "abnormal" meander is the result of vertical movements along the Peel Boundary Fault Zone caused by severe earthquakes at the beginning of the Late Glacial (14,700 – 11,700 years ago; Van Balen et al., 2019). As the Peelhorst rose approximately 1 meter relative to the Roer Valley Graben, the bed of the Meuse was displaced by the same amount. Such a sudden displacement of the riverbed has significant consequences for the flow in the river, which affects sedimentation in the river channel and thus also influences the sinuosity of the river. Once this (temporary) obstacle in the Meuse river channel was eroded away, the course of the Meuse could return to its "normal" dynamics.

) TNO Public 21/219

Streams: A distinctive feature of the Geopark is the vast number of sites where groundwater is pushed to the surface leading to the formation of wet areas and wijst areas. This phenomenon is clearly visible on the Peelhorst in the vicinity of its bounding faults but also at the Maas Terraces. These seepages source the many streams in the area that make use of the terrain steps created by either faults or fluvial incision processes or a combination of the two. Also, the seepage water provides a constant supply of mineral-rich water, contributing to a unique flora and fauna.

5.2.3 Wet & Peaty

The geological structure and history of the Peelhorst have strongly influenced water flow and groundwater levels. During the eastward shift of the Meuse, (relatively) coarse river sediments were deposited on the Peelhorst. Combined with the (relatively) shallow position of marine (poorly permeable) Miocene deposits on the Peelhorst, a unique groundwater system with high groundwater levels established.

In the originally sparsely vegetated landscape of the Peel at the end of the last ice age (Figure 10), the wind created sand dunes that blocked the meltwater valleys, especially at the eastern site of the Peelhorst. As water was unable to penetrate the poorly permeable subsurface, small lakes were formed (e.g. the "Kom van Griendtsveen") that became vegetated as the climate warmed. With this vegetation, peat formation began. Because the soil had been frozen for a long time, few nutrients had been leached out. This led to lush vegetation. The peat that forms in this nutrient-rich environment is called fen peat. Gradually, the vegetation became denser, and forest peat developed. The thickening peat layer made the nutrient-rich groundwater increasingly difficult for plants to access. This eventually ended the formation of fen peat. The high elevation of the Peel also contributed to this process. There seems to be a relationship with the maximum peat distribution and some of the internal faults on the Peelhorst that delineate elongate depressions (minigrabens) that were important for retaining the water and focused the thick accumulaton of peat (see Figure 10Figure 10).

While most fen peat vegetation began to die off during the mid-Holocene (~6000-4000 yrs ago), the environment became suitable for sphagnum moss. Sphagnum thrives in nutrient-poor conditions and retains rainwater well due to its fine branching. Sphagnum mosses secrete acid, which inhibits bacterial growth and thus the decomposition of plant material. With the arrival of sphagnum mosses and acidification, the formation of raised bog peat began. The sphagnum moss areas expanded from the original lakes over the surrounding dry lands. At the peak of peat growth, about 4,000 years ago (Subboreal period, Figure 10), raised bogs with diameters of kilometres formed on the Peelhorst, and the raised bog eventually reached an area of 30,000 hectares. At 1000 BC, the climate became colder and wetter again. These conditions were favourable for renewed expansion of the raised bog, even further than it had during the Subboreal period. This expansion was likely also encouraged by human activity. Since the Iron Age, and especially from the early Middle Ages, many forests were cleared. Numerous villages and hamlets around the Peel owe their names to the felling ("rooijen") of trees: Castenray, Venray, Tienray, Leveroy, Stramproy, Tungelroy, Wanroy, and Nistelrode all refer to forest clearing (*Joosten, 1991*).

) TNO Public 22/219




Figure 10 – Peat development in the geopark. At 9000 BC (shortly after the last glacial), no peat had developed; at 3850 the peat had is largest extent; At present, because of the exploitation of bog, only some peat remnants remain. Modified from Vos et al., 2020.

For a long time these wet peat areas were difficult to access but eventually **reclaimed** for the exploitation of bog. Within the aUGGP, the geosite Deurnese Peel and Mariapeel represents an unclaimed area where sphagnum moss remnants are still present. However, sphagnum moss cushions are no longer growing because the Peel has dried out due to intensive drainage and the original extremely nutrient-poor conditions no longer exist. As a result, the decomposition of the peat released nutrients into the groundwater. Additionally, the surrounding livestock farms contribute a lot of nitrogen to the atmosphere and groundwater, making the environment too nutrient-rich for sphagnum moss development.

Restoring wet nature pearls and wijst areas along the main faults along the Peelhorst has received a lot of attention from water boards, provinces, municipalities, and landowners. The past dry summers have shown that it is important to retain water on the high grounds of the Peelhorst. In various places in the large remaining peat areas with a relatively significant thickness, restoration management has successfully led to the development of raised bog vegetation. Dams and weirs are intended to ensure that rainwater remains in the Peel.

To prevent groundwater from seeping away too quickly, the surrounding agricultural businesses and the drinking water companies are allowed to extract limited amounts of groundwater.

) TNO Public 23/219

Agriculture must also become increasingly sparing with the emission of fertilizers. The measures seem to be having an effect. In the wet peat pits, sphagnum moss is expanding, and the formation of sphagnum peat has begun again. In a few decades, we will know whether the almost disappeared plant communities of the raised bog environment can develop again sustainably.

Through the efforts to restore the original peatland ecosystems in the Deurnsche & Mariapeel and the Groote Peel these areas are now Ramsar international Wetlands and European Natura 2000-regions showcasing the unique interactions of (geo)hydrological system and the ensuing biosphere.

5.2.4 Wijst

On a regional scale, the shallow groundwater and surface water in the geopark generally flows from higher areas such as the Peelhorst to the lower Maas or the Roer Valley Graben. Water also flows from the Limburg river terraces to the Maas. These regional flow directions are intersected by clusters of subsurface faults, such as the Peel Boundary Fault Zone and the Tegelen Fault. These faults have reduced permeability and hinder groundwater flow, affecting surface runoff via ditches, streams, and rivers. This characteristic 'barrier' effect of the underground faults leads to raised water levels at the higher side of the fault, sometimes reaching or even exceeding the ground surface. Across the fault, the water can flow freely again and infiltrate into the predominantly sandy soils of the Roer Valley Graben. This creates an abrupt groundwater jump at faults like the Peel Boundary Fault Zone, comparable to an underground waterfall (Figure 11). An inventory of the jump height has revealed that the groundwater jump is usually 1 to 2.5 meters, although both lower and higher jumps occur as well.

The faults thus form relatively sharp boundaries between a higher side that is often wet and marshy and a lower side that is much drier. This counterintuitive 'upside-down world' is a well-known phenomenon of the Brabant and Limburg faults. Those who live high up often suffer from wet feet, while those who live in lower areas remain dry. This is completely different in the rest of the Netherlands, and from a hydrological perspective, the situation along the Peelrand fault is unique.




Figure 11 - Schematic representation of the barrier effect of the Peelrand fault zone (in red), its impact on local groundwater flow, and the location of wijst areas fed by iron-rich seepage (modified from *Lapperre, 2025*). Note that similar phenomena occur along faults internal to the Peelhorst and along its eastern boundary (Tegelen Fault) as well.

) TNO Public 24/219

The water that surfaces at the higher side of the fault contains a lot of dissolved iron derived from glauconite-rich Miocene marine sediments that are relatively close the surface on the elevated Peelhorst. The iron is bound by iron bacteria at the bottom of ditches and in the air through reactions with oxygen, carbon dioxide, and water. The iron compounds give the seepage water a reddish-brown rust color and explain the remarkable coloration of the streams and ditches. The unique relation between of iron-rich seepage and associated faults is referred to as "wijst" and marshy areas as "wijstgronden" and hereafter referred to as wijst areas.

Locally, the iron oxide precipitates in the pore spaces of the coarse sediments close to the fault that, over time, led to cementation of the sediment and formation of **iron ore** banks (bog ore), further reducing permeability. Man has known about the high iron content of the groundwater and surface water in the Peel area for many centuries, and has profited from the bog ore by using it as a building material since Roman times. But the iron ore sometimes hindered the access to suitable drinking water for man and cattle.

The faults that obstruct the groundwater flow are usually difficult to see at the surface. However, if we look closely at specific indicators, the faults reveal themselves through the presence of elongated wijst areas, a jump or slope in the ground surface (a so-called terrain step), watercourses with a strong orange colour, ecological contrasts, and fauna differences in crop growth. The latter occur because plants develop less well in wet and cold soil than in drier and warmer soil. Also, faults and water-related cultural-historical objects are found in more places, such as the iron-containing Wonderbare Bron of Handel and significant references such as the Hooge Wijststraat (in Heesch) and the Roode Beek (Meinweg). When several of these characteristic phenomena occur simultaneously, it provides an additional indication to suspect underground faults. In this way, it is possible to indirectly see what is hidden underground.

Fauna & Flora: Where there is no terrain step, the location of the fault, if undisturbed, can sometimes still be recognized by the presence of seepage flora 'above' the fault, bordered by dry grasslands 'below' the fault. The iron-rich seepage water is typically nutrient-poor to moderately nutrient-rich, soft, and often rich in sulfate. The flora of wijst areas is typically rich in species that thrive in these conditions, such as tufted sedge, marsh ragwort, snakeweed, broad-leaved marsh orchid, and marsh marigold. In the past century, humans have largely drained the wijst areas and used them as grassland. Most of the originally marshy grounds are now characterized by a landscape with ditches, wet hay meadows, alder groves, and small swamp forests. Restoration of wijst, such as at the Sint Annabos area, led to a scenically beautiful alder carr forest on the high side of the fault. On the wet peat under the alders next to ferns and tufted sedges also the water violet grows. The latter is a plant with submerged leaves that occurs in shallow waters in shady areas and is a good indicator of seepage. Also in streams, as well as in pools and fens in the areas surrounding faults, plants typical of seepage areas grow, such as rushes and sedges or species of moist grasslands, such as field sorrel and cuckoo flower. The area around the fault offers birds a wide variety of habitats. Wet and dry, nutrient-rich and nutrient-poor, wooded and open areas lie close together. Among the many different bird species, the rare spotted crake is the most notable.

In most areas on the high side of the fault, the quality of the vegetation has come under pressure due to the increasing nutrient richness due to the rise of artificial fertilizers and intensive farming. Plants like nettle, hogweed, thistle, and reed appeared. This nutrient richness promotes the growth of willows, nettle, hogweed, thistle, and reeds at the expense of alders and tufted sedges. The wijst areas of the Sint Annabos, however, proved that the dissolved iron in the seepage water is able to bind phosphates, retaining the relatively nutrient poor conditions.

) TNO Public 25/219

5.2.5 The force of Water

The topography caused by fault activity influences the drainage system of streams, rivers, and groundwater. Throughout history, this has sometimes been used beneficially.

Starting from the Middle Ages, water mills and water wells were built in many places (*Bon, 1972*). Their location is often determined by the underground fault system. In places with a fault in the subsurface, the impermeable fault facilitates groundwater seepages that are often the source for streams (brooks) that can be used for the operation of water mills. The strong seepage near the faults provided a more or less constant water supply, while the natural elevation difference in the landscape created additional water-level gradient. The most suitable location for a water mill is on or slightly downstream where a stream crosses a fault. Along the Peelrand fault zone, only one watermill has been preserved: on De Vlier brook in Deurne at Haageind. However, research showed that once there were six more watermills. All known locations are, as expected, situated on the Peel Boundary fault or one of its secondary faults. At the eastern site of the Peelhorst, the water mill in Oploo is situated on the Tegelen fault. The watermill at Neer uses the natural fall of the Neerbeek stream into the Maas

In addition to watermills, the choice of location for castles and terrains surrounded by ditches, often referred to by the English term 'moated site', surprisingly also appears to be related to faults and springs. These castles and moated sites were built in the 13th, 14th, or 15th century as noble residences. Their once existence is partly derived from the presence of a moat. Such a moat was not supposed to dry up in summer and was therefore almost always situated along a brook. In the municipality of Gemert-Bakel, the locations of ten moated sites are known and it turns out that seven of them are located near one of the three existing fault branches of the Peel Boundary Fault Zone. Always on the side of the downthrown block and all within a distance of 100 meters from the fault.

The most intense manifestation of the spring phenomenon was at places that centuries ago were named 'put' (well). These wells were so prominent that everyone knew these places, and for that reason, they were designated as fixed boundary points between villages on the edge of the Peel in the Middle Ages. Just as sources of brooks and rivers often gained religious significance, this was also the case with the so-called Willibrord wells. Saint Willibrord is said to have baptized there. At the border of Deurne and Meijel, we find a Willibrord well, and also at the border of Oss and Heesch. Both wells are associated with a fault. In other places, the spring phenomenon was less intense but still strong enough to be the source of a natural brook. Based on the oldest topographical maps, the natural watercourses can be reconstructed. This has been done for Gemert; ten places have been found where a brook begins. It turns out that all natural watercourses there are sourced from a spring area. In places where a well was present or at the source of a brook, the spring phenomenon was so strong that the water and marshy character dominated, and people were more troubled by the spring than they benefited from it. The advantage of the spring was found in places where the spring was less intense.

5.2.6 The human factor and the landscape

The Peelhorst & Maasvallei aUGGp represents a diverse landscape that is influenced not only by geological forces but also by humans that reclaimed the originally wet and difficult-to-access area. In a country like the Netherlands, where water always shapes the environment, even on the higher sandy grounds, geological history is crucial for our living environment. It serves as a valuable tool in addressing one of this decade's challenges: managing the right amount of water for agriculture, nature, and drinking water

Both faults and rivers have a significant impact on the development of the landscape. This is especially true for the Peelhorst, where the Rhine, Maas, and Peelrand fault contributed to the

) TNO Public 26/219

formation of the landscape. The Peel, with its extensive, almost impenetrable peatlands for humans, and the Maas to the east and north, formed a border area for hundreds of years. With only a limited number of trade routes over sand ridges, the marshy Peel was well-defensible and an ideal natural border zone. For centuries, territorial boundaries lay here. For the local population, it was an area for peat extraction, and land was reclaimed on a small scale. The mysterious phenomenon of wijst gave rise to stories and legends about miraculous springs. Wijst water was practically used for agriculture, the construction of moats around castles, and the operation of water mills. But sometimes the wijst was so marshy that it simply couldn't be used for agricultural purposes. Especially at the eastern part of the fault zone, the wijst was most intense and disruptive. The unique geology made the area also interesting for the extraction of minerals. In the Peel, iron ore, sand, and gravel were extracted, and even coal was successfully drilled, but never extracted.

Iron ore: The water that surfaces in wijst areas contains a lot of dissolved iron. The iron is bound by iron bacteria at the bottom of ditches and in the air through reactions with oxygen, carbon dioxide, and water. The iron compounds give the wijst water a reddish-brown rust color and the term "rood in de sloot" (red in the ditch) is commonly used to describe the phenomenon. Cementation occurs, and iron ore is formed close to the fault with a variable and complex composition. Either the iron compounds cement the sand on the higher site of the fault as iron-rich wijst water is transported across the fault under influence of the faultinduced slope, or it cements coarse-grained fluvial deposits on the lowerl site of the fault. Both processes occur, explaining the wide distribution of iron ores. In the past, iron ore chunks were extracted for various purposes. The chunks of iron ore were used, among other things, as building material. They are regularly found in the foundations of churches in the region. They were also used for the piers on which the wooden beams of farms were founded. After bricks became affordable, this tradition disappeared. It is notable that iron ore chunks continued to be used in the construction of communal religious structures such as Marian grottos and mounds on which roadside crosses were placed. The ores were also extracted for iron production. Archaeological research has shown that iron production already existed in prehistoric times, based on the findings of charcoal kilns and iron slag (charcoal is produced in charcoal kilns, which is necessary for iron production). The extraction of iron ore remained important in historical times as well. Municipal reports from Deurne and Gemert in the 19th century mention the "export" of iron ore for iron production. The trade in iron ore was more prevalent on the eastern side of the Peelhorst. In the marsh between Wanroy and Rijkevoort, a lot of so-called marsh stone was found, and in 1857 the municipality sold a concession for its extraction. A horse-drawn tram line was constructed to transport the material from this marsh to a depot along the Maas near Beugen, where the iron ore could be shipped.

Peat: Between about 4000-1000 years ago the peat growth reached its greatest extent and from the early Middle Ages (500 – 800 AD) onward the Peelhorst area changed drastically because from then on, peat was extracted on an increasingly large scale. The first historical records of peat extraction in the Peel date from around 1400. The destruction of the last forests around the Peel likely forced people to look for alternative energy sources. Peat extraction was an obvious choice—not only because people in the Netherlands already had experience with it elsewhere, but also because the plaggen agriculture practiced on the sandy soils at the time had already introduced some of the techniques and tools needed for peat cutting (Joosten, 1995a). The wild land, where, it was said, wandering lights lured travelers into the swamp, transformed in about a century into an extensive, efficiently organized agricultural area. In the northern part of the Peelhorst, the raised bog had already significantly decreased by the 19th century. In the southern part, large-scale industrial peat extraction began at that time. Until then, the Peel was a large, almost impenetrable, marshy area, resulting in it being not only a natural border but also an administrative and, to some extent, a cultural boundary (Van den Brand, 1982; Iven and van Gerwen, 1978). Even in the 1930s, the Peel was so inaccessible that the government considered the area suitable as a natural

) TNO Public 27/219

barrier for the construction of the Peel-Raam Line, a defensive line against a German invasion before WWII. With a canal, casemates, obstacles, and the last remnants of peat, it was hoped in the late thirties to stop or slow down a German invasion close to the border.




Figure 12 – The situation in 1853, with the old villages around the Peelhorst are indicated. Map originally presented by *Van den Munckhof & Joosten, 1990*

Sand and Gravel

Due to the uplift of the Peel Horst and erosion of younger sediments, Pleistocene, Pliocene, and Miocene sand and gravel formations are relatively close to the surface in this area. In many places on the Peel Horst—such as Milheeze, Zeilberg, Liessel, and Neerkant—extraction has taken place in the past. Most of the sand was and still is extracted from either fluvial deposits of the Beegden and the Kreftenheye Formations which are rich in gravel and coarse sand, or marine deposits of the Breda Subgroup and Kiezeloolite Formation, which are rich in green- and quartz sand, respectively.

The coarse sand is characterized by angular grain shapes. Locally, this sand is referred to as "skèrrep sand" (sharp sand) and is ideal for use as construction material. For road builders, glauconite-bearing sand is a good material; glauconite grains are angular and interlock tightly, a property that makes the green sands particularly suitable for constructing motorway embankments. Sand and gravel extraction has ranged from very small-scale to large-scale commercial operations. The small-scale extraction mainly involved masonry sand and gravel, which was collected in small quantities 'in the wild' by farmers, civilians, and contractors at

) TNO Public 28/219

numerous locations. Commercial sand extraction took place in two major concessions on either side of the Peel Horst. At the western side in Hoogdonk near Liessel and on the eastern side at the "De Kuilen" near Langenboom (Figure 13).




Figure 13 – Southeast-northwest cross section through the subsurface model Geotop (v1.6.1) across De Kuilen sand extraction site situated on an elevated fault block on the Peelhorst where the Miocene is closer to the surface. See inset map for location.

In 1954, the Kalkzandsteen (lime sandstone) factory Hoogdonk was established in Liessel. The Peel Boundary Fault runs directly in front of the factory gates. On the factory grounds, quartz sand of the Kiezeloolite Formation are close to the surface, making it easy to extract. These "silver sands" consists mainly of fine to medium-grained quartz grains that are well sorted. This pure sand is ideal for binding with lime to produce lime sandstone bricks. In addition to silver sand, deeper drillings in Hoogdonk have revealed greenish marine sands from the Groote Heide Formation at depths of 50 to 136 metres. These older sands are not typically used for lime sandstone production but are of geological interest. In 2000, a permit was granted to expand the extraction site, allowing the factory to continue production until around 2030. However, sand extraction was halted in 2022, although Xella Kalkzandsteen resumed sand extraction for lime sandstone products at a different location in 2024. The excavation permit is valid until the end of 2025; further plans remain unknown.

De Kuilen near Langenboom are water bodies formed by the extraction of "green sand." On the Peel Horst, green sands from Miocene age lie less than 1 meter below the surface, whereas elsewhere in the Netherlands they are found at much greater depths. The sand is extracted using a suction dredger that draws the sand from the bottom of the lake.

Both in Hoogdonk and De Kuilen, sand extraction (down to about 30 metres deep) has led to the discovery of numerous fossils, particularly from the late Pliocene and Miocene. The two sand extraction sites are paleontological mirror images on either side of the Peel Horst and are of exceptional national importance.

In Hoogdonk, fossil finds include shark teeth, whale vertebrae, bird remains, sea urchins, and starfish. Whale remains and many other marine fossils provide evidence that this part of the Netherlands was once a sea (Miocene, Breda Subgroup). Remains of mastodons and other fossils date from a time when the area was land (Pliocene and Pleistocene, respectively Kieseloolite and Beegden formations). Occasionally, large erratics ("vlincken" in Limburg an Noord-Brabant) are found in the Pleistocene river deposits (Beegden Formation) representing

) TNO Public 29/219

large blocks and stones that were transported by floating ice via the River Meuse.. In Liessel, for example, the "Jaap van de Leijsing" and the "Buntse Kei" are displayed in the town square. Plant seeds and artefacts indicating human presence have also been found, suggesting some archaeological significance.

De Kuilen are known for fossil shells from the Pliocene, which can still occasionally be found on the shores of the lakes, but are especially renowned for the wide variety of fossils excavated there in the past. Langenboom is recognized across Europe as one of the most productive fossil sites from the Miocene and Pliocene—holding the top position for fossil birds. Paleontological finds include the hoofed mammal Chalicotherium, four new auk species, and the short-snouted dolphin (Protophocaena minima). At Langenboom, the extracted sand—including fossils—was sprayed onto land, allowing amateur geologists and paleontologists to easily collect specimens on a large scale.

Settlement: The most important geological feature of the area is, of course, the presence of faults. In the past, people were unaware of their existence but they adapted to and utilized the landscape with all its elements without realizing their interrelationship. The interaction between humans and the landscape is clearly visible on historical maps (see Figure 12). The Peelhorst, the higher area in between the Peelrand and Tegelen fault zones, was originally a wet and empty area where peat formation occurred in marshes. However, the edges of this area were ideal for habitation: here people were always assured of controllable water. At the northwestern edge of the Peelhorst, south of Oss, the prehistoric inhabitants made use of the existing elevation difference. For those coming from the low-lying clay area in the north, the higher Peelhorst must have been a striking feature. At Vorsel and Zevenbergen, several urn fields and many burial mounds were present on the high edge, including the famous princely graves of Oss. Burial mounds also served as markers of the "property rights" of a local community. They had to be noticeable and were erected in prominent places. Research has shown that the burial mounds around the princely grave of Oss were grazed by sheep at that time. This prevented the growth of shrubs that could obscure the burial mounds from view. As a result, and due to their high location, the mounds were visible from a great distance (Jansen and Van der Laan, 2011).

The peatland of the Peel has long been unattractive for habitation. Archaeological finds have mainly been made on sand ridges along the edge. However, tools from hunter-gatherers from the Paleolithic (2.5 Myr - 10.000 BC) and Mesolithic (10.000-8000 BC) have been found in various places. In these early periods, the Peel was not yet covered with peat; it expanded further in later times. Flint axes have been found at Ospel, Venray, and Heibloem. These tools were likely offered at water-rich places. A settlement from the late Stone Age was discovered at Millheeze. Finds at the edge of the Peel near Bakel and Deurne indicate habitation during the Iron Age and Roman Period. On June 15, 1910, the peat cutters Gabriël Smolenaers, Driekske Slaats and Loeves Pier made a spectacular discovery in the Peel marsh. At Helenaveen, they found the personal belongings of a Roman officer: a gilded silver cavalry helmet, a cloak pin, a cavalry spur, bells from a horse harness, and Roman coins. These finds date from the early 4th century. This remarkable find initiated a period in which many more finds were made in the Peel, the oldest dating back to the Mesolithic and Neolithic periods. After a period of decline, the population in the southern sandy area, and thus in the Peel region, began to increase again from the 6th-7th century. Habitation took place at some distance from the peat. During the Middle Ages, a ring of villages formed around the vast peatland. In Brabant, these included Someren, Asten, Liessel, Deurne, Milheeze, Bakel, Gemert, Handel, Boekel, Volkel, Uden, Zeeland, Nistelrode, and Schaik. To the east are Mill, Wanroij, St.-Anthonis, Oploo, and Overloon. The Limburg part of the ring includes Venray, Horst, Panningen, Helden, Weert, Nederweert, and Heythuysen. Often, the villages lay as an isolated strip of cultivated land amidst the wild grounds of the peat or the extensive marsh

TNO Public 30/219

forests and heathlands to the east towards the Maas. In addition to villages, smaller hamlets emerged, originating from a farm or a mill.

Due to the water seepage at the higher sites of faults, streams originated that were utilized for moated sites such as castles, houses and farms that could be perfectly defended thanks to the natural waterline.

Agriculture: The medieval cultural landscape consisted of arable land and grasslands around the old villages. The fields were generally located on the higher, well-drained parts of the valley slopes. There were individual reclaimed fields, surrounded by hedgerows or hedges, and contiguous open field complexes or fields.

However, in the thirteenth century, the population grew rapidly, and farmers from nearby villages began to use the cover sand ridge. They cleared the forests, and on the heathland that replaced them, they grazed their sheep for centuries. In sandy areas, it was difficult to farm without sufficient manure. Therefore, forest and heath sods ("plaggen") were cut, which were mixed with animal manure to increase the fertility of the fields. The nutrients in the manure only partially returned to the heath. At night, the manure was collected in the stable — the potstal — on a bedding of cut heath or grass. The sods were returned to the fields that became increasingly higher. These fields are referred to as "esgronden and these anthropogenic soils classify as Plaggen soils.

The advantage of manure-rich heath sods on the fields lay in the fact that a few weeks of soaking the sods with urine could kill the seeds of potential arable weeds. Because potstal manure was used to fertilise the naturally poor arable land, the already nutrient-poor sandy soil beneath the heath became even more depleted.

After the Middle Ages, as the population continued to grow, the use of the heath intensified. Cutting heath shrubs was no longer sufficient to produce enough fertiliser for the fields. Increasingly, the heath was stripped of its sods. Excessive and deep sod-cutting accelerated the degradation of the heath. The soil became exhausted, and the heath could no longer recover.

Anthropogenic wind deposits:

The severe impoverishment of the heathlands led to one of the earliest large-scale environmental disasters in history. Where vegetation failed to recover in time after cutting or sod removal, the wind took hold of the bare sand. Just as in the Ice Age, the sand began to drift again. People were virtually powerless against this natural force. Large areas of heathland, farmland, and even entire villages were buried under drifting sand.

Parts of the cover sand landscape transformed into sand drift landscapes. These drift dunes can be recognized in the landscape by their much steeper and higher profiles compared to the cover sand ridges and parabolic dunes formed during the Ice Age. The reason for this is that they developed in a landscape where scattered trees, shrubs, and hedgerows were present. These obstacles slowed the wind and trapped the sand, which was then deposited locally in steep dune ridges. The danger of sand drifts was always lurking, as the Bedafse Bergen (Figure 14) show.

To prevent new farmers from settling on the heath, commons were established to regulate the use of the land: everyone's rights and duties were laid down regarding the number of sheep to be grazed, the amount of sod and peat to be cut, and so on. Only in the French period did the common lands come into the possession of the municipalities.

) TNO Public 31/219

The common heathlands were subsequently sold, new farms were built, and part of the heathlands turned into agricultural land and forest when artificial fertilizer was introduced around 1900.



Figure 14 - The Bedafse Bergen to the west of Uden. In this sand drift landscape, inland dunes have formed that rise up to 15 metres above the original cover sand level. On top of the dunes stands coppiced oak woodland, which was planted to protect the nearby fields from drifting sand and has grown upwards along with the accumulating sand. Photograph presented in *Koomen et al., 2007.*

In the previous century, many of the sand drifts were stabilized by planting pine trees. The steep drift dunes are still clearly visible within the forests today.

Reclamation and consolidation: Ysselsteyn was founded in 1920 as a model reclamation village. The village had a Y-shaped road system (Figure 15). Various facilities, including a school and a church, were located around the junction. The farms were spread out; originally, 15 farms were built, and more followed after 1920. In 1938, 58 farms were established with state support.

After the Second World War, various developments significantly influenced the landscape (Figure 16). These were not specifically regional developments but national issues that had a regional impact. These included the ongoing scaling up in agriculture, including land

) TNO Public 32/219

consolidations, the growth of villages, the expansion of infrastructure, and the rise of recreation.



Figure 15 – The reclamation village of Ysselsteyn with its typical "Y"-shaped road pattern. Note the geometrical land-consolidation pattern radiating from the village.

In agriculture, the ongoing scaling up led to specialization in dairy farming or poultry and pig farming. Additionally, the average size of farms increased significantly. The narrow, often strip-shaped plots with alder hedges in the stream valleys were replaced by larger block-shaped parcels. The scattered plots of farmers were thus concentrated. New farms were built in places where there had never been farms before. In the region, a total of 40 land consolidations were carried out, covering at least 135,000 hectares (not all of the mentioned projects were entirely within the region). Some started in the 1930s, most took place between 1960 and 1980. Parts were consolidated twice. The parcelling was often changed, and the water management was thoroughly addressed, making arable farming possible even in the lowest parts of the landscape. The low-lying grassland areas became as open as the fields.

Restoration: Due to the reclamations and land consolidations after World War II, the Peel became a modern agricultural landscape. Straight roads and canalized watercourses cut through the fields and meadows. Thanks to the Land Reclamation Service, followed by land consolidation, the Peelhorst became one of the most important intensive agricultural and bioindustry areas of the Netherlands. New villages were founded, and existing settlements grew (Figure 16). In the 1970s, however, there was a shift in thinking about the malleability of the landscape.

Parts of the Peel have been transformed into Natura 2000 areas in recent decades. This happened in some cases as part of land consolidation or reallotment projects. This mainly concerned the non-peat areas, the remaining heathlands, and pieces of forest. Besides the National Park De Groote Peel, this includes the Deurnese Peel-Mariapeel, Boschhuizerbergen, Srasven-De Banen, and the Weerter- and Budelerberg-Ringselven.

There was an increased appreciation for unique elements in the landscape such as the traces of the Peelrand fault, associated wijst and the unusual contrast of high, wet grounds and low, dry grounds. With growing environmental awareness, more attention was paid to the wijst phenomenon. There was also increasing (scientific) interest in the wijst and the faults of the Peel Boundary Fault Zone. As a result, wijst areas were mapped and qualified as "earth science values" (see *Provincie Noord-Brabant, 2004*). In various places, efforts are now being made to restore wijst. Old, more or less existing seepage phenomena are being revived through smaller and larger interventions in the landscape. Overall, there is now quite a bit to see, more

) TNO Public 33/219




Figure 16 – Until well after the Middle Ages, the Peel was an empty and inaccessible area without any villages (left; the peat areas are marked in grey)). The vast barren lands and small-scale reclamation of the Peel came to an end in the 19th century. From that time onwards, large-scale peat extraction began and new villages were established (right). The peat cutting was also a form of land reclamation: the intention was to use the land exposed by excavation for agriculture and horticulture. The maps cover approximately the same area (modified after *Van Balen et al., 2022*).

than in the heyday of land consolidation when rapid water drainage was necessary to enable the use of heavy agricultural machinery. Meanwhile, due to climate change, priorities have shifted, and water retention is becoming increasingly important. Those who now follow the fault lines through the landscape will encounter not only the geological and cultural-historical traces of the fault but also new wijst restoration projects. Restoring the water retaining function of the fault system, which in many places was destroyed by agricultural activity and ground levelling for urban development, can help combat the drought of recent years and also contribute to water storage during wet periods. Knowledge of the faults and the surrounding subsurface is therefore important for construction projects and sustainable agriculture. In the past, people knew that they did not had to build on the fault and adapted crop choices to the subsurface and specific conditions. Farmers were pleased to own a wijst plot, a refuge in dry times. But local knowledge has disappeared, but thanks to all the recent attention to the fault, people realize (again) that it is time to take the fault into account. The faults are even considered for new developments, such as underground cold-heat systems and the use of geothermal energy that can make use faults.

The Geopark demonstrates how the landscape changed over time: from poor fields to nature reserves, from heathland to forests, and from farmland to recreational opportunities

) TNO Public 34/219

6 Geosites

This part gives an overview of the selected 40 geosites (for the time being 41 are selected). For the selection of the geosites a geographical and topical balance is sought for. The many different types of manifestations at the geosite are grouped according to "Interests" that, in turn, are grouped in either "Geological", "Geosite", or "Other" categories. Internation, National, or Regional significance is indicated for all geosites and a level of Unicity is assigned. Figure 17 summarizes these criteria.

Manifestation	Interest	Cat.
Elevation differences due to faults and/or river terraces	Geomorphology	99
Seepage, artesian flow, hydraulic head, waterworks (mills, moats, canals)	Hydrology	Geological interest(s)
Ores, minerals, fossils, glacial erratics	Depo & Foss	interest
Abandoned river courses: terraces, streams	Rivers & Streams	(s)
Geological age of the manifestation(s)	Geologic period	
Historical structures: castles, farmsteads, military constructions	Cultural, Hist., Archeo.	eosite i
Vegetation: reeds, alders, oaks, reptiles, birds, butterflies	Flora & Fauna	Geosite interest(s
Land use: bog iron as building material, water bodies for sand and gravel extraction, peat	Industrial or Economic	s)
Has research been conducted on the specific site?	scientific	o
Information boards, visitor experience, education	educational	Other interes
Does the site have visitor appeal?	(geo)tourist	est
Significance = unique internationally, nationally, or regionally	I = international N = national R = regional	Significance Unicieit
Unicity within the aUGGp	A = unique B = few other sites equal C = many (>5) sites equal	Unicieit

Figure 17 – Ranking and grouping criteria for the 40 selected geosites

TNO Public 35/219

Table 2 - Listing of 41 selected geosites. The yellow highlighted sites represent sites with multiple manifestations combined. Geological Elements: PF = Peel Boundarry Fault Zone, MV = Maasvallei, TF = Tegelen Fault, PH = Peelhorst

	Geositenr	naam potentiele Geosite	Geological interest(s)		Geosite interest(s)				Other interest			Type of p	Significance	Uniciteit		
element	Selected gesosites	Markage	Geomorphology	Hydrology	Depo & Foss	Rivers & Streams	Geologic period	Cultural, Hist, Archeo.	Flora & Fauna	Industrial or Economic	scientific	educational	(geo)tourist	pm - door TNO nader te benoemen	I = international N = national R = regional	A = unique B = few other sites equal C = many (>5) sites equal
MV	GS_01	Maasheggen	x			х	х	х	х		х	х	х		1	A
TF	GS_02	Steilrand Mill	х		х	x		х	х		х	х	х		1	A
PF	GS_03	Bijzonder Brabant	х	×					х		х	х	х			Α
PH	GS_04	Griendtsveen - Toon Kortoomspark-		х				х		х		х	х		1	A
PF	GS_05	Vorstengraf	х					х				х	х			A
TF	GS_06	Peelraamstelling						×				х	х			A
PF	GS_07	Sint Annabos	х	х	х	х		х	х	х	х	х	х		1	В
PF	GS_08	Hoogdonkseweg Liessel	х	х	х	х	x			х	х	х	х		-	В
MT	GS_09	Leudal	х	х	х		İ	×	х	х		х	х		N	A
	GS_10	Maasdoorkruising	х	х		х		х	х		х		х		N	A
PH	GS_11	Maria- en Deurnsche	х						х						,	В
TF	GS_12	Peel Waterzuiverende		×					х			×			N	_
25	GS_13	wiistboerderii Zeeland Slabroek	х	×	×	×		×	х			×	×			A
PF PF	GS_14	Gemert, Breuken	x	х	х	х		х	х			х	х		N	В
	GS_15	Uden - Stadswijst	×	×	x	<u> </u>		×	х			x	×		N 	Ť
PF	GS_16	Geneneind	x	×		x		×	x		х	x	×		N 	В
	GS_17	Neer	×	×	\vdash	×		x	×		×	×	×		N	В
	GS_18	Helenaveen	^	×	×	×		×	^	×	^	×	x		N	В
PH	GS_19	Wijstbos Zeeland		X	X	^		×	х	^		X	×		N	В
	GS_20	Langenboom/de Kuilen		_ X				X	Х						N	В
TF	GS_21	Sint Hubert			х		Х				х	х	х		1	В
TF	GS_22	Kasteel Deurne.	х						х	Х		х	Х		N	В
PF	GS_23	Heiakker Maashorst	х	x		_		×		х		х			N	В
PF	GS_24	De Burgt		x					х			х	х		R	Α
PF	GS_25	Donzel	Х	х	_			x				х			N	A
PF	GS 26	De Specht /Putakker		x	х				х						N	С
PF	GS_26 GS_27	De Perekker	Х	Х		X			х		х	X	х		R	В
PF			Х	×		X			х			х	х		R	В
PF MV	GS_28	Meijel Schuitwater	х	x				×				x	х		R	В
MV	GS_29	Broekhuizen-Lottum	х			х	х		х		х	х			R	В
PH	GS_30	De Bult	х					x	х		х	х	х		R	В
TF	GS_31	Loobeekdal	х						х			x	х		R	В
MV	GS_32	Maaspark Noord Limburg	х			×	х		х		х				R	В
PH	GS_33	Stippelberg Klotterpeel/De Rips	х	×				×	х						R	В
PH	GS_34	Zinkske Molentje						х	х			х			R	В
TF	GS_35	Hooge Raam		х		х			х						R	В
MV	GS_36	De Vilt				х			х			х			R	В
TF	GS_37	Watermolen Oploo													R	В
PF	GS_38	De Berkt	х	х	х	х		×	х			х	х		R	С
PF	GS_39	Het Loo	х	х	х	х			х				х		R	С
PF	GS_40	Waterbloem	х	×				×	х						R	С
PH	GS_41	De Krim	х						х			х	х		R	С
PH	GS_42	De Elsbeemden		×	х	x		×	х	x	х		х		R	В

) TNO Public 36/219

7 General geological description

The geological history of the aUGGp dates back to the Carboniferous period, hundreds of millions of years ago. From this distant past, traces have been left in the deeper subsurface. Due to movement of tectonic plates, this particular part of Northwestern Europe travelled all the way from the southern hemisphere to its current position. For example, the coal deposits that lie more than a thousand meters deep under the Peelhorst are remnants from a time in which Northwest Europe was located around the equator and was covered with extensive mangrove forest that developed under a tropical warm climate. However, for the description of the ground directly beneath our feet and the formation of the current landscape, we limit the story to the last 10 million years. The materials from this period are now at the surface or at a depth of at most a few hundred meters. It was mainly a time of sedimentation; a time of deposition of sand, gravel, clay, and sometimes peat. Additionally, erosion occurred where wind, rivers, and the sea carried away previously deposited material. The movements of the earth's crust and the climate, particularly the alternation of cold and warm periods, had a significant influence on the interplay of sedimentation and erosion.

The maps and timelines in this chapter serve as tools for the story of the geological history of the Geopark. The **geological map** (Figure 18), shows the deposits that are now at the surface, while the **cross-section** through the Geopark (Figure 19) provides an impression of the deeper deposits.

The Earth's crust is moving

The 'solid ground' beneath our feet may seem stable and unchanging at first glance. However, earthquakes and fault lines within the Geopark show that the Earth's crust is constantly in motion. The Earth's crust, the outer solid part of the Earth, is about 80 kilometres thick and consists of different plates. These plates move relative to each other across the globe at a rate of a few centimeters per year (this process is called plate tectonics). Where plates move towards each other, collision mountains are formed, and

) TNO Public 37/219




Figure 18 - DRAFT: Geological map of the aUGGP

) TNO Public 38/219

where they move apart, oceans can form. In the Mediterranean region, the Eurasian Plate and the African Plate collide. This pushed Italy northward and formed the Alps. At the same time, the North American Plate and the Eurasian Plate moved apart along the Mid-Atlantic Ridge, causing the Atlantic Ocean to grow larger. This separation creates pressure in a southeasterly direction. Due to the pressure from two sides, the Dutch part of Europe is being compressed. As a result, parts of the Earth's crust rise and others fall, such as the subsidence area to which the Dutch subsurface belongs. This subsidence area extends across Western Europe, from Spain through eastern France, Germany, to the Dutch Roer Valley Graben system. While midmountain ranges such as the Ardennes in Belgium and the Rhenish Massif in Germany are gradually being pushed up, the subsurface of the Netherlands is slowly spreading apart. Therefore, the surface of our country is continuously getting a little larger.




Figure 19 – Geological cross section through the subsurface model GeoTOP v1.6.1 (for location see the geological map in Figure 18)

As a result of these pulling and pushing forces, the Earth's crust comes under tension, and faults are formed. Most of these faults are closely related to structures that formed during the Caledonian and Variscan orogenies. These phases of plate-tectonic movement led to the collision of several micro-continents during the Late Cambium to Early Devonian and Late Devonian to Late Carboniferous periods, respectively. Eventually, the Pangea supercontinent was formed. Most of the structures formed at these phases became reactivated during several later tectonic phases, including the Cenozoic rifting. These faults form systems that branch out and eventually die out. For this reason, it is more accurate to speak of fault zones, the bundling of fault segments that together form a displacement zone. On one side of a fault (zone), the Earth's crust rises or falls faster than on the other side. In this way, a landscape with horsts and grabens can form. A horst is an area bounded by faults that rises relative to its surroundings and is therefore higher; a graben (relatively) sinks and is therefore lower.

) TNO Public 39/219




Figure 20 – Typical horst and graben structure with branching fault geometry representative for the Peelhorst and Peel Boundary Fault Zone. Modified from *Childs et al., 2009*

The Peelhorst forms the backbone of the Geopark and is bounded to the south by the Roer Valley Graben and to the north by the Venlo Graben (or Venlo Block). The Roer Valley Graben and the Peelhorst are separated by the relatively narrow Peel Boundary Fault Zone (PBFZ). The Peelhorst and Venlo Graben are separated by the Tegelen Fault Zone.




Figure 21 - The major uplift (+) and subsidence (-) areas in the Ruhr Valley Rift System and the main fault lines resulting from the movements of the Earth's crust. Modified from *Koomen et al., 2007.*

The effect of movements along the PBFZ is sometimes strongly felt when earthquakes occurs and the fault itself is sometimes clearly visible as a terrain step. Currently, the subsidence of the Roer Valley Graben relative to the Peelhorst is about one centimeter per century. This may seem small, but in a geologically brief period of one million years, it results in a height difference of one hundred meters! This cumulative effect is very evident in the more than 1 km difference in the depth of the base of the Miocene (30 million years old) on either side of the Peel Boundary Fault. However, due to continuous erosion on the horsts and infilling of the graben, the height differences caused by fault movements are largely - but not completely compensated. In the landscape of the Geopark, the graben remains visible as an area that is lower relative to the horsts. The faults that are locally visible in the landscape in Brabant and

) TNO Public 40/219

Limburg can also be traced far to the northwest in the Dutch subsurface. Because rivers and wind have always transported and deposited substantial amounts of sediment (gravel, sand, clay) here, the resulting height difference at the surface is often completely or largely filled. Therefore, the horsts and grabens in the Netherlands are mainly underground.

Fault patterns

As mentioned before, Alpine and Atlantic tectonic forces make that the Roer Valley Rift System is not only slowly spreading apart but that within the rift fault blocks move alongside each other in a right-lateral manner. Such combination of deformation is called "transtension." The fault lines shown in Figure 18 make up a complex pattern of faults and appear to be never straight lines. Faults have limited extent and can converge and diverge resulting in an anastomosing arrangement. Such an arrangement is not typical for the aUUGP, but is a commonly observed feature in faulted transtensional terrains. It is a clear indication that faulting is a dynamic process during which faults segments can start to develop at nuclei, subsequently grow in length and vertical displacement, and may eventually merge with other fault segments. Consequently, fault patterns are not simple and straightforward, something important to consider when trying to locate faults in the field

Sandbox experiments have tried to mimic the development of faults under transtension and show fault patterns that resemble those seen in the aUGGP and adjacent Roer Valley Graben (Figure 22). Those experiments also show that faults not only branch in the horizontal plane, but also in the vertical plane and may converge into a principal zone of deformation at kilometres depth. Most likely, the fault segments of the Peel Boundary Fault Zone that we observe at the surface nowadays are in fact the most shallow tips of deeper-seated faults. Further geophysical research of the deep subsurface is needed to proof this. Such deep-seated faults are likely to be inherited from much older deformation phases and probably date back to the Carboniferous period. During subsequent phase of plate-tectonic puulling and pushing, e.g., during the Triassic, Jurassic and Cretaceous periods, these faults were re-used. From the Oligocene onward (~25 Myr) the current phase of deformation started of which the aUGGP forms an area with multiple lines of evidence.

Earthquakes

If movements along faults in the Earth's crust occur in pulses, they are referred to as earthquakes. These are usually light to moderate in the Netherlands. Nevertheless, the Peel Boundary Fault Zone is considered an exceptionally active fault structure by Central European standards. The most powerful earthquake observed at a fault in the Netherlands was the Roermond earthquake in 1992, with a magnitude of 5.8 on the Richter scale (Mw 5.3). Even though there was no surface rupture, the damage in the Netherlands alone amounted to 77 million euros. The last known earthquake before that was the one in Uden in 1932, with a magnitude of 5.0. In the years following the Roermond earthquake, research was conducted in Belgium, Germany, and the Netherlands to gain more insight into the frequency and intensity of earthquakes. More specifically, to answer the question if it is possible that earthquakes from a more distant past were even stronger. In the Netherlands, the Peel Boundary Fault Zone near Neer and the Feldbiss Fault Zone near Born were investigated with trench studies.

) TNO Public 41/219




Figure 22 – DRAFT: Three-dimensional geometry of an idealized early stage pull-apart basin developing in 5° transtension based on the results of analogue modelling. From: *Wu et al., 2009*. This geometry is with enechelon arrangement of branching faults is representative of the Roer Valley Graben (central part of figure) and the Peelhorst (left part of figure).

Results from trenching studies located near Bakel and Neer, in the central and southeastern parts of the PBFZ, respectively, provided evidence for two surface rupturing paleoearthquakes. The largest earthquake had an estimated magnitude of Mw \sim 6.8 and a surface rupture length of at least 35 km, approximately 10-20 times stronger than the one in Roermond (Van Balen et al., 2019). As it took place around the Late Pleniglacial - Late Glacial transition there is likely a connection between the large-scale surface movements caused by the melting of ice caps in Scandinavia, Great Britain, and the North Sea, and the local, severe earthquakes of that time (e.g. Houtgast et al., 2005; Van Balen et al., 2019, 2021)). Results from a new trench situated (Van Balen et al., 2024) at the northwestern part of the PBFZ, near Uden, shows evidence for three to four paleo-earthquakes, of which three were surface rupturing. These comprise two normal faulting- and one, younger trans-tensional displacement. The normal faulting events have ~1 m vertical displacements each, which translate into magnitudes of Mw ~7. Like the previous results, they occurred during the Late Pleniglacial-Late Glacial transition, at \sim 15 ka and \sim 14 ka. The younger transtensional event occurred sometime during the Holocene, pre-dating an unaffected, 13th century man-made paleo-channel on the hangingwall. The potential fourth, non-surface rupturing earthquake is indirectly evidenced by loading deformations of a sand layer and a collapsed brick-wall in the infill of the paleo-channel. Comparison of the trenching results along the PBFZ, show that for one event a correlation is possible and would indicate a surface rupture length of at least 55 km. Combined, all trenching results indicate that the characteristic maximum rupturing displacement is ~1 m, corresponding to Mw ~7 as the maximum magnitude of paleoearthquakes along the PBFZ. Based on observations, the period between large earthquakes appears to be thousands of years. Thus, it is not ruled out that severe earthquakes can occur today. Moreover, lighter earthquakes occur once every few hundred years and can also cause considerable damage.

Vertical motions and rivers

The movements of the Earth's crust have had a significant impact on the development of the river system in southern Netherlands. For instance, 2 million years ago, the Meuse (Maas)

) TNO Public 42/219

flowed eastward at the height of South Limburg and emptied into the Rhine north of Aachen. Due to the uplift of the Ardennes-Eifel Massif, the Meuse, like the Rhine, had to shift its course westward.

The continuous subsidence of the Roer Valley Graben led to the accumulation of a substantial sediment package, particularly in the Roer Valley Graben. The Roer Valley Graben, and to a lesser extent the Venlo Block and the northern part of the Campine Block, have been filled with relatively fine fluvial sediments from the Late Miocene and Early Pleistocene (Kiezeloöliet, Waalre and Stramproy Formations (*Westerhoff et al., 2008*). These deposits are (mainly in the Roer Valley Graben) overlain by coarse river deposits from the Rhine and Meuse dating to the early part of the Middle Pleistocene (Sterksel Formation). During the later part, the Cromerian (850,000 – 465,000 years ago), the Rhine shifted to a more northerly course, thereby leaving the Roer Valley Graben. As a result, the Meuse became the dominant river in the southern Netherlands. The coarse fluvial sediments of the Meuse in the Roer Valley Graben system belong to the Beegden Formation.

From 400,000 years ago, after filling the Roer Valley Graben, the Meuse also shifted north-eastward, over the now rapidly rising an tilting Peelhorst. The Meuse then descended from this emerging horst and has since flowed into the Venlo Graben located to the east. Due to this shift in the Meuse's course, and the much lesser subsidence compared to surrounding tectonic blocks, the fine, shallow marine and nearshore sediments from the Miocene (Breda Subgroup) on the Peel Block are directly overlain by the coarse Meuse sediments (Figure 23).

In addition to Meuse deposits, local river and large-scale aeolian sediments were also deposited in the Roer Valley Graben system (Boxtel Formation). Generally, the sediment package belonging to the Boxtel Formation is thicker in the relatively strongly subsiding Roer Valley Graben than on the other blocks of the Roer Valley Graben system.




Figure 23 – Blockdigram showing a cross-section throug the Roervalley Graben and adjacent uplifte fault blocks (based on DINOloket), HH = Heerlerheidebreuk, GF = Geleenbreuk, FF = Feldbissbreuk, KB = Koningsbosch breuk, BE = Beegdenbreuk, PBFZ = Peelrandbreukzone. TFZ = Tegelenbreukzone, VFZ = Viersenbreukzone. (modified after *Woolderink, 2021*).

) TNO Public 43/219




Figure 24 – Due to the movements of the Earth's crust and glaciations, the Rivers Rhine and Meuse were repeatedly forced to change their courses. The maps indicate the locations of the major rivers in southern Netherlands at different points in the past. Modified from *Busschers et al., 2025* and references therein.

High is old, Low is young

The Geopark is located at the transition from uplift to subsidence areas, and through the presence of horsts and grabens, has a great variety in the composition of the subsurface. Old deposits lie close to the surface in the relatively rising areas such as the Peelhorst (Figure 25). Here, river deposition stopped long ago, and as a result of erosion, some of the older deposits have disappeared. In the subsiding areas, the old sediments are buried under thick layers of younger deposits. This occurred in the north and west of the province and in the Roer Valley and Venlo Graben.




Figure 25 - Schematic representation of landscape development due to uplift of fault blocks (horsts) and subsidence of fault blocks forming grabens. This causes the younger (lilac) geological layers on the horst to be eroded, exposing the older deposits (pink) at the surface. In the subsiding graben, younger river and eolian deposits (salmon pink) are subsequently deposited. Modified from *Koomen et al., 2007.* Consequently, the younger deposits are at a lower elevation and positioned adjacent older deposits at higher elevations ("high is old, low is young").

Time travel through the Miocene

Subtropical seabed deposits

The oldest deposits found close to the surface in the Geopark date back to the Miocene (Fig. 14). This geological period lasted from 23 million to 2.6 million years ago. During the Neogene, the Netherlands had a temperate warm to subtropical climate, and the Geopark area was covered by a shallow sea. Green to green-black colored loamy sand was deposited on the seabed. The color is due to the mineral glauconite, which forms in warm, calm, shallow seas. Due to subsidence, this 'green sand formation' lies more than 300 meters deep in the Roer

) TNO Public 44/219

Valley Graben. However, on the Peelhorst, these sediments are found almost at the surface in some places, where they can be easily extracted. From the Pliocene (about 5 to 2.6 million years ago), the sea slowly retreated westward and became shallower. Shell and sand banks were mainly deposited on the seabed. Subsequently, the shallowing continued, and parts of Brabant and Limburg fell dry, allowing rivers coming from the south and east to shift their course to the northwest. The seawater was replaced by fresh water infiltrating into sand layers in the subsurface, these now form an important source for drinking water supply.

The first river deposits

Due to the continuous uplift of the Peelhorst, parts fell dry, and the rivers coming from the south and east extend their course to the northwest. The material brought by these predecessors of the Rhine, Meuse are very rich in quartz; the sand is almost white in color. This is due to the prolonged, intensive weathering of rocks in the source areas under the warm climate conditions during the Neogene. Under these conditions, quartz is more resistant to weathering than other minerals, so after a long time, almost only pale quartz grains remained.

Time travel through the Pleistocene

Ice ages

The Pleistocene epoch, which lasted from 2.58 million to 11,700 years ago, is characterized by an alternation of ice ages and warmer 'interglacial' periods (Figure 26). This highly variable regime of cold and warm times had a significant impact on the geological development of the Dutch subsurface. During ice ages, which typically lasted from several tens of thousands of years to about 100,000 years, the temperature across the entire Earth dropped. The main cause was changes in the Earth's orbit around the sun and the associated major changes in ocean currents. Due to the temperature drop, the snow in the polar regions and high mountains like the Alps no longer melted, resulting in the formation of ice caps several kilometres thick. Under its own weight, the ice in such an ice cap deforms and flows very slowly. Consequently, the ice caps could expand enormously during the coldest periods. During the penultimate ice age, the Saalian (370,000 to 130,000 years ago), the Scandinavian ice cap reached the central Netherlands. The Geopark area was never covered by ice, but it had the climate of a polar desert during the coldest phases. Enormous amounts of water were trapped in the ice caps. As a result, the sea level dropped to more than 100 meters below the current level. Consequently, the North Sea was largely dry land, and the rivers flowed hundreds of kilometres further to the west into the sea. Between the ice ages, the climate was similar to that of our time. The ice caps melted, the sea level rose again, and vegetation returned, initially as tundra and taiga landscapes. These warmer periods lasted about 10,000 to 20,000 years. The current warm period, the Holocene (from about 12,000 years ago to the present), is also a warm interphase between two ice ages. When exactly the next ice age is to be expected has become rather unpredictable due to human interference with climate.

Rivers

On the Peelhorst and Maasvallei are, the currently visible deposits from the Pleistocene largely date back to the Weichselian Stage. Especially during its glacial periods, the rivers transported a lot of material from the hinterland, significantly expanding the delta in the North Sea basin. This delta mainly consists of sandy and gravelly sediments from the predecessors of the Rhine and Meuse. The fact that rivers transport large masses of sand and gravel during ice ages is related to the worsening climate during those periods. The soils were completely or partially frozen, and vegetation significantly declined. Under these conditions, substantial amounts of meltwater are released in a short time during spring or summer, greatly enhancing the runoff of soil material. This led to the formation of so-called 'braided rivers,' that are characterized by a wide floodplain with many channels that are almost dry for much of the year (Figure 27).

) TNO Public 45/219

During wet periods, the fast-flowing river carries large amounts of water and sediment for a short time. The floodplain is then largely underwater, and the channels shift rapidly, with sand and gravel bars (shoals) being continuously eroded and rebuilt elsewhere. The appearance of the floodplain can thus change significantly in a short time.

In the Roer Valley Graben, the coarse sandy deposits of the braided Rhine and Meuse are about 70 meters thick. They lie here 25 to 50 meters below the surface (see also Figure 23). In the relatively rising areas of the Peelhorst, these river deposits are thinner. The river deposits from the ice ages now form an important source of coarse sand and gravel for use as construction materials and are therefore exploited at various places.

Cover sand

Only in a few parts of Geopark do the Pleistocene river sands lie at the surface. Almost everywhere else, they are covered by a sand layer that was deposited by the wind at the end of the Late Glacial, from about 30,000 to 12,000 years ago. Under extremely cold and dry climate conditions, the wind in the then 'polar desert' took hold of the sparsely vegetated soil, causing it to drift. The sandy valley floors of the braided rivers and streams were particularly susceptible to drifting. Almost the entire landscape became covered with this sand, which is therefore called 'cover sand.' A lot of cover sand ended up in the Roer Valley Graben (Figure 18). Here, on the gravelly deposits of the braided Meuse (Maas) and Rhine, there is a sand layer up to 30 meters thick, consisting of cover sand, interspersed with loamy sand deposited by streams and surface runoff. Only the upper part of these sediments dates from the last ice age; at the bottom are cover sand and stream deposits from previous ice ages. On the Peelhorst, the cover sand is at most a few meters thick and sometimes completely absent.

Time travel through the Holocene

Climate gets warmer

The Holocene is the warm period that began about 12,000 years ago and continues to this day. The improvement in climate brought significant changes to the landscape. Due to the milder climate, vegetation could thicken into a closed forest that covered the entire landscape. The water discharge through rivers and streams became more uniform, and they changed from braided to meandering watercourses. Globally, the ice caps melted, causing the sea level to rise by more than 100 meters. The North Sea area disappeared underwater, and the sea eventually reached as far as the west of the Province of Noord-Brabant.

) TNO Public 46/219




Figure 26 – DRAFT: Geological timeline form the Miocene to Recent, i.e. of the last 10 million years. Just as in archaeology, in geology, names are given to certain time periods and to the depositional layers in the subsurface. On the temperature curve under the heading 'Climate,' the blues represent the ice ages, and the reds represent the warmer interglacial periods. The column 'Geological Elements' shows the approximate ages of the Meuse deposits (Beegden Formation) in the corresponding element. Modified from *Koomen et al., 2007.*

TNO Public 47/219




Figure 27 - Diagram of a braided river. Modified from Koomen et al., 2007.

Natural levees and floodplains

At the relative warm onset of the Late Glacial (Allerød and Bølling), the Meuse changed from a braided to an exceptionally broad meandering river. Subsequently, during the Younger Dryas glacial, incised braided river valleys formed for a relatively short time (~1200 yr). At the beginning of the Holocene, warmer conditions returned and the Meuse is a meandering river ever since. Although less extensive than during the Allerød and Bølling, the meandering river formed distinct bends, or meanders. In the outer bend, the water flows faster, causing the cut bank to be eroded. In the inner bend, the water flows more slowly, and sand and clay are deposited on the point bar. This process causes the bends to become wider, creating large loops in the river. Eventually, the meander bends meet, and the river cuts off its own meander loop. The cut-off meander loop remains in the landscape as an abandoned river arm (a socalled horseshoe lake). Over time, the abandoned channel becomes completely filled and remains visible only as a slightly lower, wet strip in the landscape. A meandering river also experiences periods of high water discharge, for example, when there is a lot of rain in the hinterland. The river then overflows its banks. Just outside the riverbed, the speed of the inland-flowing water decreases, and mainly fine sand settles. Over the years, this process creates natural levees (ridges) directly next to the riverbed, which are relatively high and can be considered as more or less natural dikes. The old villages, roads, and fields are located on the natural levees of the Meuse, which are several tens to several hundreds of meters wide. In the lower floodplains behind the levee, river clay is deposited. The floodplains have wet, heavy clay soils and are now open grassland areas.

) TNO Public 48/219




Figure 28 – Diagram of a meandering river. Modified from Koomen et al., 2007.

Raised bogs

Most of the Peelhorst, the area known as the Peel, has the shape of an extensive, very flat plateau. This flat terrain is the main reason why extensive marshes could form here because there were no slopes or stream valleys to drain the water. This, however, was not always the case. During the last ice age, there were still meltwater valleys on the horst. At the end of the last ice age, however, these valleys were filled with blown-in cover sand. A plateau with an irregular relief of shallow basins and low cover sand ridges was formed. When it got warmer and more humid after the last ice age, the basins filled with water, creating a landscape with numerous small lakes. After the Last Ice Age, plant growth also returned. The lakes (fens) became overgrown with reeds, sedges, and rushes, and peat began to form (Figure 29). As plant remains in the water do not decompose, the fen slowly filled with peat and became overgrown. This peat, which is nourished by mineral-rich stream or groundwater, is called 'low peat' (laagveen). Low peat can develop into raised bog (hoogveen) if the peat layer becomes thicker and the groundwater is out of reach of the roots of the marsh plants. The plants then only have access to rainwater, which contains few nutrients. Sphagnum moss is one of the plants that can grow under these poor conditions. The dying moss plants are not decomposed in the wet environment and form a good substrate for new moss plants. Sphagnum moss also acts like a sponge. It retains water very well, keeping the marsh soaking wet even as it rises higher. This creates a so-called raised bog cushion that can grow to more than ten meters thick. Raised bog cushions (or domes) can also expand laterally over originally drier areas, such as cover sand ridges. At the peak of peat growth, about 4,000 years ago, raised bogs with diameters of kilometres formed on the Peelhorst, and the raised bog eventually reached an area of 30,000 hectares. Only remnants of the raised bog domes now remain. Medieval people managed to drain the extensive peatlands by digging ditches and canals. Moreover, peat proved to be an excellent fuel. Many peatlands were completely lost due to peat extraction.

) TNO Public 49/219




Figure 29 - The formation of low peat and raised bogs. A body of water becomes overgrown and fills with low peat. This nutrient-rich peat develops into an increasingly nutrient-poor type of peat. Eventually, the marsh is fed by rainwater and becomes nutrient-poor enough for sphagnum moss to establish itself and dominate other plant species (Modified from *Koomen et al., 2007*, after *Visscher, 1949* and *Berendsen, 2005*).

) TNO Public 50/219

The human touch

Early habitants

About 6000 years ago, humans in Northwest Europe began to leave their mark on the development of the landscape. Until then, our prehistoric ancestors lived off what the primeval forest had to offer. They hunted wild animals and gathered fruits and roots. They did not live in fixed places but in hunting camps. These were set up near ponds and streams, where water was available and where the game gathered to drink. Humans gradually began to use the forest more intensively. During the Stone Age (before 4100 years ago), more and more trees were cut down for the production of firewood and construction wood. During the Bronze Age (4100 to 2600 years ago), shifting cultivation became popular, and patches of forest were cleared for the creation of small plots of land. However, fields on the nutrient-poor cover sand soils were exhausted after a few harvests. Farmers had to constantly move to new plots of land. It took about 25 years for the abandoned fields to become sufficiently fertile again for reuse under secondary forest.

In the Iron Age, starting around 2600 years ago, deforestation accelerated. A huge amount of oak wood was needed to produce charcoal for smelting iron. The forests could no longer recover, so by Roman times, large areas of forest had degraded into heathlands. After the Roman period, the forests somewhat recovered as the population decreased. In the Middle Ages, the population increased again, and land use intensified further. Many predecessors of the current villages and towns appeared on the cover sand ridges. For centuries, soil fertility remained the greatest concern of the farmers was how to make productive fields on poor soil? The wet parts, such as stream valleys, marshes, and low sand grounds, would remain almost uninhabited and unused for hundreds of years. Between the tenth and twentieth centuries, these areas were gradually drained and reclaimed

The peat invades

Until about 4,000 years ago, the cover sand ridges in the Peel landscape were habitable. After that, people were driven out of the area as the peat domes completely overgrew the cover sand ridges. In the Middle Ages, the Peel marshes became a gigantic and inaccessible marsh barrier on the border of Brabant and Limburg. Archaeological finds still remind us of life on the 'drowned' cover sand ridges. When habitation was no longer possible, pre-Christian people still came to the marshes to connect with the world of the gods. In 1910, a gilded silver Roman military parade helmet was found near Helenaveen. This so-called 'Helmet of Deurne' is probably an offering to a peat god Figure 30).

) TNO Public 51/219



Figure 30 – The Golden Helmet of Deurne was discovered in 1910 by Gebbel Smolenaar, Gebbel Smolenaars, Loeves Pier en Driekske Slaats, peat cutters from Meijel. The helmet is currently exposed in the Rijksmuseum of Oudheden in Leiden

Peat extraction by farmers

The peatland only became attractive to humans again when wood as a fuel became scarce. From the Middle Ages, farmers began to dig peat on a small scale at the edges of the marshes. They dried the peat and burned or sold the resulting turf. Over the centuries, large peat marshes were dug away down to the underlying sand. The excavated areas were abandoned, leading to the formation of extensive heathlands. In addition to peat extraction by farmers, there was also systematic peat extraction. For example, from the monastery Padua, founded in 1742, the peatlands around Boekel were systematically excavated. The monks transformed the excavated sandy soils into agricultural land.

Industrial Peat Extraction

Large-scale peat extraction in the Peel began much later than in, for example, West Brabant, where turf was already being extracted on a large scale in the Middle Ages. This was due to the proximity of markets and the better transport options in West Brabant and South Holland. The construction of canals such as the Zuid-Willemsvaart, the Noordervaart, the Helenavaart, and the Deurne Canal was a significant stimulus for peat extraction in the Peel. In 1853, the Van de Griendt brothers bought 610 hectares of peatland from the municipality of Deurne and started the first industrial peat company. Large areas on the edges of the Peel had already been excavated by farmers and monasteries, but the thickest peat layers remained. The industrial method of peat extraction followed a strict pattern, employing hundreds of peat workers. In the peak year of 1893 alone, about 700 peat workers worked for the Griendtsveen company. That year, they excavated 107 hectares of peat marsh, producing 960,000 m³ of turf.

) TNO Public 52/219




Figure 31 – Map from around 1850 showing the fens on the Peelhorst and the Groote Peel along the watershed between the provinces of Limburg and Noord-Brabant.

Changing agricultural landscapes

Until 1900, a relatively small part of the Brabant landscape consisted of fields. A much larger part consisted of vast heathlands, which initially arose from deforestation and forest degradation. To maintain the fertility of the fields, farmers used manure from sheep that grazed on the heath. In the evenings, the flocks returned to the barns in the village, where the manure was collected on a bed of heather litter. The farmers spread the mixture of litter and manure on the fields. If there was a shortage of manure, the heath was not only mowed but also turfed. The entire sod was then removed and used as litter in sheepfolds. This method succeeded in giving the fields good fertility. However, grazing and turfing made the heath soil increasingly poorer.

) TNO Public 53/219



Figure 32 - The higher elevation and bulging shape of old fields are usually clearly visible in the landscape, as seen here near Bakel (source: https://www.landvandepeel.nl).

The old arable lands are known in the south of the Netherlands as 'es'. In North Brabant, they are often referred to as 'akkers' or 'oude akkers'. The applied manure contained not only humus but also sand grains. By applying sandy manure to the field for centuries, the field layer became increasingly thicker, sometimes even more than a meter. The old fields therefore lie visibly higher than the immediate surroundings and are also called 'bulging fields' (Figure 32). Occasionally, various objects and waste residues were added to the field layer with the manure. Because many remains of old habitation have been preserved under and in the 'enk' covers, they are archaeologically valuable pieces of the soil archive.

Until 1900, the villages with their fields lay like cultural landscape islands in large open heathlands. Most of these heathlands were converted into farmland or forest in the last century. However, some beautiful heathlands still remain.

Nowadays, a large part of the Peelhorst consists of young agricultural landscapes. Sometimes these areas are the result of the planned completion of industrially excavated peatlands. In other cases, heathlands were reclaimed that had sometimes been created centuries earlier from farmer peat extractions. In yet other cases, heathlands were reclaimed that had never been covered by peat.

) TNO Public 54/219



Figure 33 - The large-scale, geometric parceling pattern of the landscape of the recent reclamations on the Peelhorst is clearly recognizable on the topographic map, as seen here in the Wanroijsche Peel north of the village of Landhorst.

Just like in the cover sand area, the heathlands could be used as agricultural land from 1850 onwards thanks to the introduction of artificial fertiliser. The first agricultural reclamations were carried out by farmers, resulting in a small-scale landscape. From the end of the 19th century, municipalities increasingly involved themselves in the reclamation of heathlands and raised bogs. The so-called 'young reclamations' from this time have large plots and widely scattered farms with names like Annahoeve, Rosinahoeve, and Reginahoeve. These reclamations usually have a rectangular and regular layout of plots and roads (Figure 33). They resemble the young reclamations in the cover sand landscape but are much larger in scale. Unlike in the cover sand landscape, they are not interspersed with old fields or stream valleys. Just like in the Dutch polders of the 19th and 20th centuries, this is a new landscape of tens of thousands of hectares. In the lower-lying areas, where the first peatlands once formed, it can still be wet. Here, there is often a dense network of ditches and trenches to promote drainage.

Peat Remnants

Not all the peat has disappeared. Especially in the southern part of the Peel, remnants have been preserved. These are the most extensive raised bog remnants in the Netherlands, but there are no longer large, growing raised bog cushions. The peat has been too drained and harvested, and precipitation and groundwater have become too rich in nutrients due to bioindustry. Nature management in the Groote Peel, as well as in the Deurnsche- and Mariapeel, aims to restore the raised bog landscape. These nature reserves now give an impression of the former vast wildernesses. The canals and peat pits bear witness to the hard life of the peat workers.

TNO Public 55/219

Key references

The references listed here are the ones specifically mentioned in the text and figures of this document and can be considered as key references. Note that a full Bibliography (compiled by the Scientific Board of the Geopark Peelhorst & Maasvallei) is presented as Appendix B.

- Ahorner, L. (1994) 'Fault-plane solutions and source parameters of the 1992 Roermond, the Netherlands, mainshock and its stronger aftershocks from regional seismic data', in: Geologie en Mijnbouw 73 (2-4, Special Issue), 199-214.
- Berendsen, H.J.A., 2005. Landschap in Delen. Van Gorcum, Assen, the Netherlands, 340 pp. Bon, J. (1972), 'Hydrologische veldkenmerken langs de Westflank van de Peelhorst', in: Tiidschrift van het Koninklijk Nederlands Aardrijkskundig Genootschap, 65pp.
- Busschers, F.S., Cohen, K., Wesselingh, F.P., Schokker, J., Bakker, M., Van Balen, R. & Van Heteren, S., 2025. Quaternary. In: J.H. Ten Veen, G.-J. Vis, J. De Jager, J. & Th.E. Wong (eds): Geology of the Netherlands, second edition. Amsterdam University Press (Amsterdam): 333-391. DOI: 10.5117/9789463728362 ch10
- Childs, C., Manzocchi, T., Walsh, J. J., Bonson, C. G., Nicol, A. & Schopfer, M. P. J. (2009). A geometric model of fault zone and fault rock thickness variations. Journal of Structural Geology, 31, 117–127.
- Ettema, N. (2021) Ecologische iconen van de Wijst Wat breuken ons bovengronds laten zien. Geopark Peelhorst en Maasvallei i. o., Uden, 121 pp.
- Ettema, N. (2022) Breuken in het land van Peel en Maas: veldgids' Geopark Peelrandbreuk en Maashorst i.o. Uitgeverij Matrijs (Utrecht), 152 pp.
- Houtgast, R.F. & Van Balen, Ronald & Kasse, Cornelis (Kees) & Vandenberghe, J.. (2003). Late Quaternary tectonic evolution and postseismic near surface fault displacement along the Geleen Faut (Feldbis Fault Zone Roer Valley Rift System, the Netherlands), based on trenching. Geologie en Mijnbouw/Netherlands Journal of Geosciences. 82. 10.1017/S0016774600020734.
- Iven, W. and van Gerwen, Th. (1978) Mensen in de Peel, Distel, 78 pp.
- Jansen, R., en K. van der Laan (2011), Verleden van een bewogen landschap, Landschapsen bewoningsgeschiedenis van de Maashorst, Mattrijs (Utrecht), 176 pp.
- Joosten, J.H.J. (1991). Die Geschichte der Morre und der Moornutzung im Peelgebiet (Niederlande). Temla, Band 21, 75-100.
- Koomen, A. J. M., Kiden, P. & Verbauwen, E. (eds.) (2007). Van beekdal tot stuifduin Aardkundige waarden in Noord-Brabant. Provincie Noord-Brabant ('s Hertogenbosch): 120 pp.
- Lapperre, R. E. (2025). Fault-Zone Hydrogeology in Unconsolidated Sediments: Cross-Fault Groundwater Levels, Fault Sealing, and Permeability Distribution in the Roer Valley Rift System. [PhD-Thesis Research and graduation internal, Vrije Universiteit Amsterdam]. https://doi.org/10.5463/thesis.1223
- Lapperre, R.E., Bense, V.F., Kasse, C. et al. (2022) Temporal and spatial variability of cross-fault groundwater-level differences: the impact of fault-induced permeability reduction, precipitation and evapotranspiration. Hydrogeol J 30, 1233–1257. https://doi.org/10.1007/s10040-022-02465-w

) TNO Public 56/219

- Munckhof, P.J.J. van den & Joosten, J.H.J. (1990): Een breuk net een verleden. Beheers- en ontwikkelingsvisie voor de Grote Eenheid Natuurgebied zuidelijke Peelhorst.- 181 s., 58 Abb.; Naluur Milieu en Faunabeheer Tilburg/Roernond.
- Provincie Limburg, (2009) Landschapskader Noord- en Midden-Limburg, 68 pp.
- Provincie Limburg (2021) Omgevingsvisie Limburg, 204 pp.
- Provincie Noord-Brabant (2004) Aardkundig Waardevolle Gebiedenkaart Noord-Brabant, 173 pp.
- Stouthamer, E. & Berendsen, H.J.A.. (2001). Avulsion Frequency, Avulsion Duration, and Interavulsion Period of Holocene Channel Belts in the Rhine-Meuse Delta, The Netherlands. J. Sediment. Res.. 71. 10.1306/112100710589.
- Vos, P., Van der Meulen, M., Weerts, H. & Bazelmans, J. (2020) Atlas of the Holocene Netherlands landscape and habitation since the last ice age. Amsterdam University Press (Amsterdam), 96 pp.
- Van Balen, R. T., Lapperre, R. E., Woolderink, H. A. G., Wallinga, J. & Kasse, C. (2024), In: Tectonophysics. 879, p. 1-15 15 p., 230322.
- Van Balen, R.T., Bakker, M.A.J., Kasse, C. & Woolderink, H. (2019) A Late Glacial surface rupturing earthquake at the Peel Boundary fault zone, Roer Valley Rift System, the Netherlands. Quaternary Science Reviews 218:254. Van Balen, R.T., N. Ettema, H. Kuijpers, R. Lapperre, B. Nelemans, J. Timmers, G. Verbeek en E. Weerman (2022), Breuken in het land van Peel en Maas' Geopark Peelrandbreuk en Maashorst i.o. Uitgeverij Matrijs. 176 pp.
- Van Balen, R.T., Kasse, C., Wallinga, J., Woolderink, H.A.G. (2021) Middle to Late Pleistocene faulting history of the Heerlerheide fault, Roer Valley Rift System, influenced by glacioisostasy and mining-induced displacement, Quaternary Science Reviews, Volume 268, 107111
- Van den Berg, M., Vanneste, K., Dost, B., Lokhorst, A., Van Eijk, M., Verbeeck, K. (2002). Paleoseismic investigations along the Peel Boundary Fault: geological setting, site selection and trenching results. Netherlands. J. Geosci. Geol. Mijnb. 81, 39–60.
- Visscher, J. (1949) Veenvorming. Noorduijns wetenschappelijke reeks (Gorinchem). nr. 33, 115 pp.
- Westerhoff, W.E., Kemna, H.A. & Boenigk, W., 2008. The confluence area of Rhine, Meuse, and Belgian rivers: Late Pliocene and Early Pleistocene fluvial history of the northern Lower Rhine Embayment. Netherlands Journal of Geosciences 87: 107-125. DOI: 10.1017/S0016774600024070
- Wu, J. E., McClay, K., Whitehouse, P., & Dooley, T. (2009). 4D analogue modelling of transtensional pull-apart basins. Marine and Petroleum Geology, 26(8), 1608-1623.

TNO Public 57/219

) TNO Public) TNO 2025 R11935

) TNO Public 58/219

Signature

TNO) Energy & Materials Transition) Utrecht, 24 September 2025

Denise Maljers Research manager Johan ten Veen Author

Patrick Kiden Review

) TNO Intern 59/219

) TNO Intern) TNO 2025 R11935

) TNO Intern 60/219

Appendix A

Geosites descriptions

) TNO Intern 61/219

) TNO Intern) TNO 2025 R11935

) TNO Intern 62/219

Geosite: GS_01	Name
Significance : International	Maasheggen
Coordinates (RD): 197704, 404015	
Municipality	Land van Cuijk
Province and country	Noord-Brabant
Protection status	UNESCO MAN & BIOSPHERE status, Aardkundig
	waardevol gebied (AWG)
Site management organizations	Staatsbosbeheer; Steering and Core Group De
	Maasheggen
Accessibility	The area can be accessed via several "gateways", where parking is available. At these locations, Maasheggen Ambassadors are present, who can provide further information about the area. The area can only be entered on foot or by bicycle
Relevance for tourism	Numerous signposted walking and cycling routes traverse De Maasheggen.
Relevance for education & information	Family routes, Museum Ceuclum in Cuijk, Maasheggen weaving workshops, treasure hunts. Various children's activities.
Teaching equipment	
Scientific interest	Woolderink et al., 2018

Description

1. Location

De Maasheggen represents a landscape and comprises a chain of nature reserves along the River Meuse between the Noord Brabant towns of Vierlingsbeek and Cuijk. It is a 25 km long, elongated area of 6,700 ha, directly adjacent to the Meuse. The Maasheggen are best preserved in Noord Brabant, between Vierlingsbeek and Sambeek; this area forms the geosite.

2. Landscape formation and description

Origin of the landscape:

Since the last Ice Age, the Meuse has shifted in a north-easterly direction on the tectonically tilting Venlo Block. During the Holocene, there have been above-average shifts in the river valley of the Meuse, recognisable by abandoned meander bends. The stream valleys, which developed mainly after the Younger Dryas, indicate a drainage system caused by tilting, originating at the Tegelen Fault and discharging into the Meuse. The Maasheggen lie in the western floodplains of the Meuse and display a landscape where, for thousands of years, woven hedges have been used to demarcate pastures. This has created a unique landscape with a mosaic of hedges, meadows, drinking pools, and ancient pollarded trees.

Geomorphology:

This geosite lies in the floodplains of the present-day Meuse and also features some abandoned Meuse meanders due to human intervention.

) TNO Intern 63/219

Hydrology:

The hydrology of the Maasheggen is strongly influenced by the proximity of the Meuse and the presence of drinking pools, which serve as water sources for livestock and other animals. Water safety is one of the greatest challenges for the Maasheggen, situated along the Meuse. The dense hedge structure, consisting of hawthorn and other shrubs, helps retain rainwater and reduce surface runoff, thus preventing the soil from drying out quickly. The damming effect caused by additional hedges in the Meuse valley must be compensated by water-lowering measures. The hydrology of the Maasheggen is therefore a complex interplay between the Meuse, the hedge structure, drinking pools, and the soil. The challenge is to take the necessary water safety measures, taking into account regional ambitions and the cultural-historical and ecological values in the northern Meuse valley.

Deposits & Fossils:

The subsoil of the Maasheggen mainly consists of Holocene Meuse terrace deposits (Oost-Maarland Member, Beegden Formation). These comprise moderately to extremely coarse and gravelly sand, fine to coarse blue-grey gravel, locally even boulders and blocks. To a lesser extent, sandy loam and clay layers are present.

River & Streams:

The Meuse is a major factor in the area's hydrology. The river influences the groundwater level and can cause flooding, which in turn affects the hedges and surrounding farmland.

3. Geosite Interests

Cultural History

The Maasheggen were once found along all the floodplains of the Meuse. However, with the invention of barbed wire (and later electric fencing), the decline of hedges as field boundaries began. Especially shortly after the Second World War, there was a sudden surplus of barbed wire, and the woven hedge virtually disappeared everywhere as a field boundary. The Maasheggen area on the border of Noord Brabant and Limburg has, however, been preserved and thus forms the oldest cultural landscape in the country. Nowhere else in Europe are nature, landscape, economy, and culture so visibly and enduringly intertwined. The Maasheggen is the only biosphere reserve in the Netherlands.

Romans: In Roman times, Cuijk was an important crossing point for the Romans. At Cuijk—then called Ceuclum—there was a bridge that enabled the Romans to cross the Meuse. In Roman times, there was a busy traffic route along the Meuse, from Tongeren and Maastricht to Nijmegen. In some places, the road surface has been found during excavations, allowing the Roman road to be largely reconstructed. The road can be followed over a greater distance in certain places by the differences in elevation in the landscape, which can be experienced along the designated cycle route "the Roman road".

Flora & Fauna:

Thanks to the shelter and protection provided by the kilometres of woven hawthorn and blackthorn hedges, De Maasheggen is home to many special and sometimes even endangered animals (beaver, badger, fieldfare, lesser spotted woodpecker, little owl, black redstart, butterflies) and plants (various varieties of hawthorn and blackthorn, spindle, cuckoo flower).


The drinking pools in the meadows are not only important water sources for livestock, but also home to toads, frogs, and newts. These amphibians lay their eggs in the pools in spring. The most characteristic is the rare great crested newt.

) TNO Intern 64/219

De Maasheggen is an important source for seeds. Of the more than one hundred authentic Dutch seeds preserved in the world seed bank in Svalbard, Norway, over sixty come from De Maasheggen.

Industrial & Economic:

- 4. Relation with other geosites
- **5. Documentation & Resources** www.maasheggenunesco.com
- 6. Maps, photos and figures



Meanderende Maasbedding →
(Allerød)

Afwateringsbeek in oude →
Maas meandet

Apvateringsbeek in oude →
Maas meandet

Left: AHN image of the current Meuse bed with the Maasheggen alongside abandoned Meuse beds. Right: reconstruction of old Meuse courses (Woolderink et al., 2018). Green – meandering Meuse before the last Ice Age, Yellow = braided Meuse during the last Ice Age, orange = meandering Meuse after the last Ice Age

) TNO Intern 65/219



Characteristic landscape of the Maasheggen with the Meuse in the foreground. (Source: www.vierlingsbeek-groeningen.nl/recreatie/maasheggen)



"Maasheggen" (Source: www.vierlingsbeek-groeningen.nl/recreatie/maasheggen)

) TNO Intern 66/219

Significance: International Coordinates (RD): 181365, 412593 Municipality Land van Cuijk Province and country Noord-Brabant (NL) Protection status Aardkundig waardevol gebied (AWG) Site management organizations Accessibility The area is easily accessible from all directions. Relevance for tourism Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features Teaching equipment Guided educational activities.	Goosito: GS 02	Name
Municipality Land van Cuijk Province and country Noord-Brabant (NL) Protection status Aardkundig waardevol gebied (AWG) Site management organizations Accessibility The area is easily accessible from all directions. Relevance for tourism Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features	Geosite: GS_02	1101110
Municipality Land van Cuijk Province and country Noord-Brabant (NL) Protection status Aardkundig waardevol gebied (AWG) Site management organizations The area is easily accessible from all directions. Relevance for tourism Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features	Significance : International	Steilrand Mill
Province and country Noord-Brabant (NL) Aardkundig waardevol gebied (AWG) Site management organizations Accessibility The area is easily accessible from all directions. Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features	Coordinates (RD): 181365, 412593	
Protection status Aardkundig waardevol gebied (AWG) Site management organizations Accessibility The area is easily accessible from all directions. Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features	Municipality	Land van Cuijk
Protection status Aardkundig waardevol gebied (AWG) Site management organizations Accessibility The area is easily accessible from all directions. Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features		
Site management organizations Accessibility The area is easily accessible from all directions. Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features	Province and country	Noord-Brabant (NL)
Accessibility The area is easily accessible from all directions. Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features	Protection status	Aardkundig waardevol gebied (AWG)
Relevance for tourism Numerous marked and unmarked cycling and walking paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features	Site management organizations	
paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and information panels. Relevance for education & information The area contains information panels, mainly about ecological and landscape features	Accessibility	The area is easily accessible from all directions.
ecological and landscape features	Relevance for tourism	paths provide excellent access to the relatively undulating, varied, and attractive landscape. The infrastructure is well developed, with rest benches and
Teaching equipment Guided educational activities.	Relevance for education & information	
	Teaching equipment	Guided educational activities.

Description

1. Location

This geologically valuable area comprises the prominent, up to 7-metre-high steep edge between the Peelhorst and the Pleistocene Meuse river landscape to the northwest and southeast of Mill. The area is relatively long and narrow: approximately 10 km in a northwest-southeast direction but only 1 to 2 km in a southwest-northeast direction.

2. Landscape formation and description

Origin of the landscape

The steep edge was formed by river erosion (of an existing fault-determined terrain step) during the last two ice ages in the Middle and Late Pleistocene, between approximately 370,000 and 30,000 years ago. The Meuse in the current Meuse valley northeast of the Peelhorst was occasionally joined by the Rhine during cold periods. Erosion by both rivers shifted the steep edge, originally further northeast, southwestwards to its current position. Geologically, the Steilrand van Mill also marks the boundary between older, Middle Pleistocene Meuse deposits on the higher Peelhorst and younger, Late Pleistocene Meuse and partly Rhine deposits to the northeast..

Geomorphology:

Prominent steep edge up to 7 metres high: A uniquely visible steep edge at the transition from the Peelhorst to the Meuse valley, intersected by stream valleys. The Graafseweg runs from Grave to Mill and features several fine slopes. As shown on the fault map, this elevation difference lies between two faults.

Hydrology:

Deposits & Fossils:

) TNO Intern 67/219

River & Streams:

Stream valleys (formed as meltwater valleys), gullies and flats (remnants of a braided river system).

3. Geosite Interests

Cultural History

Valuable landscape elements include the prominent windmills 'De Korenbloem' in Mill and the Heimolen southeast of Mill. The settlement of Bruggen contains several characteristic farms. The military defences of the Peel-Raamstelling along the Defensiekanaal (Peelkanaal) are a well-preserved and coherent example of a casemate line in the tradition of permanent fortress construction from 1939–1940. The low-lying open stream valley of Mill, between the old village centre and post-war expansion districts and included in the Peel-Raamstelling, has high cultural-historical value.

Flora & Fauna:

Valuable historic green structures include scattered woods and hedgerows on the northwest edge of the area (Reeksche Bergen), on the Russendaal estate, the hedges, hedgerows and scattered small woods of the De Tongelaar estate near the Kammerberg, the wetland areas along the Graafsche Raam and parts of the Beerse Overlaat along the northeast boundary. Linear green structures of high cultural-historical and ecological value include the old railway embankment east of Mill and the hedgerows and tree belts along the Peelkanaal. Ecologically valuable landscape elements are found near the Graafsche Raam, Langven and Kammerberg. Much of the woodland on the steep edge slope and on the high side of the steep edge forms part of the Ecological Main Structure. The Peelkanaal and valley landscape at Mill form an ecological corridor.

Industrial & Economic:


4. Relation with other geosites

Northeast and at the foot of the steep edge, the geologically valuable area adjoins the existing AWG 'Maasterrassen'. This landscape is characterised by a relief of 'gullies' and 'flats', remnants of a braided river system from the last ice age.

5. Documentation & Resources

6. Maps, photos and figures

) TNO Intern 68/219



) TNO Intern 69/219

Consider CC 03	Nome
Geosite: GS_03	Name
Significance : National	Bijzonder Brabants
Coordinates (RD): 184676, 384046	
Municipality	Deurne
Duranings and sounting	Neard Dishart
Province and country	Noord-Brabant
Protection status	
1 Total Status	
Site management organizations	Private
Accessibility	Access to the garden is via the barn side. If arriving by
	car, you may park under the trees on the grass by the
	barn, then walk into the garden and enjoy the many
	special vegetables cultivated by farmer Henk and the
	Peel Boundary Fault.
Relevance for tourism	Unforgettable vegetable garden on the Peel Boundary
	Fault. An excellent location to discover the Peel
	Boundary Fault. Tickets can only be purchased in
	advance via the website. Accredited guides from VVV,
	Heemkunde, IVN, Staatsbosbeheer, schools and the
	Geopark organisation may visit the Peel Boundary Fault
	at Bijzonder Brabants and lead group tours.
Relevance for education & information	Vegetable farm Henk Kerkers: Information boards
	explaining the geology of horsts and grabens.
	Observation wells to demonstrate the difference in
	groundwater levels on either side of the fault. There is
	also a very large photographic print of a
	geohydrological fault profile.
Teaching equipment	Lessons can be booked that provide detailed
	information about the Peel Boundary Fault and its
	effects on the landscape and natural environment.
	Educational programmes for schoolchildren.
Scientific Interest	Recent (May 2025) research focuses on the activity of
	the geological fault system in the subsurface of (among
	others) Deurne using the fibre optic network
Description	

1. Location

The vegetable garden Bijzonder Brabants is situated on the Peel Boundary Fault, just outside the village of Deurne.

2. Landscape formation and description

Origin of the landscape

Of the five research locations, this is the site where the impact of tectonics in this area is most clearly demonstrated.

Geomorphology:

Uniquely, the Peel Boundary Fault runs directly through the vegetable garden. This is a geological

) TNO Intern 70/219

fault line that runs from Roermond to Uden in the Netherlands. In the garden, the Peel Boundary Fault and its underground waterfall are made visible by the installation of a metal plate.

Hydrology:

At the fault, a steel plate has been driven into the ground. Above the fault, with its high groundwater level, the soil remains intact, while below the fault, part has been excavated against the fault/steel plate. At the excavation site, you stand below the fault at a lower level and face the fault. The steel plate contains holes through which water flows or seeps. Wells have been installed on both the horst and graben sides, with well rings and an internal measuring scale showing the actual groundwater level. This allows visitors to see for themselves the significant difference in groundwater levels.

Deposits & Fossils:

River & Streams:

3. Geosite Interests

Cultural History


Flora & Fauna:

Around the visible underground waterfall, an informative and educational landscape has been created with plantings characteristic of the Peel Boundary Fault on both the horst and graben sides. In farmer Henk's vegetable garden, around 100 special vegetables and 100 different herbs are grown each year. Farmer Henk cultivates the most remarkable vegetables and herbs as naturally as possible (climate-conscious and sustainable) in the open air..

Industrial & Economic:

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures

) TNO Intern 71/219



) TNO Intern 72/219



The farm Bijzonder Brabants is located on the Peel Boundary Fault (photo: www.landvandepeel.nl)



Groundwater wells illustrate the difference in groundwater levels on either side of the fault (photo: www.landvandepeel.nl); information board in the background.

) TNO Intern 73/219

Geosite: GS_04	Name	
Significance : International	Griendtsveen and surroundings.	
Coordinates (RD): 189748, 383860	, and the second	
Municipality	Horst aan de Maas, Deurne,	
Province and country	Noord-brabant, Limburg	
Protection status	Bechermd Dorpsgezicht	
Site management organizations	Landschap Horst aan de Maas?	
Accessibility		
Relevance for tourism	Toon Kortooms park (peat cutting demonstrations, museum). Starting point for walking and cycling routes through the Peel, such as the "Verheven Peel" route.	
Relevance for education & information	Toon Kortooms museum. Peelmuseum – America, also with peat cutting demonstrations.	
Teaching equipment	Walking route with information boards featuring quotes from Kortooms and explanations about his life and work. "Expedition De Peel" is an interactive activity immersing visitors in the world of Peel workers, combining team building with knowledge about life in the Peel in the past	
Description		

1. Location

Griendtsveen and the Toon Kortooms museum are located on the northern side of the Deurnse and Mariapeel, around the border between Noord-Brabant and Limburg. The protected area covers 87.7 hectares and lies entirely within the province of Limburg. The Peelmuseum – America is situated on the main road from Griendtsveen to America (corner of Griendtsveenseweg and Middenpeelweg) in Limburg and is housed in the former D.U.W. camp. The relatively small Peel remnant, the Grauwveen, lies north of the Helmond–Venlo railway and falls within this geosite.

2. Landscape formation and description

Origin of the landscape

Water was unable to penetrate the poorly permeable subsurfacesoil of the Peelhorst, resulting in the formation of small lakes that became vegetated as the climate warmed after the last Ice Age. This vegetation initiated the formation of low peat. As the peat layer thickened, the groundwater level dropped, making it increasingly difficult for plants to access nutrient-rich groundwater, ultimately ending the formation of raised bog. The high elevation of the Peel also contributed to this process. For a long time, these wet peatlands were difficult to access, but eventually they were reclaimed for peat exploitation. Within the aUGGP, the Deurnsche Peel and Mariapeel represent unexploited areas where remnants of peat moss are still present. Griendtsveen is a reclamation village located on the northern edge of the peatland. The Grauwveen is a small unexploited area just north of Griendtsveen and is one of the few places where the upper peat layer has not been systematically excavated. This layer is known as bonkaarde, pale peat or "grauwveen". In most of the Peel, this layer was excavated in the early 20th century for the production of peat litter. The area has geological value because a relatively complete raised bog profile can still be found here, although the soil has changed due to

) TNO Intern 74/219

desiccation and mineralization.

Geomorphology:

Hydrology:

Griendtsveen belongs to Horst (Limburg) and here the water is pumped up so that it can flow to the Meuse. The drainage of the village of Griendtsveen takes place via the Griendtsveen Kanaal in an easterly direction. This is remarkable because Griendtsveen lies on the west side of the watershed through the Mariapeel, and the natural gradient thus goes westward. The explanation is that this is a kind of 'political drainage'. The village belongs to the Limburg municipality of Horst and the Limburg water board, and therefore the drainage has been artificially directed eastward. This is done by a small pumping station that pumps the water over a weir into the canal. After that, it is high enough to flow eastward naturally

Deposits & Fossils:

In Grauwveen, the raised bog profile has been preserved, but desiccation and mineralization have changed it.

River & Streams:

The Helenavaart between Griendtsveen and Meijel is 15.7 km long and was dug to transport peat from the Peel and also served for drainage of the peatland. The Helenavaart runs between the Mariapeel and the Deurnsche Peel after Helenaveen.

3. Geosite Interests

Cultural History

In 1853, the brothers Van de Griendt purchased 610 hectares of peatland from the municipality of Deurne and started the first industrial peat extraction. The village of Griendtsveen was founded in 1885 by the brothers Eduard and Jozef Griendt. Griendtsveen has a very picturesque location and a protected village view. The history of Griendtsveen is closely linked to the reclamation of the Peel. The Eindhoven–Venlo railway, built in 1866, was crucial for the reclamation of the peatland. The contractor for this railway (from Helmond to Venlo) was Jan van de Griendt (1804–1882), a merchant from Den Bosch. In 1853, he was one of the founders of the "Society for the Reclamation and Peat Extraction of the Peel", later renamed "Society Helenaveen". Van de Griendt founded the village of Helenaveen. His sons Jozef and Eduard continued his work and founded the village of Griendtsveen; both villages are connected by the Helenavaart. Helenaveen is in Noord-Brabant, Griendtsveen in Limburg. In Griendtsveen, the Griendtsveen Peat Litter Company was active.

Over time, all the peat was excavated and the village became residential. It is perhaps the only village in Limburg that has never had an agricultural function. In 1956, the monopoly of the Van de Griendt company ended. With this normalisation, the reclamation of the Peel was completed. The entire village is a monument, as it is a fine example of a Peel reclamation village. Griendtsveen is one of the very few places in Dutch Limburg with a pure Brabant dialect.

The writer Toon Kortooms is buried in Griendtsveen. His father was director of the Deurne municipal peat company, which, along with their family home, was located "just" in the municipality of Deurne, but close to Griendtsveen. His social life therefore took place in the village of Griendtsveen. Kortooms' novels are mainly set in the Brabant Peel and are often based on his own experiences. The Toon Kortooms park includes a museum. The museum has grown on the site where the Kortooms family lived, immediately next to the foundations of the Deurne peat litter factory, which was a competitor

) TNO Intern 75/219

of the similar factory of the Van der Griendt company. In the Peelmuseum, visitors can experience the atmosphere of the hard life and heavy work of the diligent Peel workers during peat extraction and reclamation of the Peel from around 1900. The museum increasingly shows what happened in earlier years in this rugged landscape, and how people lived and worked there.

Flora & Fauna:

The Grauwveen features dry and wet heath, deciduous forest and marsh. There are still small areas with living raised bog and regenerating raised bog in the early stages. Much of the vegetation consists of birch-oak forest and bracken. Scattered small-scale peat excavations, so-called farmers' pits, are moist biotopes where raised bog has a chance to recover. The relatively untouched terrain is a refuge for various animal species. Management is also aimed at favorable conditions for the smooth snake and butterflies such as the large heath and the silver-studded blue.

Industrial & Economic:


Griendtsveen is a typical reclamation village founded when industrial peat extraction began. The industrial method of peat extraction followed a strict pattern, employing hundreds of peat workers. In the peak year 1893, about 700 peat workers worked for the Griendtsveen company, excavating 107 hectares of peat bog, yielding 960,000 m³ of peat.

4. Relation with other geosites

Helenaveen (GS_18) is another peat colony on the south side of the Deurnse and Mariapeel (GS_11).

5. Documentation & Resources

6. Maps, photos and figures



Map showing the location of GS_04 – Griendtsveen and surroundings (blue outline)

) TNO Intern 76/219

Geosite: GS_05	Name	
Significance: International	Vorstengraf en Paalgraven	
Coordinates (RD): 167986, 416079	vorsterigial en radigiaven	
Municipality	Oss	
Province and country	Noord-Brabant	
Protection status	Archaeological monuments (2): Vorstengraf and	
	Paalgraven	
Site management organizations	Municipality of Oss	
Accessibility	Via Zwaardweg in Oss; The Vorstengraf is part of the	
	Brabant Nature Network, an initiative of the province to	
	create more nature reserves.	
Relevance for tourism	Finds from the Vorstengraf are exhibited in the National	
	Museum of Antiquities (Leiden). Finds from the	
	Vorstengraf (mound 7) at Zevenbergen have been	
	transferred to Museum Jan Cunen	
Relevance for education & information	The graves are equipped with information boards.	
	and graves are equipped man members assured	
Teaching equipment		
0.1.10	A	
Scientific interest	Archaeological research: From 1997 to 2005, over 5	
	kilometres of trial trenches were dug in the surrounding	
	area.	
Description		

1. Location

The archaeological monuments Vorstengraf Oss and De Zevenbergen are two areas with burial mounds, urn fields, and flat graves from the Dutch Bronze Age. The Vorstengraf near Oss is the largest burial mound in the Netherlands. The two areas are separated by a motorway and are 400 meters apart, but are accessible via a pedestrian bridge. The Paalgraven are also located to the east of the motorway.

2. Landscape formation and description

Origin of the landscape

The princely grave at Oss dates from the early Iron Age, with finds from around 700 BC. The burial mound was built on an older burial mound from the Bronze Age (circa 1800 BC). It is the largest burial mound in the Netherlands and originally had a height of about 3 meters and a diameter of 54 meters. A wealthy person of great esteem was buried there, as evidenced by the rich grave goods.

In 1932, a bronze bucket was discovered on the former Osse Heide while leveling a large hill, containing the cremated remains of the 'prince of Oss' and the grave goods that were given to the deceased. The most famous is the ritual around the bent sword. Swords from the Iron Age are a rarity. In addition, a knife, horse bits and harness, an axe, a whetstone, two razors, jewelry, and clothing were also found.

During the construction of the A50 near Zevenbergen, a second princely grave was found. In the vicinity of the princely graves, burial mounds with post circles (paalgraven) from the early Iron Age

) TNO Intern 77/219

were excavated. Some have been reconstructed, and many are located at the traffic junction aptly named 'Paalgraven'..

Geomorphology:

The area around the Vorstengraf lies on the Peel Boundary Fault fault and descends from 26 meters to 11 meters within a distance of a few hundred meters. The Vorstengraf is situated in a relatively high, well-visible location: precisely where the Maashorst, an extension of the Peelhorst, transitions into the lower terrain of the Maaskant. This last relief was formed by the incision of the Meuse.

Hydrology:

Deposits & Fossils:

River & Streams:

3. Geosite Interests

Cultural History

During the period from the early Bronze Age to the early Iron Age (circa 2000 to 700 BC), there was a notable concentration of habitation on the northwestern edge of the Peelhorst, to the south of Oss. In prehistoric times, the existing elevation differences were utilized here. For those coming from the low-lying clay area in the north, the higher Peelhorst must have been a striking feature. At Vorsel and Zevenbergen, there were several urn fields and many burial mounds on the high edge, including the well-known princely graves of Oss. Burial mounds also served as markers of the 'property rights' of a local community. Therefore, they had to be conspicuous and were erected in prominent locations. Research has shown that the burial mounds around the princely grave of Oss were grazed by sheep at that time. This prevented the growth of shrubs that could obscure the burial mounds from view. Consequently, and due to their high location, the mounds were visible from a great distance (Jansen and Van der Laan, 2011).

Flora & Fauna:

In the area, natural processes, such as the decay of dead trees, are not actively countered. This offers opportunities for the development of diverse ecosystems. The province of Noord-Brabant aims to acquire agricultural lands and convert them into nature within the Brabant Nature Network. This process will proceed gradually, focusing on the preservation of archaeological heritage and the promotion of biodiversity.

Industrial & Economic:


4. Relation with other geosites

Slightly further south on the border of Uden and Nistelrode, the grave of the Prince(ss) of the Maashorst was found in 2005. Burial mounds and finds were also made in the area where the Peelhorst crosses the Meuse valley, for example in Baarlo (De Bong) and in Swalmen (Bosberg). Here too, the burial mounds are located on higher edges.


5. Documentation & Resources

Jansen and Van der Laan, 2011

) TNO Intern 78/219



Map with the location of GS_05 – Vorstengraf and Paalgraven (blue outlined)



Cartoon map of the geosite GS_05 with the location of the Vorstengraf and the Paalgraven (source: Liselore Burgmans)

) TNO Intern 79/219



The monuments Vorstengraf (left) and Paalgraven (right). By Watisfictie (Own work, CC BY-SA 3.0) and Encyacht (Own work, Public domain,

https://commons.wikimedia.org/w/index.php?curid=8876351)

) TNO Intern 80/219

Geosite: GS_06	Name
Significance: International	Peelraamstelling
Coordinates (RD): 182518, 401913	
Municipality	Land van Cuijk
Province and country	Noord Brabant
Protection status	Rijksmonument
Site management organizations	
Accessibility	Roijendijk, Mill , Landhorst
Relevance for tourism	High experiential value
Relevance for education & information	
Teaching equipment	

1. Location

The Peel-Raamstelling was a Dutch defensive line constructed in 1939 to halt or delay a potential German attack. On 10 May 1940, the first day of the Dutch war, the Peel-Raamstelling fell. The line was situated behind the Meuse Line, varying from 9 km to 21 km apart. It began at the Meuse near Grave, ran via Mill, through the Peel, and along the Zuid-Willemsvaart to the Belgian border near Weert. The Asperge Monument still commemorates one of the first German acts of war on Dutch soil. In the Land van Cuijk, between Grave and Landhorst, the remnants of the Peel-Raamstelling are still clearly visible:

- Stadsmuseum Grave
- The route of the former "Duitse Lijntje (German Railroad)
- Asperge Monument in Mill
- Waterweetjeswandeling in Landhorst
- Guided tour by Stichting Sporen van de Oorlog in Mill

2. Landscape formation and description

Origin of the landscape

The geology of the Peelhorst directly influenced the choice of location and nature of the defences, thus having strategic value. Between Meijel and De Rips, especially around Griendtsveen, the line benefited from natural protection by the marshy peat bogs of the Peel and several existing waters, such as the Graafse Raam, Helenavaart, and Noordervaart. Along the northern part, an artificial barrier was created in the form of the Defensiekanaal. Nearly 400 casemates were also constructed, along with barbed wire obstacles and clear fields of fire.

Geomor	<u>pho</u>	logy:

<u>H'</u>	yc	ro	log	y :

) TNO Intern 81/219

Deposits & Fossils:

River & Streams:

The line could, between Meijel and De Rips, especially around Griendtsveen, benefit from natural protection by the marshy peat bogs of the Peel and several existing waters, such as the Graafse Raam, Helenavaart, and Noordervaart. Along the northern part, an artificial barrier was created in the form of the Defensiekanaal.

3. Geosite Interests

Cultural History

The Peel-Raamstelling remains largely intact and is still very recognisable in the landscape of the northern Peel due to a defensive canal with many casemates. Notable sections include the area between Griendtsveen and Vliegbasis de Peel (especially within the Heidsche Peel) and the site at Mill where the line crossed the so-called "Duitse Lijntje". Here stands a monument commemorating the famous "armoured train incident". The monument consists of a pair of bent rails near the bridge, with the infamous "asparagus" obstacles (the actual derailment site is several dozen metres westwards), as well as the artillery monument, the restored casemate, and the monument 1-3RI. Casemates in the municipalities of Mill and Sint Hubert, Venray, Horst aan de Maas, and Deurne are protected as national monuments.

Flora & Fauna:


Industrial & Economic:

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures



Map showing the location of GS_06 – Peel-Raamstelling (blue outline). The geosite represents

) TNO Intern 82/219



) TNO Intern 83/219

Geosite: GS_07	Name
Significance: International	Sint Annabos
Coordinates (RD): 168052, 409679	
Municipality	Maashorst / Bernheze
Province and country	Noord-Brabant
Protection status	The wijst area between Uden and Vorstenbosch was declared a geological monument in 2004. The area is one of five wijst reserves and has the status of "natte natuur parel"
Site management organizations	Staatsbosbeheer
Accessibility	Via Schansweg no. 10 in Uden, with a large car park.
Relevance for tourism	Circular walk along various phenomena related to the Peel Boundary Fault.
Relevance for education & information	Many manifestations in the Annabos are explained on information boards. Excursion opportunities. For education about Sint Annabos, one can contact IVN Uden, which organises nature activities and walks in the area.
Teaching equipment	
Scientific interest	Research trench (2018) at Lageburchtweg by VU to study various properties of the Peel Boundary Fault. The research in the trench examined fault structures, fault displacement, the age of displaced deposits, and variation in fault impermeability. Samples were also taken for geohydrological laboratory research.

1. Location

The nature reserve "de Wijstgronden Uden" consists of three wet, seepage-rich sub-areas along the road from Uden to Vorstenbosch in Noord-Brabant. These are the Loose Beemden, the Raktse Beemden, and Kooldert, together covering about 41 hectares. The well-known Sint Annabos, west of Uden, is part of the (sub)area the Raktse Beemden and is the place where the Peel Boundary Fault is most visible through a great variety of manifestations. The wijst grounds of the Loose Beemden are similar and lie south of the Karperdijk.

2. Landscape formation and description

Origin of the landscape

Due to the poor water transmissibility of the faults, a narrow, wet zone has formed along the faults, the wijst grounds, with a unique variation in wetness and groundwater composition for the Netherlands. As a result, a great diversity of plants can be found in the narrow strip along the fault. Characteristic species from twenty plant communities grow on these wijst grounds. The water that rises along the faults is rich in iron.

Geomorphology:

) TNO Intern 84/219

Sint Anna Bos is the place where the Peel Boundary Fault is most visible, as the landscape suddenly rises about three metres via a steep slope. The height difference is bridged by following the boardwalk with the wooden bridge, where the landscape suddenly rises about three metres. Sometimes the fault itself is visible at the surface. In the asphalt of the biking lane on the Karperdijk in Uden (GS_07e), which lies over the Peel Boundary Fault, there is a crack of two centimetres, showing that the fault is still active. In the vicinity of faults, cracks in the asphalt are often seen.

Hydrology:

The Sint Annabos (GS_07d) is a moist deciduous forest, rich in alders, ferns, and mosses. It is a wijst area where wijst flora can be seen along the deep ditches through which the seepage water on the high side of the fault flows towards the graben on the west side. The rusty brown colour shows that the water in the seepage ditches is rich in iron. The Revennekes (GS_07b) is a created nature area and lies exactly on the Peel Boundary Fault. The fault line is not directly visible in the landscape but can be recognised by the rusty brown water in the ditch (next to the bike lane). The height difference over the fault can also be seen at the large drop at the two weirs in the deep watercourse.

Deposits & Fossils:

The area west of the fault is characterised by several sand drifts, including the Achter de Berg Nature Reserve (GS_07a) and Bedafse Bergen (GS_07f). In the Middle Ages, drifting sand ridges formed here. The drifting sand ridge, up to 25 metres high, is well maintained.

River & Streams:

Various seepage ditches east of the fault with marsh flora.

3. Geosite Interests

Cultural History

The wijst grounds in this area were discovered through flora and fauna research in connection with the construction of the A50 route through the area. The wijst grounds have been designated as a Geological Monument and the route of the motorway was diverted.

Flora & Fauna:


The Raktse Beemden (GS_07c) form an elongated moist grassland area at the transition from dry to wet, with meadows, hedgerows, and ditches. The rising seepage water makes the area a hotspot for special plants and animals. Beautiful species such as marsh marigold and bistort bloom here. Marsh birds and amphibians such as the moor frog thrive here. In the Revennekes, there are characteristic tree species and moisture-loving vegetation typical of this relatively wet nature area. On the other side of the cycle path is a hedgerow mainly of downy birch, which thrives here. This tree species is characteristic of wijst grounds.

Industrial & Economic:


In the last century, much drifting sand was stabilised with pine planting. The wood was mainly used in coal mining as pit props.

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures

) TNO Intern 85/219



Map showing the location of GS_07 – Sint Annabos (blue outline) and surroundings along the Peel Boundary Fault (in red)



AHN image of the geosite Sint Annabos showing the height difference over the Peel Boundary Fault. The morphology of the drifting dunes is clearly visible west of the Peel Boundary Fault in the Achter de Berg Nature Reserve (GS_07a, top of the image) and the Bedafse Bergen (GS_07f, bottom of the image)

) TNO Intern 86/219



Wijst water, recognisable by its typical rusty brown colour, flows over the Peel Boundary Fault near Uden (source: "Breuken in het land van Maas en Waal", Van Balen et al., 2022)



Distinct vegetation boundary on the steep edge of the Peel Boundary Fault.(Source: "Ecologische iconen van de Wijst", Ettema, 2010.)

) TNO Intern 87/219

Geosite: GS_08 Significance: International Coordinates (RD): 185730, 382105	Name Hoogdonk (Den Brink)
Municipality	Deurne
Province and country	Noord-Brabant (NL)
Protection status	Natuurmonument
Site management organizations	Xella Nederland
Accessibility	Various roads run around the lake, but the site itself is not accessible.
Relevance for tourism	Walking routes along and around the Den Brink lake
Relevance for education & information	Fossil finds can be viewed at the Klok and Peel Museum in Asten.
Teaching equipment	
Scientific interest	The site is recognized wordwide as a fossil site for the Miocene and Pliocene

1. Location

The Hoogdonk (Den Brink) sand extraction site is located north of the village of Liessel at the end of Hoogdonkseweg, at the intersection of Snoerstebaan/Berktsedijk. The lake is about 1 km long and up to 50 m wide.

2. Landscape formation and description

Origin of the landscape

The Hoogdonk sand extraction site is situated on the Peelhorst, just east of the Peel Boundary Fault. Due to the uplift of the Peelhorst and erosion of younger sediments, Pleistocene, Pliocene, and Miocene sands are relatively close to the surface here. This explains why so many fossils from the late Pliocene and Miocene have been found at this sand extraction site (up to about 30 m deep)..

Geomorphology:

The sand extraction site is bounded on the west by the Peel Boundary Fault, which is marked by a small terrain step. More obvious, however, are the differences in vegetation on either side of the fault. At the north-western boundary of the sand pit along the Berktsedijk, the Peel Boundary Fault can be experienced at the Oude Aa. The height difference is evident from the closely spaced weirs in the Oude Aa.

Hydrology:

The lake is fed by rain and seepage water and, to preserve the fish population, is not in open connection with surrounding water. As Den Brink is mostly deeper than four metres, it can be considered a deep-water type lake. Since September 1968, the lake has been used as a fishing pond by the De Peel Angling Association.

) TNO Intern 88/219

Deposits & Fossils:

Over time, numerous palaeontological finds have been made at the sand extraction site up to 2002. Hoogdonk worked with two museums, the Klok and Peel Museum in Asten and the Oertijdmuseum De Groene Poort in Boxtel. Groups of amateur palaeontologists regularly searched for special finds in the dredged sand together. Shark teeth, whale vertebrae, bird remains, sea urchins, and starfish have been found. Whale remains and many other marine fossils are evidence of a time when this part of the Netherlands was sea (Miocene, Breda Subgroup). Remains of mastodons and other fossils date from a period when the area was land (Pliocene and Pleistocene, Kiezeloolite and Beegden formations, respectively). Occasionally, larger erratic boulders are found in the Pleistocene river deposits. In Liessel, for example, the "Jaap van de Leijsing" and the "Buntse Kei" are on the market square. Seeds of plants and artefacts indicating human presence have also been found, so to some extent, archaeological finds have also been made. Most of the dredged fossils were made available by director Van den Brink to the Klok and Peel Museum in Asten, where the palaeontology department displays a large and beautiful collection of the many fossils. After 2002, the dredging method changed, ending palaeontological research at the sand-lime brick factory. Initially, dredging was done with a floating crane ship. Due to changing environmental requirements, this was done after 2002 with a hired electric suction dredger...

River & Streams:

The Oude Aa stream runs along the north side of the excavation over the Peel Boundary Fault. The Oude Aa is also known as the Vlierdense Aa, Vloeieinds(ch)e (Water)Loop and Raktse Loop and originates in the Deurnese Peel.

3. Geosite Interests

Cultural History

Many residents of the municipality of Deurne, mainly from the Sint-Jozef parish (district) and Liessel, worked in the production at this factory, often in two shifts (the stokers in three shifts) at Den Brink, as the factory was soon popularly called. Den Brink is also the name of the lake created by sand extraction.

Flora & Fauna:

The flora and fauna are extremely varied due to the alternation of wet and dry areas. The area contains flora characteristic of wijst areas, such as Dwarf Hair Grass. Other less common species that prefer base-rich seepage include brown flatsedge, pennyroyal, shortawn fox-tail, and the Cavernous Crystalwort

Industrial & Economic:

Two farmers sold 15 hectares of farmland to Gijberrtus Abtonius (Bert) van den Brink and 20 hectares of peatland to the municipality of Deurne, and in 1954 the Hoogdonk Sand-Lime Brick Factory was established in Liessel. The Peel Boundary Fault lies in front of the factory gate. On the factory site, Miocene (?) sand is at the surface, making it easy to extract. In 2000, a permit was obtained to expand the lake, allowing the factory to continue production until about 2030. Sand extraction was stopped in 2022, but Xella Kalkzandsteen resumed sand extraction for sand-lime brick products in 2024. The permit for excavation runs until 2025; further plans are unknown.

4. Relation with other geosites

The Hoogdonk sand extraction site has no direct relationship with nearby geosites. In De Kuilen (GS_20), forty kilometres north of Liessel, fossils from the Miocene and Pliocene have been found in a sand extraction pit. This location is also on the Peelhorst.

) TNO Intern 89/219


5. Documentation & Resources

- Peters, A., Lammers, Th.& Mol, D. (1991). Mastodonten=kiezen uit Liessel (Noord-Brabant). Cranium 8 (2), 89-96.
- Peters, N., (2009), Brabant tussen walvissen en mastodonten, Fossielen uit Liessel [Asten/Boxtel]

6. Maps, photos and figures




Map showing the location of GS_08 – Hoogdonk-Den Brink (blue outline) on the high fault block just east of the Peel Boundary Fault (in red)



AHN image of the geosite GS_08 — Hoogdonk-Den Brink showing the height difference over the Peel Boundary Fault. The higher area at the bottom of the image lies west of the fault and represents a drifting sand area.

) TNO Intern 90/219



) TNO Intern 91/219

Geosite: GS 09	Name
Significance: National	Leudal
Coordinates (RD): 193927, 362351	Leadai
Municipality	Leudal
Province and country	Limburg
Protection status	Natura-2000
Site management organizations	Staatsbosbeheer
Accessibility	Het Leudalmuseum, Roggelse weg
	. 55
Relevance for tourism	Leudal is a very popular walking area and is part of the
	Hertogenpad, but there are also many shorter walking
	routes; these start near the Leudalmuseum. There are
	various cycle paths as part of the cycling Hroute
	network.
Relevance for education & information	The Leudalmuseum, located in the Sint Elisabethshof.
	This museum provides information about the flora and
	fauna, archaeology, local history, and some history
	(especially the Second World War) of the area
Teaching equipment	(copedian) the sessing trong that the drea
reaching equipment	

1. Location

Het Leudal is a nature reserve in the Dutch province of Limburg. It is located between the village centres of Haelen, Roggel, Neer, Nunhem, and Heythuysen in the municipality of Leudal. The total nature reserve covers about 900 hectares, a large part of which is managed as a reserve by Staatsbosbeheer. The area is particularly known for its stream valleys. The geosite consists of an area of 315 hectares designated as a Natura 2000 area.

2. Landscape formation and description

Origin of the landscape

Het Leudal is a relatively high-lying cover sand area on the west bank of the Meuse, with deposits from the last ice age. There are stream valleys, land dunes, and forest ponds. Due to its location in a landscape with Meuse terraces and fairly large height differences, the streams flow quickly, and the stream valleys are strikingly deeply incised into the sand.

Geomorphology:

The stream valleys were formed in the last ice age and have been further eroded over the centuries. Another clearly visible remnant from the distant past are the metre-high sand dunes. In combination with the deep, almost ravine-like stream valleys, they create significant height differences.

Hydrology:

Due to the height differences in Het Leudal, there are large differences in groundwater levels. Together with the differences in soil composition, this creates a great diversity in vegetation. Groundwater steps are found locally in the stream valleys. This is due to peat layers in the subsoil,

) TNO Intern 92/219

which prevent the groundwater from sinking away. Clay layers can cause wet spots locally. There are also several diffuse seepage zones in old meander bends, in the stream valley of the Roggelsebeek on the side of the high hinterland.

Deposits & Fossils:

River & Streams:

The Tungelroyse Beek (locally also called Leubeek) and the Zelsterbeek flow through the area. Between Nunhem and Neer, they merge into the Neerbeek. Just a few hundred meters further, the Haelense Beek joins them, which runs for kilometres just south and east of the reserve. The Leubeek was largely canalised in the past. But at the beginning of the 21st century, an important part of the old stream course with its meanders was restored. Upstream of Het Leudal, various stream sections have been or are being restored, particularly parts of the Tungelroyse Beek and the Roggelse Beek.

3. Geosite Interests

Cultural History

In the middle of the area lies the Sint-Ursulamolen or Leumolen from 1773 (named after Saint Ursula). It now is a water mill and was once used for grain and oil.

Flora & Fauna:

The vegetation in the stream valleys is very varied, especially in the deepest valleys due to the seepage and the associated vegetation. The wet forests in the deep stream valleys belong to relatively special forest types (alder carr, bird cherry-ash forest, hornbeam forest, and birch carr). They mainly contain species that indicate deep seepage, such as large amounts of golden saxifrage, marsh marigold, and water violet. In pools and marshes, further grow snake-root, alder sedge, stiff sedge, bog myrtle thickets, and adder's tongue. In relatively nutrient-poor places in the stream valley, slender primrose, wood anemone, Solomon's seal, and lily of the valley bloom in spring. East of the Elizabeth Monastery are poor grasslands with, among others, spotted orchid. The higher grounds also form a varied nature reserve with forest (consisting of deciduous and coniferous trees), small heathlands, and dry grasslands. There are also fields and meadows.

The fauna is very varied. In addition to fox and roe deer, there are badgers and other mustelids; a group of wild boars roams in the area. Beavers were released in 2002. Special bird species include black woodpecker, green woodpecker, middle spotted woodpecker, bullfinch, grey wagtail, sparrowhawk, and hobby. Moreover, kingfishers can be found, which have an ideal habitat in the area with steep sandbanks and fish-rich streams. The phegea moth occurs in the area.

Industrial & Economic:

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures

) TNO Intern 93/219



Map showing the location of GS_09 – Leudal (blue outlined) located on the low-lying fault block southwest of the Peel Boundary Fault (in red). The Leubeek and Zelsterbeek (white arrows) drain to the northeast. After the confluence of the streams, they flow further as the Neerbeek towards the Peel Boundary Fault, which they then cross at Neer.



The Leubeek in the nature reserve Het Leudal (source: Wikipedia)

) TNO Intern 94/219



The Sint-Ursulamolen in Het Leudal (source: Wikipedia)

) TNO Intern 95/219

Geosite: GS_10	Name
Significance: International	Maasterrassen
Coordinates (RD): 201809, 367049	
Municipality	Leudel, Beesel en Venlo
Province and country	Limburg
Protection status	Aardkundig waardevol gebied
Site management organizations	
Accessibility	
Relevance for tourism	
Relevance for education & information	
Teaching equipment	

1. Location

The most accentuated Meuse terraces lie roughly between Roermond (Swalmdal) and Tegelen, characterising the area where the River Meuse crosses the Peelhorst..

2. Landscape formation and description

Origin of the landscape

Where the area of the Meuse south of Roermond is a geological subsidence area (Roer Valley Graben), North Limburg (including the Peel Horst, but also the Venlo Graben) is actually being pushed further upwards. Near the village of Neer, the uplift currently amounts to about 2 cm per century (almost half a metre since Roman times, 2 metres since the beginning of the Holocene). This is evidenced by the earthquakes that occasionally occur around the Peel Boundary Fault near Roermond.

The uplift of Northern Limburg automatically meant that the Meuse had to start incising. In warm climate periods, this incision process proceeded relatively quickly, only to partly come to a halt again during ice ages and cold periods. During these glacial periods, the Meuse flowed as a braided river through an almost treeless landscape, carrying much sand and gravel. The old, braided channel relics on the lower terrace date from one of the last cold periods.

Due to climate fluctuations and river course shifts in the upstream catchment, the Meuse has experienced large differences in discharge over time. For example, until about 200,000 years ago (Saalian), the Meuse still drained the water of the Moselle, which, among other things, allowed the large meander bends on the middle terrace of the Schuytwater and Kaldenbroek to form. During glacial periods, sea level was much lower than today, and the North Sea coast lay relatively far away. During warm periods, the coast could advance almost as far as Limburg. The braided Meuse disappeared and the river took on more the character of a more sedate delta river, sometimes with broad meander bends, but sometimes also with its current narrow 'canyon bed'...

) TNO Intern 96/219

Geomorphology:

The alternation of cold and warmer periods, and thus of incision and widening of the Meuse, ultimately led to the formation of the well-known Meuse terraces, which are visible everywhere in the area like steps. Where the Meuse eroded close to the old middle or high terrace, high steep banks arose directly at the water, such as at Neer, Kessel and at the monastery village of Steijl, where a height difference of 15 to 20 metres between river and middle terrace can be observed.

Hydrology:

Instead of flooding by river water, the dominant process along this river stretch is the emergence of iron-rich groundwater. This is manifested in shallow seepage channels scattered throughout the area at various terrace levels, often far above the Meuse. These channels collect the clear groundwater from older, adjacent areas. Most channel relics are in fact 'fossil channels', no longer directly related to the present Meuse.

Deposits & Fossils:

River & Streams:

Present-day Meuse and numerous side streams

3. Geosite Interests

Cultural History

The old structure of rising river terraces still determines the use and spatial development of the Limburg Meuse valley to this day. Old Meuse villages such as Meerlo, Grubbenvorst and Arcen are still recognisable as historic bastions on the high terrace edges. These were also the logical settlement sites of the first inhabitants of the Netherlands. Right along the Meuse, we find remains from the Stone Age, Bronze Age, Iron Age, Roman period and Middle Ages, directly on top of each other. Because the Meuse has hardly shifted its position for 10,000 years, it has never 'cleared away' this old cultural heritage.

Flora & Fauna:

Due to the action of iron-rich groundwater, the channels and old Meuse arms (potentially) have a unique plant growth with species such as Marsh Marigold, Water Horsetail and even Golden Saxifrage. Where rich alder carr forests grow in the seepage areas, such as in the Old Meuse Arm near Meerlo and the Heuloërbroek near Aijen, we find Nightingale, Pied Flycatcher and sometimes even Golden Oriole. The European Tree Frog, Purple Emperor and Garlic Toad probably originally occurred in large numbers in and around the wet groundwater channels of the Meuse valley.

Industrial & Economic:


4. Relation with other geosites

5. Documentation & Resourcess


- https://www.limburgs-landschap.nl/wp-content/uploads/2021/04/2019 Maasterrassen Ir.pdf
- Woolderink, H.A.G, C. Kasse en R.T. van Balen (2020), De invloed van breuken op de Maas en Roer in het Roerdalslenk-systeem.' in: Grondboor en Hamer 74, 176-182
- Woolderink, H. A. G. (2021). Faulty Rivers: The effect of faulting on river morphodynamics and morphology. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam]. Ipskamp.

) TNO Intern 97/219

6. Maps, photos and figures




The Meuse crossing (GS_10) represents the area (outlined in blue) where the Meuse has cut through the Peel Horst between the Peel Boundary Fault and Tegelen Fault.



AHN image (shaded relief) of the Meuse crossing shows height differences due to Meuse terraces.

) TNO Intern 98/219



Palaeogeographical reconstruction of the Meuse terraces as observed in the AHN image. (source Woolderink et al., 2020)



Old terrace edges and incising streams as a unique feature of the geomorphology of the Meuse terraces. Source: www.limburgs-landschap.nl

) TNO Intern 99/219

Geosite: GS_11	Name
Significance: international	Deurnse- en Mariapeel
Coordinates (RD): 190423, 381305	
Municipality	Deurne, Horst aan de Maas
Province and country	Noord-Brabant, Limburg (NL)
Protection status	Natura-2000. UNESCO wetlands
Site management organizations	Staatsbosbeheer
Accessibility	Various nature "gateways"
Relevance for tourism	Walking and cycling routes
Relevance for education & information	Guided tours by guides (starting from, among others, nature gateway De Peel)
Teaching equipment	

1. Location

The area covers more than 2,700 hectares and consists of two sub-areas: Deurnsche (Deurnse, Deurnese or Deurnesche) Peel and the Mariapeel. Together with the nearby Groote Peel, they are remnants of what was once an extensive primeval landscape of living raised bog. The reserve is completely surrounded by agricultural land.

2. Landscape formation and description

Origin of the landscape

During the last part of the last ice age, the Peel landscape was sparsely vegetated for a long time. Much cover sand was blown away in places where small rivers cut through sandy sediments from the subsurface. Due to the predominantly south-westerly winds, the upper reaches of the Brabant rivers, which flow northwest, struggled. Entire cover sand ridges were blown straight into the valley, leaving completely or largely cut-off basins. The Limburg rivers were much less affected, as they flowed in the same direction as the drifting cover sand. The Brabant rivers had yet another handicap. The Peel Horst as a whole is tilting, with the north-eastern half subsiding and the southwestern half rising. The upper reaches of the Brabant rivers therefore subsided relative to the middle and lower reaches in the Roer Valley Graben, reducing gradient and erosive power. Due to tectonics and the drifting in of cover sand, large, more or less closed basins eventually formed in the Graben of Griendtsveen and in the sanded-up upper reaches of Brabant rivers such as the Soeloop and the Vlier. In all these basins, fen peat began to form during the last part of the ice age. During the Holocene, transitional mires developed on this, and eventually raised bog formed, which also spread outside the original basins and completely or largely overgrew surrounding cover sand ridges. Locally, elongated chains of peat lakes developed between a raised bog and a cover sand ridge or between two approaching raised bogs, such as the Negenmeren near Griendtsveen. For a long time, these wet peat areas were difficult to access, but eventually they were reclaimed for peat extraction.

) TNO Intern 100/219

Geomorphology:

Hydrology:

The watershed through the area means that the Deurnese Peel and Helenaveen drain to the west and Griendtsveen and Mariapeel to the east. Due to the location of the peat area in a fault zone, buffering of fens just above these faults is possible.

Deposits & Fossils:

Within the aUGGP, the Deurnsche Peel and Mariapeel represent undeveloped areas where remains of peat moss are still present. The entire Natura 2000 area Deurnsche Peel & Mariapeel is a more or less coherent remnant of the largely disappeared raised bog area De Peel, on the watershed between the Brabant stream the Aa in the west and the Limburg Grote Molenbeek in the east. Most of the peat has been excavated in the past, but locally thick layers of residual peat are still present. In some complexes of farm pits, there are vegetations with hummock-forming raised bog species. These Peel raised bogs were largely excavated down to the sandy subsoil. These areas are the southernmost representatives of the flat subatlantic raised bogs, which elsewhere and also in the Peel region have largely disappeared due to excavation, reclamation and peat cutting.

River & Streams:

3. Geosite Interests

Cultural History

For at least seven centuries, but probably much longer, the peat of the Deurnsche Peel and the Mariapeel was gradually excavated from the surrounding Peel villages. The oldest peat pits of the Verheven Peel are the Ronde Kuilen in the Liesselse Peel, on the western edge of the Deurnsche Peel. On the Limburg side, the peat diggers from Horst, Sevenum and Maasbree generally only reached the Verheven Peel in the course of the 19th century, so the peat pits there are somewhat younger. In the Deurnsche Peel, peat was extracted until the 1970s, and the traces of this are still clearly visible

Flora & Fauna:

Due to the different peat extraction histories of the parts of the area, there is a large and fine-scale variation in vegetation and landscape, with gradients towards a slightly more mineral-rich environment. In the oldest peat pits, raised bog growth has been present on a miniature scale for a long time. On the large residual peat units, a relatively large peat thickness is still present, on which, through restoration management, the development of raised bog vegetation is now also taking place in various places.

The faunal value of the affected raised bog mainly concerns the herpetofauna and avifauna. Herpetofauna: Smooth snake, Moor frog. Avifauna: various marsh birds (Bluethroat, Stonechat), Black-necked Grebe, Teal and Black-headed Gull, Bittern, Little Grebe, Water Rail, Spotted Crake, Cormorant and various ducks. A rare breeding bird is the Nightjar. In addition to the large numbers of cranes seen during migration, the Mariapeel is somewhat renowned as a stopover for the rare Black Stork.

Industrial & Economic:

In the Deurnsche Peel, peat was extracted until the 1970s, and the traces of this are still clearly visible.

) TNO Intern 101/219

4. Relation with other geosites

De Bult (GS_30) is also a raised bog area that has remained after reclamation and is part of the same Natura 2000 area, but is located separately. There is also a relationship with the peat reclamation geosites Griendtsveen (GS_04) and Helenaveen (GS_18).


5. Documentation & Resources

6. Maps, photos and figures



Location of geosite GS_011 – Deurnese and Mariapeel (outlined in blue) situated on the elevated Peel Horst just east of the Peel Boundary Fault.

) TNO Intern 102/219



AHN image (shaded relief) of the Deurnese and Mariapeel (outlined in blue). The image clearly shows the elongated traces of reclamation.



Geomorphological map with the areas where the peat remains are located shown in purple. The watershed here runs almost parallel to the provincial border between Noord Brabant and Limburg (black line), which is also clearly reflected in the opposite flow directions of the stream valleys (in dark green) on either side of the raised bog area.

) TNO Intern 103/219

Geosite: GS_12 Significance: National Coordinates (RD): 176654, 413192	Name Waterzuiverende Wijstboerderij Zeeland
Municipality	Maashorst
Province and country	Noord-Brabant
Protection status	
Site management organizations	Private: Fam. Bongers-Sepp
Accessibility	Langenboomseweg 103, Zeeland
Relevance for tourism	Several walking and cycling routes pass by this location (including the "Langs de Breuk) cycle route)
Relevance for education & information	There are plans for a visitor centre in the old manure silo
Teaching equipment	

1. Location

The Bongers-Sepp family are realising their dream at Heihorst in Zeeland: a dairy farm that has transitioned to nature-inclusive, organic farming. Father Piet and daughter Karen have recently dedicated 35 hectares of their land to nature and wijst restoration. As a result, the farm now and in the future contributes to strengthening the landscape, nature, and especially the water system. One of the most important steps has been the planting of around 20,000 trees on these 35 hectares. This is creating a place where people and animals can once again live in harmony.

2. Landscape formation and description

Origin of the landscape

Geomorphology:

Hydrology:

The land has been designed for optimal water retention and purification of water that isn't needed for the crops (which is the majority). There is a 1.5 km long strip of willows; broad strips of nut trees combined with other productive species, and hedges around the fields. For this, the farmers received compensation from the Brabant Green Development Fund

Deposits & Fossils:

River & Streams:

3. Geosite Interests

Cultural History

) TNO Intern 104/219

Flora & Fauna:

Industrial & Economic:

Alongside space for water and nature, agriculture remains important. Grazing livestock now provides income, as does arable farming with crops such as summer wheat with peas, silage maize, spring barley, and species-rich grassland. The crops are organic, which works well with the planted trees and shrubs that will start to yield produce in a few years' time. There are also plans for a visitor centre in the old manure silo, so that other farmers can get to know the farm and draw inspiration from the transition the family has made.


4. Relation with other geosites

GS_35 De Hoge Raam borders this geosite to the north.

5. Documentation & Resources

Website: https://werkendlandschap.nl/project/waterzuiverende-wijstboerderij

6. Maps, photos and figures



Location of geosite GS_012 – Water-Purifying Wijst Farm Zeeland (outlined in blue), situated around the northernmost fault segment of the Peel Boundary Fault (see inset map for location).

) TNO Intern 105/219



Aerial photograph of the water-purifying wijst farm (view looking east). The Peel Boundary Fault and differences in vegetation and moisture regime are indicated. Source: werkendlandschap.nl

) TNO Intern 106/219

Geosite: GS_13	Name
Significance: National	Slabroek
Coordinates (RD): 169700, 412016	
Municipality	Maashorst
Province and country	Noord-Brabant, The Netherlands
Protection status	Deels Wijstreservaat status
Site management organizations	Staatsbosbeheer
Accessibility	Well connected by a network of walking and cycling paths
Relevance for tourism	A range of activities are organised in this area from the De Maashorst Nature Centre (GS_23). In addition, several local businesses cater to tourists, offering accommodation and selling regional products.
Relevance for education & information	There's an information panel in the area, and extensive information can also be found at the nearby De Maashorst Nature Centre (GS_23).
Teaching equipment	Education team at De Maashorst Nature Centre

1. Location

This small-scale natural and cultural landscape is situated on the Peelhorst, near the hamlet of Slabroek. It comprises a shallow stream valley and adjacent heathland and drifting sand area. In the stream valley lies one of the five Wijst Restoration Projects designated by the province of Noord Brabant.

2. Landscape formation and description

Origin of the landscape

This geosite is located on the Peelhorst. It features a broad, shallow stream valley (Venloop) and, to the south, heathland and drifting sand. The area is intersected by two southeast-northwest trending branches of the Peel Boundary Fault (the Mellebreuk and the Hoevense Fault), which are not visible at ground level but do cause local wijst phenomena in the valley. The valley formed during the last ice age, when rain and meltwater, unable to infiltrate due to permafrost, ran off over the surface, carrying some of the sandy sediment with it. The present-day stream has a much smaller discharge than would be expected for the width of the valley. The heathland is a cultural landscape, created by overgrazing and soil erosion in the Middle Ages. Blowing sand formed elongated drifting sand ridges on either side of the area.

Geomorphology:

This area has hardly been subject to land consolidation, so the original relief has largely been preserved. The shallow stream valley, the cover sand relief and the drifting sand relief are therefore still clearly visible.

Hydrology:

Due to the faults, the wijst phenomenon also occurs here: clean, iron- and mineral-rich, low-acidity groundwater is pushed up to the surface. However, because of large-scale lowering of the

) TNO Intern 107/219

groundwater table, the seepage water now only appears locally at the surface ('wijst'). Efforts are being made to retain water better in the area by constructing small weirs in drainage ditches. <u>Deposits & Fossils:</u>

Coarse, gravel-rich river sand, deposited by the Rhine and Meuse (Early and Middle Pleistocene; Waalre and Beegden Formations), is overlain by a 1 to 2 metre thick layer of silty (cover) sand, deposited by the wind at the end of the last ice age (Late Pleistocene; Boxtel Formation). Along the flanks of the stream valley are elongated drifting sand dunes (Holocene; Boxtel Formation, Kootwijk Subunit), formed by human activity.

River & Streams:

The narrow Venloop lies in a broad and shallow periglacial valley, making it an example of an 'underfit river'.

3. Geosite Interests

Cultural History

On the Slabroekse Heide there are also more than 100 burial mounds from the Bronze Age and a cemetery from the Iron Age and Roman period. The heath and drifting sand area is also a fine example of a medieval cultural landscape shaped by human intervention (turf cutting, overgrazing).

Flora & Fauna:

The restored stream valley of the Venloop is very special, with rare animal and plant species having recently established themselves. The clean, mineral-rich water is now better retained in a broad, meandering stream, rather than being quickly drained away via a narrow, deep, straight ditch. Locally, there are sedge communities and cuckooflower meadows, though these are declining due to drying out. Cuckooflowers serve as a host plant for the orange tip butterfly. Grazing by European bison, Exmoor ponies and tauros cattle keeps the area open and varied. A rare species found in the stream valley is the great crested newt. The banded demoiselle dragonfly is also present, a species that only lives near clean, flowing water.

Industrial & Economic:

4. Relation with other geosites

The De Maashorst Nature Centre in Slabroek (GS_23) provides extensive information about this area. This area is one of five intended Wijst Restoration Projects designated by the province of Noord Brabant. Others include the wijst area Donzel west of Nistelrode (GS_02), the Wijstbos south of Zeeland (GS_197), the Sint Annabos in Uden (GS_07), and the source of the Esperloop in Gemert (GS_16).

5. Documentation & Resources


 Monitoring Maashorst Groote Wetering & Venloop Iris van der Laan Nico Ettema 2016 In opdracht van Stg. Natuurorganisaties De Maashorst

6. Maps, photos and figures

) TNO Intern 108/219



Location of geosite GS_013 – Slabroek. The Mellebreuk and Hoevense Fault are two northern branches of the Peel Boundary Fault.



Morphological characteristics of geosite GS_13 and the relationship with the faults

) TNO Intern 109/219



Stream valley of the Venloop (source: https://arkrewilding.nl/nieuws/2017/opening-slingerende-venloop)

) TNO Intern 110/219

Geosite: GS_14	Name
Significance: International	Gemert - Breuken Beleven
Coordinates (RD): 175790, 395242	
	Compart Police
Municipality	Gemert-Bakel
Province and country	Noord-Brabant
Protection status	Various
Site management organizations	Peelnetwerk, Mijn Mooi Brabant, ANWB (?)
Accessibility	Starting point and bike hire at VVV Gemert-Bakel
Relevance for tourism	The manifestations in and around Gemert are part of
	the "Breuken Beleven" cycle route (for example, the
	Joffer Grevenbroekpad)
Relevance for education & information	Several points along the route feature information
Relevance for education & information	
	boards
Teaching equipment	
- cacimis equipment	

1. Location

The village of Gemert is literally situated on the Peel Boundary Fault zone. As part of the "Experience the Faults" cycle route, several different manifestations of the Peel Boundary Fault zone around Gemert can be visited, including:

GS 14a: Hof van Nazareth: Lourdes made from ironstone blocks

GS 14b: Hopveld Bron:

GS_14c: Hazeldonklaan: Rood in de sloot (Red in the Ditch)

GS_14d: Joffer Grevenbroekpad – Infobord

GS 14e: Broekstraat: a reference to iron-rich ijzerbroek

GS 14f: Daalhorst

GS_14g: Sint Antoniusstraat de Mortel: Wayside cross on ironstone

GS_14h: Lochterweg - de Mortel elevation difference

2. Landscape formation and description

Origin of the landscape

The area around Gemert is characterised by the presence of several branches of the Peel Boundary Fault, namely the Gemert, Gemert-Zuid, Handel and Millheeze faults. The area not only displays morphological and hydrological landscape features, but also tells the story of human interaction, as evidenced by a large number of cultural-historical features.

Where iron-rich seepage water (wijst) emerges, iron oxides can be deposited in coarse-grained sediment near the surface. This cements the sediment, forming iron ore. Either the iron compounds cement the sand on the lower side of the fault, as iron-rich wijst water is transported over the fault due to the slope caused by the fault, or it cements coarse-grained fluvial deposits on the higher side of the fault. Both processes occur, which explains the widespread presence of iron ore. Pieces of iron ore were used in the past as building material and for iron production (since prehistoric times). They are often found in the foundations of churches in the region. Once bricks became affordable, this

) TNO Intern 111/219

tradition faded. Interestingly, iron ore blocks continued to be used in the construction of communal religious structures such as Marian grottos and hills where wayside crosses were placed.

Geomorphology:

GS_14f: Daalhorst: A striking feature of the Daalhorst is the elevation difference of 0.5 metres over a distance of 100 metres, caused by the Gemert-Zuid Fault, a side branch of the Peel Boundary Fault.

Hydrology:

In the immediate vicinity of the Peel Boundary Fault and its side branches, groundwater rises right up to the surface. Because the fault is impermeable, groundwater cannot pass through it. This creates upward pressure, causing water to well up from the subsurface. As a result, a narrow strip of wet and sometimes marshy ground forms along the fault. This phenomenon is known as wijst. The water, which sometimes comes from great depth, is very rich in iron. The iron in the groundwater oxidises when it comes into contact with oxygen. The iron oxide (rust) turns the water a reddish-brown colour. In various places in Gemert, this red water is clearly visible in the roadside ditches. This is also the case in Hazeldonklaan, where the wijst phenomenon at the Gemert-Zuid Fault causes this red water.

<u>Deposits & Fossils:</u>

River & Streams:

3. Geosite Interests

Cultural History

GS_14a Lourdes Grotto Gemert: This Marian grotto is modelled on the grotto in the French pilgrimage site of Lourdes and is made from rough blocks of ironstone or slag (waste from metal smelters). Inside the grotto is a small empty podium and several empty niches, possibly once intended for holy water stoups.

GS_14d: Joffer Grevenbroekpad: The information board describes the history of the Grevenbroekpad and the role of Joffer Grevenbroek in the region. It also provides information about the local flora and fauna.


GS_14g: On Sint Antoniusstraat in de Mortel stands a wayside cross on blocks of ironstone.

Flora & Fauna:

Industrial & Economic:

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures


) TNO Intern 112/219



) TNO Intern 113/219



Lourdes Grotto Gemert



Slope in the Daalhorst (source: www.landvandepeel.nl)

) TNO Intern 114/219

Geosite: GS_15	Name
Significance: Regional	Uden - Stadswijst
Coordinates (RD): 171297, 407764	ouen otausmjot
Municipality	Maashorst
Province and country	Noord-Brabant
Protection status	
Site management organizations	Municipality of Maashorst
Accessibility	All locations are situated in public parks
Relevance for tourism	All locations can be visited on a circular walk around
	Uden, possibly combined with sites in the Sint Annabos
	(GS_07).
Relevance for education & information	Various information boards: Mellepark ("De wijst op
	Melle")
Teaching equipment	

1. Location

The municipality of Uden is rich in faults belonging to the Peel Boundary Fault zone. Several city parks have been created along the faults in Uden, resulting in a green network running through the urban area, with ecological corridors connecting to the surrounding countryside. The guiding principle in their design was to restore the natural situation, allowing seepage water to flow down the slope into the grasslands below. The following city parks have been established in Uden:

GS_15a – Stadweide

GS_15b - Mellepark (named after the Mellebreuk)

GS_15c - Raampark

GS 15d - Zuiderpark

2. Landscape formation and description

Origin of the landscape

GS_15a: Commissioned by the (former) municipality of Uden, an "Urban Meadow" was created in autumn 2008, located near the Wilhelminalaan/Lippstadtsingel roundabout in Uden. Here, the Peel Boundary Fault was made visible by excavating over 15,000 m³ of soil and sand. This fault causes a significant difference in groundwater level within the site, which is clearly visible in the ponds and streams that have been dug there. After excavation, the meadow was entirely sown by hand with a special herb mixture. The intention is that in future, the area will be managed with grazing sheep. GS_15b: Mellepark is a park in the Melle district of Uden. The ribbon-shaped park was laid out in the 1970s and broadly follows the Melle Fault, a side fault of the larger Peel Boundary Fault. The park is characterised by its changes in elevation and numerous water features with rusty-brown water.

Geomorphology:

When the Pleistocene Meuse still flowed here, fine sand was sorted by the current in the riverbed, leaving gravel in pebble banks. In the shelter of these pebble banks, crescent-shaped sand dunes

) TNO Intern 115/219

(barchans) formed. After the last ice age, iron-rich seepage water began to flow through these pebble banks; the barchans trapped the iron-rich water, causing oxidation. This led to the formation of ironstone banks, which, like the compacted fault, push up the seepage water. In the Urban Meadow in Uden, several ironstone banks have been exposed by removing the topsoil.

Hydrology:

To manage high groundwater levels along the high side of faults, many retention basins have been constructed in Uden in the past, recognisable by their brown colour. This also reduced the seepage pressure on the Wijst soils. In addition, swales have been created on industrial estates to disconnect rainwater from the sewer and allow it to infiltrate, helping to prevent drying out and restore seepage pressure in the wijst areas. City parks have been established along the faults in Uden, creating a green network in the urban area with ecological corridors to the countryside. Uden demonstrates that urban wijst along faults within the built-up area is not so much a problem, but rather offers solutions for excessive urban heat; moreover, urban nature provides space for relaxation and helps relieve overcrowded nature reserves. Urban wijst is often characterised by orange ponds and ditches, as well as distinctive flora and fauna.

The water in Mellepark stands out for its rusty-brown colour and is full of iron particles that turn orange-brown when they come into contact with oxygen. In calm, dry weather, the particles settle as a layer on the bottom and the water is clear. But during a rain shower, the water is stirred up, the particles rise, and the water turns brown.

Deposits & Fossils:

Mellepark and the Urban Meadow hide an ancient secret: a former branch of the Meuse once flowed here, leaving behind gravel banks and sand dunes. Gravel and sand together form a permeable subsurface, allowing iron-rich groundwater to rise. In these places, a hard, dense layer of ironstone forms. Few plants grow on the layer itself, but the wet, nutrient-poor soil around it is attractive to rare species. Scattered throughout the parks, you'll find collections of ironstone blocks.

River & Streams:

3. Geosite Interests

Cultural History

Uden is a place where the wijst is less intense and mainly provided benefits, as the fields remained moist in summer. In Uden, large arable complexes lie between the faults, where settlement took place from the Iron Age through to the early Middle Ages. The presence of wijst certainly had a positive influence on settlement in these locations. In some places, later reclamations from the 12th and 13th centuries are located on or right next to a fault. Today, many place names in Uden recall the faults and wijst phenomena: for example, the bus stop at Mellepark is called Wijsthoek, and a residential area east of Mellepark on the high side of the Melle Fault is called "Breukrand".

Flora & Fauna:

Urban nature within Uden's built-up area often has higher biodiversity than the countryside. When the water features in the parks were created, seed banks were exposed, resulting in a wealth of special flora such as White rampion, Reed and Spotted orchid, Star of Bethlehem, Devil's-bit scabious, Marsh violet, Bistort and Great burnet. In the water you'll find Amphibious bistort, Floating club-rush, Bladderwort, Pillwort, Frogbit, Water violet and Whorled water-milfoil. The base-rich seepage allows rich woodland species of Bird cherry—Ash woodland and Oak—Hornbeam woodland to thrive, such as Hazel, Hornbeam, Maple, Small-leaved lime, Large-leaved lime, Bird cherry, Red and Yellow dogwood,

) TNO Intern 116/219

and Yew. Planted woodland bulbs and escaped garden plants are spreading spontaneously, such as Sweet and Pale wood violet, Bluebell, Snowdrop, Dutch crocus and Naples cyclamen. Thanks to the growth of Cuckooflower and Garlic mustard, the Orange tip butterfly is a common day butterfly in parks and gardens around the faults.

Other notable wijst areas with characteristic wijst vegetation in Uden's urban environment include:

- Park Hoeven Uden. Hoeven Fault with Bladderwort, Frogbit and Amphibious bistort in ditches.
- Zoggel Burenstraat Uden. Orang-brown waters.
- Kersenboomgaard Patrijsweg Uden. Ivy-leaved Crowfoot in cleared ditches.
- Landschapspark A50 Uden forms a long strip parallel to the Peel Boundary Fault. In the water features, special wijst flora grows such as Bladderwort, Amphibious bistort, Slender pondweed, Small pondweed and Great burnet occur.
- Duits lijntje at the Eikelheuvelweg Uden. Behind the ambulance station, a pool with seepage flora and ironstone bank.

Industrial & Economic:

4. Relation with other geosites

Sint Annabos (GS_07)


5. Documentation & Resources

6. Maps, photos and figures



Location of GS_15 – Uden Urban Wijst and the various city parks where wijst phenomena can be observed.

) TNO Intern 117/219



) TNO Intern 118/219

Geosite: GS 16	Name
Significance: National	Geneneind-Esp
Coordinates (RD): 178952, 391766	Continue Lop
Municipality	Gemert-Bakel
Widincipanty	Gement-baker
Province and country	Noord Brabant
, , , , , , , , , , , , , , , , , , , ,	
Protection status	
Site management organizations	Stichting Gemert-Bakel Toerisme en Recreatie,
	Heemkundekring Gemert-Bakel, VVV Gemert-Bakel.
	Wijstherstel: Waterschap AA & Maas, Gem. Gemert-
	Bakel, Prov N Brabant, particulier
Accessibility	All points are accessible by bicycle. Cars can park in
	Bakel around Sint Wilbertsplein.
Relevance for tourism	Geneneind and Esp are part of the "Breuken Beleven"
	route through the Gemert-Bakel area. The starting
	point is at Kasteel Gemert.
Relevance for education & information	IVN Gemert-Bakel Nature Education Centre (GS_16b)
	and information board.
Teaching equipment	
Scientific interest	As part of the Integral Area Programme Bakel-
	Milheeze North (IGP), extensive geohydrological
	research has been conducted in the Geneneind area
	in recent years to increase the opportunities for wijst
	restoration in this area. Among other things, three
	rows of piezometers have been placed at different
	depths, perpendicular to the faults in the subsoil. In
	this way, the fluctuation in groundwater levels is
	made clear.
Description	

1. Location

Geneneind is a hamlet in the rural area north of Bakel.

2. Landscape formation and description

Origin of the landscape

The Peel Boundary Fault runs through the Geneneind area. The horst is characterized by wijst phenomena. Groundwater rises strongly here, but due to human modifications of the soil, both the wijst grounds and the height differences are less visible. Various parties are therefore working to restore the wijst grounds. Because the groundwater emerging before the faults can be centuries old, the water is very clean, low in calcium, and characterized by low(er) concentrations of nutrients such as nitrates and phosphates. Where the groundwater comes into contact with oxygen (in the outside air), oxidation occurs, followed by complex chemical processes in which, among other things, iron, nickel, phosphate, and nitrate can be bound and precipitated. Oily films on these watercourses indicate the presence of iron bacteria.

Geomorphology:

) TNO Intern 119/219

On the Geneneindse Heide near Bakel, the Peel Boundary Fault can be recognized with some difficulty by a terrain step and by the vegetation.

Hydrology:

Groundwater jump of almost 2 m over the Peel Boundary Fault. Hydrologically considered, the incoming wijst water at the Peel Boundary Fault is a collection of various separate water flows, each with its own travel time (age) and composition. Geohydrological measures that intervene in the water management situation of wijst areas will generally take "a longer period" to have an effect. This is a direct result of the fact that the wijst water is on the way for at least several years, but often decades and for some part even centuries before it emerges near underground faults. This also applies to the Geneneind area. The Esperloop wijst restoration project aims to restore the hydrological situation and sets the following goals: upgrading the Esperloop to an Ecological Connection Zone (EVZ); stream restoration; creating a robust and resilient water system (sufficient water and future-proof); improving the ecological potential; enabling fish migration; stream restoration of the Esperloop and creating added value for wijst restoration.

Deposits & Fossils:

River & Streams:

At Geneneind in Bakel, the Esperloop (GS_16c) runs through the restored wijst area on the Peel Boundary Fault. The Esperloop flows further to the Grotelsche Heide. The Esperloop is a meandering stream fed by wijst water. The water of the Esperloop is very clean but contains a lot of iron. The Esperloop splits at Milheeze in the Stippelberg area from the Snelle Loop and is largely unchanneled. As such, it is one of the rare examples of a more or less intact 'heath brook'. This narrow stream meanders through the Grotelsche Forest to rejoin the Snelle Loop near the hamlet of Grotel, south of De Mortel.

3. Geosite Interests

Cultural History

Geneneind used to belong to the very old hamlet of 'Esp'. The cadastral map shows that various farms and small buildings formed a fairly large settlement. There is still one real old long-gable farm standing in the street, which is the farm at No. 18, built in 1846. The map shows a stream, which is the Esperloop, which still lies in the same place today. In 2015, the course of this stream was restored. At Geneneind, there are even still steep edges and a so-called hollow road (in combination with "bulging fields"), the only one still preserved in Noord-Brabant. The Espse Hoeve (Esp 3; GS_16a) is a moated farm from the 15th century. The moats, which have been partially restored, are filled with rising wijst water. The rest of the site has the protected status of 'archaeological monument'. The rising groundwater in the streams along the arable land was important for the farmers. The relatively warm water ensured that the frost stayed in the ground for a shorter time, allowing the farmer to use the land for up to two months longer.

Flora & Fauna:

Both locally on the banks of the Esperloop and at the bottom of the steep banks, there are stream valley plants such as marsh marigolds and black sedge, which do not occur further in the Grotelsche Forest (an old heath field) and are relatively rare in the area. A rare situation, which can possibly be explained by phosphate fixation caused by the high iron content. The rare common dragonfly also benefits from the pure water. The stream also flows along some floristically interesting grasslands.

) TNO Intern 120/219

Industrial & Economic:

4. Relation with other geosites

Geneneind is a beautiful wijst area along the Peel Boundary Fault, comparable to Sint Annabos (GS_07) but located further south.


5. Documentation & Resources

 Lapperre, R. E. (2025). Fault-Zone Hydrogeology in Unconsolidated Sediments: Cross-Fault Groundwater Levels, Fault Sealing, and Permeability Distribution in the Roer Valley Rift System. [PhD-Thesis, Vrije Universiteit Amsterdam]. https://doi.org/10.5463/thesis.1223

6. Maps, photos and figures



Location of GS_16 indicated with a light blue line



Left: crop marks and soil moisture contrasts on a satellite image indicating the exact location of the fault at Geneneind, between the wet foot wall (FW) and the drier hanging wall (HW) agricultural plots. Right: overview of the regional isohypses pattern (groundwater-level contours)

) TNO Intern 121/219

on 1 April 2016 (m NAP), with the fault modelled as an (almost) impermeable boundary, and corresponding groundwater flow path (blue arrows) on both sides of the fault using the regional groundwater model GRAM2.0 from Water Authority Aa en Maas. (from Lapperre, 2025)



"Bulging fields" near Geneneind

) TNO Intern 122/219

Geosite: GS_17	Name
Significance: National	Neer
Coordinates (RD): 195969, 364656	
Municipality	Leudal
Province and country	Limburg
Protection status	
Site management organizations	
Accessibility	rom Neer, head west via Keizersloop towards
	Keizersbos
Relevance for tourism	Walking area
Relevance for education & information	Informationpanel at De Kwir
Teaching equipment	
Scientific interest	Along the Reverskamp was one of the five research locations (KNMI, 2000). The trench is no longer visible. Results were published in Netherlands Journal of Geosciences / Geologie en Mijnbouw 81 (1): 39–60 (2002).

1. Location

The Peel Boundary Fault crosses the River Meuse near the village of Neer. Along the Peel Boundary Fault near Neer, several different manifestations of the fault zone and its relationship with streams can be observed, namely:

GS_17a: Terreintrede Peel Boundary Fault

GS_17b: Seepage site Keizersbos

GS_17c: Research trench at Raverskamp

GS_17d: De Kwir GS_17e: Het Sprunske

2. Landscape formation and description

Origin of the landscape

In the southern part of the Peel Horst, where the Meuse Valley cuts through the horst, the relief and hydrology are far more complex than elsewhere along the Peel Boundary Fault. Here, not only the Peel Horst shapes the relief, but also the river terraces in the Meuse Valley. These terraces sometimes have metre-high scarps and old river channels, which now often host streams. Streams may also originate at the foot of the Peel Boundary Fault, where they are fed by groundwater seepage and rainwater.

Geomorphology:

West of Neer, between Vlaas and Schooldijk (GS_17a), the Peel Boundary Fault is slightly visible in

) TNO Intern 123/219

the topography but clearly recognizable in the AHN (elevation model). The fault can also be located using cropmarks (caused by differences in vegetation due to wet versus dry soils). This height difference mainly resulted from several strong earthquakes around 12,000 BC, as shown by research at the trench near Neer-Raverskamp. The fault offsets a Meuse terrace from the Late Pleniglacial (abandoned by the Meuse around 18,000 BC), as indicated by the height difference between the Peel Block and the Roer Valley Graben. This difference is absent in younger Meuse terrace deposits. Based on OSL dating, Van den Berg et al. (2002) concluded that two relatively large faulting events occurred within a short time span (less than 3,000 years) between the Last Glacial Maximum (c. 15,800 years ago) and the onset of the Younger Dryas (c. 13,000 years ago), while only one very minor displacement occurred during the Younger Dryas and the Holocene.

Hydrology:

In the Meuse Valley, places with iron-rich seepage are locally called "Sprunk". It is not always clear whether such seepage is caused by the Peel Boundary Fault or by elevation differences associated with river terraces. The seepage in the area of De Kwir (GS_17d) occurs exactly where the Keizer stream intersects the Peel Boundary Fault. Behind St Martinus Church in Neer (GS_17e), seepage water also used to emerge, but this phenomenon is no longer visible due to artificial drainage. On the Ferraris Map (1771–1778), it is still clearly shown that the seepage fed a north-flowing watercourse (now known as the Wijnbeek).

Deposits & Fossils:

River & Streams:

The Keizersloop (or simply Keizer) originates at the foot of the Peel Horst and is partly fed by groundwater from the fault. The Keizer roughly follows the base of the Peel Horst and has cut deeply through cover sand ridges and the relief of the Meuse river terraces. The deepest incision is visible at De Kwir (GS_17d). The Keizer is fed by rainwater from the Roer Valley Graben, which meets the Peel Horst here, and by groundwater from the Peel Boundary Fault. The Keizerbos monastery was originally surrounded by moats, remnants of which are still visible. These were probably dug later, but the location of Keizerbos undoubtedly relates to the continuous water supply. The Keizer touches the north-eastern corner of the monastery grounds, where seepage water still occasionally surfaces. The Keizer joins the Neerbeek in Neer, which ultimately flows into the Meuse. The Neerbeek itself is formed by the confluence of the Leubeek (GS_09) and the Haelense Beek north of Nunhem.

3. Geosite Interests

Cultural History

Although the origins of the village of Neer date back to the early Middle Ages, the area upstream along the Keizer remained forested until after the year 1000 and was only reclaimed in the High Middle Ages (1000–1250). At the Keizer(sloop), the outlying farm Cherserburch is mentioned in 1185. On the same site, Norbertine monks later (1230–1246) founded the Keizerbos monastery, of which only a few buildings remain.

The Friedesse watermill on the Neerbeek is an undershot mill, now functioning as a corn mill. The original mill was built in 1343, but the current mill house and miller's house date from 1717. The mill belonged to the Abbey of Thorn. After years of disuse and decay, it became operational again following restoration in 2000–2002. The mill was designated a national monument in 1973.

) TNO Intern 124/219

Flora & Fauna:

Industrial & Economic:

4. Relation with other geosites

5. Documentation & Resources


Berg, M. van den, K. Vanneste, B. Dost, A. Lokhorst, M. van Eijk and K. Verbeeck (2002), 'Paleoseismic investigations along the Peel Boundary Fault: geological setting, site selection and trenching results', in: Netherlands Journal of Geosciences; Geologie en Mijnbouw 81

6. Maps, photos and figures



Location of GS_17 and the various manifestations along the Peel Boundary Fault.

) TNO Intern 125/219



) TNO Intern 126/219

Geosite: GS_18	Name
Significance: National	Helenaveen
Coordinates (RD): 192070, 377851	
Municipality	Deurne
Province and country	Noord-Brabant
Protection status	Protected village view (Beschermd dorpsgezicht)
Site management organizations	
Accessibility	Soemeersingel 127, 5759 RD Helenaveen
Relevance for tourism	Part of the Geopad Verheven Peel cycle route
Relevance for education & information	Information board between Helenaveen and Sevenum about the history of coal drilling in the Peel
Teaching equipment	

1. Location

Helenaveen is a peat colony situated at the easternmost point of the Deurne Peel and is one of the most easterly villages in the province of Noord-Brabant. Helenaveen was the first peat colony, founded by peat contractor Jan van de Griendt (and named after his wife Helena). It is a protected village view, partly due to its numerous canals and side channels (wijken) and the Koningshoeven, farms built after the peat extraction period.

2. Landscape formation and description

Origin of the landscape

During the latter part of the last Ice Age, the Peel landscape was sparsely vegetated for a long time. Due to tectonic activity and the drifting of cover sands, large, almost closed basins eventually formed in the Griendtsveen Graben and in the blocked upper reaches of Brabant streams. In these basins, fen peat began to develop during the final stage of the Ice Age. During the Holocene, these fens evolved into transitional mires and eventually into raised bogs, which expanded beyond the original basins and largely covered the surrounding sand ridges. For a long time, these wet peatlands were difficult to access, but they were eventually reclaimed for peat exploitation. Helenaveen is an example of large-scale industrial peat cutting.

Geomorphology:

Hydrology:

Parallel to the Helenavaart canal, a dyke was built 200 metres away, containing an impermeable barrier to create a high and stable water level behind it, ideal for raised bog restoration. Along the Helenavaart, emphasis was placed on the peat-colonial cultural landscape with its Koningshoeven and three-pronged canal system. Remarkably, this spatial planning process involved exemplary

) TNO Intern 127/219

cooperation between authorities, recreation entrepreneurs, and nature and agricultural organisations.

Deposits & Fossils:

The Van de Griendt brothers commissioned engineers from the Water Authority to conduct various studies on the feasibility of canal construction. Based on boreholes, profiles were constructed showing four soil layers: Subsoil, 'Black peat' or *klot*, 'Pale peat' or *vale moer*, topsoil. The black peat was sometimes classified as 'heavy', 'light klot', or 'less good'. Due to the undulating subsoil, peat thickness varied greatly. For the first 30 years, peat cutting focused on black peat, as pale peat (grey peat) was unsuitable as fuel. Peat litter, made from grey peat, had not yet been invented.

River & Streams:

The Helenavaart was the lifeline of the company 'Maatschappij Helenaveen'. Typical of this peat colony is that from the main canal, side canals (wijken) were dug at 600-metre intervals, with two cross-canals branching from the main canal, creating a three-pronged system. This allowed more space for buildings and reduced the number of bridges needed. This system is still visible in several places along the canal. This distinctive peat-colonial structure earned Helenaveen its designation as a protected village view in 1998.

3. Geosite Interests

Cultural History

Peat cutting required thousands of labourers, recruited not only locally but also from afar (even from Prussia). Many came from the north-east of the Netherlands, where peat extraction was declining. Helenaveen became a thriving village with a mixed population, evidenced by the presence of both a Catholic and a Protestant church. Offices and various industrial buildings were also established, including a beautifully restored former peat litter factory.

Flora & Fauna:

Industrial & Economic:

To exploit his purchased section of the Peel, Jan van de Griendt, a manufacturer from 's-Hertogenbosch, founded the peat company Maatschappij Helenaveen, named after his second wife, Helena Panis. The company quickly grew into a huge enterprise, a highly profitable multinational selling peat (or peat litter) across much of Europe.

In 1903, the State Mineral Exploration Service (ROD) was established to search for coal outside South Limburg. After studying the subsurface of neighbouring North Rhine-Westphalia and the relief of Brabant and Limburg, ROD director Van Waterschoot van der Gracht concluded that the Peel was a tectonically elevated, fault-bounded area. The first coal drilling took place in Helenaveen in 1906 and struck coal at a workable depth of 914 metres. This success was followed by more successful drillings in the Peel. Due to economic uncertainty after the First World War, exploitation did not proceed. During the 1930s crisis, there were unsuccessful calls to develop Peel coal mines as a means of combating unemployment.

After the Second World War, rising energy demand led to the establishment of the Peel Commission. Seismic surveys and new drillings confirmed earlier ROD estimates. In 1963, the Peel Commission concluded that there were sufficient coal reserves to exploit two mines (Peel Mines I and II). Construction of the State Mine Beatrix began in 1954 but was halted in 1962, mainly due to

) TNO Intern 128/219

the discovery of the Groningen gas field in 1959. The monument on Helenaveenseweg – opposite Toverland amusement park – commemorates the fact that between 1913 and 1915, the State Mines drilled for coal at the site of the current monument.


4. Relation with other geosites

Griendtsveen (GS_04) is the other peat colony on the northern side of the Deurne Peel.

5. Documentation & Resources


- Theo van de Mortel Boortorens in de Peel (https://natuurtijdschriften.nl/pub/1021589)
- André Vervuurt: Helenaveen van Boven tot Koningslaan 1900-1925





Location of GS_18 Helenaveen indicated with a light blue line.

) TNO Intern 129/219



) TNO Intern 130/219

Goosito: GS 10/2 h)	Name
Geosite: GS_19 (a,b)	113
Significance: Regional	Wijstbos Zeeland
Coordinates (RD): 176158, 411902	
Municipality	Maashorst
Province and country	Noord-Brabant, the Netherlands
Protection status	Aardkundig monument
Site management organizations	Land van Cuijk
Accessibility	Accessible from the junction of Langenboomseweg and Veldzuring.
Relevance for tourism	Part of the cycle route "Langs de breuk (Along the Fault) through the municipalities of Maashorst and Bernheze.
Relevance for education & information	Walking trail and information panels
Teaching equipment	

1. Location

The Wijstbos in Zeeland is one of five Wijst Restoration Projects designated by the province of Noord-Brabant. The area lies to the south-east of Zeeland and has been redeveloped in phases. It consists of a woodland section with balsam poplars and black alder (GS_19a) and several grasslands near the hamlet of Graspeel (GS_19b), featuring old pollarded willows and alder/ash scrub.

2. Landscape formation and description

Origin of the landscape

This geosite is located on the eastern side of the Peel Horst, on a slightly lower fault block. It is therefore the only one of the five Wijst Restoration Projects that drains eastwards towards the Meuse Valley. Older Rhine and Meuse deposits are overlain by a layer of cover sand and intersected by several south-east to north-west trending fault lines, which have displaced the subsurface deposits stepwise.

Geomorphology:

The Wijstbos lies on a slightly lower fault block of the Peel Horst. However, the landscape has been heavily re-parcelled, leaving little trace of the original relief with its cover sand ridges and shallow stream valleys. Just east of the Wijstbos, a terrace step of approximately 0.5 m is visible.

Hydrology:

Deep ploughing has disrupted the area's original water regime, causing desiccation.

Deposits & Fossils:

Coarse, gravelly river sand deposited by the Meuse (Middle Pleistocene; Beegden Formation) is overlain by a 1–2 m thick layer of silty cover sand deposited by wind at the end of the last Ice Age (Late Pleistocene; Boxtel Formation). Locally, this sand was reworked by wind during the Holocene induced by human activity.

) TNO Intern 131/219

River & Streams:

This geosite lies on the eastern side of the Peel Horst, on a lower fault block. It is therefore the only Wijst Restoration Project that drains eastwards towards the Meuse Valley. Drainage occurs via straightened ditches.

3. Geosite Interests

Cultural History

The area contains ancient field complexes that have been in use since the Iron Age. Their location is partly determined by the orientation of the fault lines and the resulting hydrological conditions.

Flora & Fauna:

As part of the Graspeel land consolidation, the edges of poplar woods were cleared and the banks of watercourses were made shallower. Several ponds were also created. The ditches host one of the largest populations of bistort in Brabant. The area is home to the little owl and many other breeding birds such as blackcap, icterine warbler and nightingale. In winter, siskins, redpolls, goldfinches, fieldfares and redwings forage here.


Industrial & Economic:

4. Relation with other geosites


This area is one of five designated Wijst Restoration Projects in the province of North Brabant. Others include the wijst area of Donzel west of Nistelrode (GS_25), the Venloop valley near Slabroek (GS_13), the Sint Annabos in Uden (GS_07), and the source of the Esperloop in Gemert (GS_16).

5. Documentation & Resources

6. Maps, photos and figures



) TNO Intern 132/219



AHN (shaded relief) image of the GS_19 site



Information panel at the entrance to the Wijstbos near Zeeland

) TNO Intern 133/219

Geosite: GS_20	Name
Significance: International	De Kuilen, Langenboom
Coordinates (RD): 179989, 412109	
Municipality	Land van Cuijk
Province and country	Noord-Brabant (NL)
Protection status	Natuurmonument
Site management organizations	Natuurmomumenten, private owners
Accessibility	The area is easily accessible from Fazantenweg. Parking is available.
Relevance for tourism	De Kuilen ceased sand extraction in 2012 and now features one of the cleanest sandy beaches in the Netherlands. There is a beach pavilion, a playground, and numerous cycling and walking paths in the adjacent nature reserve. Natuurmonumenten has established a circular walking route around the lake and the adjoining Langenboomse Woods.
Relevance for education & information	Fossil finds can be viewed at the De Kuilen visitor centre in Mill, which also provides information about the surrounding nature reserve.
Teaching equipment	

1. Location

De Kuilen lies east of the village of Langenboom, on the crest of the Peel Horst.

2. Landscape formation and description

Origin of the landscape

De Kuilen at Langenboom consists of lakes formed by the extraction of "green sand". Langenboom is situated on the crest of the Peel Horst. Here, green sands from Miocene and Pliocene formations occur less than one metre below the surface, whereas elsewhere in the Netherlands they lie at much greater depths.

Geomorphology:

Hydrology:

Green sand contains glauconite, which releases iron into groundwater. Oxidation of this iron gives the seepage water a brown colour. Although De Kuilen lies between two faults, iron-rich seepage from point sources is present. The water is exceptionally clear.

Deposits & Fossils:

De Kuilen is renowned for fossil shells from the Pliocene, which can still occasionally be found along the lake shores, but it is especially famous for the wide variety of fossils excavated here in the past. Langenboom is recognised in Europe as one of the most productive sites for Miocene and Pliocene fossils – it even ranks first for fossil birds. Palaeontological finds include the ungulate Chalicotherium,

) TNO Intern 134/219

four new auk species, and the short-snouted dolphin (Protophocaena minima). At Langenboom, the dredged sand, including fossils, was deposited on land, allowing amateur geologists and palaeontologists to collect specimens on a large scale.

River & Streams:

3. Geosite Interests

Cultural History

Flora & Fauna:

The flora and fauna are highly diverse due to the alternation of wet and dry habitats. The area supports plant species typical of wijst areas, such as needle spike-rush (Eleocharis acicularis). Other less common species that thrive on base-rich seepage include brown sedge, pennyroyal, foxtail, and the rare water moss Fontinalis antipyretica..

Industrial & Economic:

De Kuilen originated from the extraction of green sand from Miocene and Pliocene formations. For road construction, glauconite-bearing sand is an excellent material; glauconite grains are angular and interlock firmly, a property that makes the green sands of Langenboom highly suitable for building motorway embankments.


4. Relation with other geosites

De Kuilen has no direct relationship with nearby geosites. In Liessel, forty kilometres south of Langenboom, a sand pit (GS_08) has yielded Miocene and Pliocene fossils. This site also lies on the Peel Horst.


5. Documentation & Resources

6. Maps, photos and figures

) TNO Intern 135/219



) TNO Intern 136/219



South-east to north-west cross-section through the GeoTOP subsurface model (v1.6.1) at the De Kuilen sand extraction site, located on an elevated fault block of the Peel Horst where Miocene deposits lie closer to the surface (see inset map for location)



Finds from the local history society "Felix Walter" – Langenboom.

) TNO Intern 137/219

Geosite: GS_21 (a, b, c) Significance: National	Name Sint Hubert
Coordinates (RD): 184471, 409842	
Municipality	Land van Cuijk
Province and country	Noord-Brabant
Protection status	Aardkundig Waarveol Gebied (Peel Boundary Fault System)
Site management organizations	GS_21a: Land van Cuijk GS_21b: Heemschut Sint Huybert GS_21c: Het Duitse lijntje
Accessibility	All sites (GS_21a,b,c) are easily accessible and located along public roads.
Relevance for tourism	This geosite forms part of several walking and cycling routes around Sint Hubert and Mill.
Relevance for education & information	Informationpanel at Broekstenen Huuske Informationpanel about Het Duitse Lijntje at the Scheiwalweg crossing with the old railway line
Teaching equipment	

1. Location

This geosite comprises three locations in and around the village of Sint Hubert: the church square (GS_21a), the IJzerbroekseweg (GS_21b), and the German railway line (Het Duitse Lijntje, GS_21c), situated about 1 km from the village centre. These sites illustrate the history of iron ore extraction.

2. Landscape formation and description

Origin of the landscape

Sint Hubert lies west of the Tegelen Fault on a high fault block, as evidenced by wijst phenomena and iron ore deposits in the relatively coarse-grained old Meuse deposits (Beegden Formation). However, it is located east of the morphological scarp of Mill.

Geomorphology:

The geosite lies east of the Mill scarp (see also GS_02), which was formed by river erosion during the last two Ice Ages in the Middle and Late Pleistocene. The morphology of this geosite shows an atypical relationship between tectonic activity and river processes, as the greatest elevation difference (the scarp) does not coincide with the fault.

Hydrology:

Occurrence of iron ore (bog iron) formed by precipitation of iron oxides from iron-rich seepage (wijst) on the elevated block along the Tegelen Fault.

Deposits & Fossils:

River & Streams:

) TNO Intern 138/219

3. Geosite Interests

Cultural History

On the square in front of Sintt Hubert's Church (GS_21a) stands the 1994 sculpture 'De Broeksteenwerker' by Marian van Puyvelde, commemorating iron ore extraction in Het IJzerbroek. In this area, between Sint Hubert and Haps, bog iron was mined between about 1825 and 1860 using pickaxes. Blocks of iron ore found locally are displayed near the sculpture. These so-called "broekstenen" are conglomerates of iron oxide, sand, clay, and gravel.

In the church of Sint Hubert, blocks of iron ore are clearly visible in the masonry. On IJzerbroekweg once stood the "Broekstenen Huuske" (1825), built from iron ore blocks. The house disappeared around 1950; the information board tells the story of the subsurface and the house (GS_21b). The iron ore blocks were quarried locally to reclaim farmland. The hard iron ore layer prevented rainwater from draining downwards and groundwater from rising. Crops suffered from waterlogging in wet weather and drought in dry periods. Special pickaxes were used to break the ore, and large pieces were removed for practical use. The red clay was brought to the surface and, mixed with peat sods, formed the new topsoil. For water management, a ditch was dug and fitted with sluices to regulate water levels.

Het Duitse Lijntje (The German railway) was opened in 1873 by the North Brabant–German Railway Company (NBDS). It connected the seaports of Vlissingen, Antwerp, and Rotterdam with the German hinterland and was part of the fastest route between London, Berlin, and St Petersburg. During the First World War, the line was heavily used for domestic workers and refugees entering the Netherlands. In the Second World War, it played a major role: on 10 May 1940, German troops invaded the Netherlands via this line at 5 a.m. Near Zeeland, soldiers disembarked after breaking through the Peel-Raam defensive line. At the end of the war, the Allies used the railway to invade Germany. Passenger services ceased in 1950, and freight services ended in 2004. Most of the track was removed in 1973, but traces remain in the landscape.

The section between Scheiwalweg and Achterdijk near Sint Hubert is elevated because the area was once frequently flooded. The embankment is still visible. Both Scheiwalweg (GS_21c) and Achterdijk had signalman's cottages. Two have been preserved in nearby Mill. Mill station has been converted into a youth centre (De Wissel). A few metres of track, a loading quay, and an artwork symbolizing the railway remain. At the former station site in Haps, an original signal post still stands.

Flora & Fauna:

The old railway line now serves as a wildlife corridor. Species include badgers, foxes, buzzards, kestrels, roe deer, and various amphibians. The tree stands and ponds along the route make the area attractive for nature enthusiasts.

<u>Industrial & Economic:</u>


4. Relation with other geosites

Alongside the sites in Mellepark and Stadsweide in Uden (GS_15), this geosite illustrates the historical importance of iron as a valuable raw material, dating back to the Iron Age. The site lies at the base of the Mill steilrand (escarpment, GS_02).

5. Documentation & Resources

6. Maps, photos and figures

) TNO Intern 139/219



) TNO Intern 140/219



Sculpture De Broeksteenwerker (GS_21a) and the Broekstenen Huuske (GS_21b)



Informationpanel "Het Duitse Lijntje" at Scheiwalweg in Sint Hubert (GS_21c)

) TNO Intern 141/219

Geosite: GS_22 Significance: National Coordinates (RD): 184072, 386789	Name Kasteel Deurne-Heiakker
Municipality	Deurne
Province and country	Noord-Brabant
Protection status	National Monument (Rijksmonument)
Site management organizations	Municipality of Deurne
Accessibility	The ruins of the Great Castle are located in a public park and are freely accessible to walkers and visitors.
Relevance for tourism	The site is part of several walking routes, such as the Deurne Castle Route. Guided tours can be booked via VVV Deurne.
Relevance for education & information	De Ossenbeemd, adjacent to the castle garden, is the leading nature and environmental centre in the Geopark area, offering by far the most information about the Peel Fault.
Teaching equipment	

1. Location

The Great Castle Groot Kasteel) is located at Haageind 37, Deurne. There is also a Small Castle and a watermill. The Peel Boundary Fault runs through Vlierpark in Deurne, just north of the castles and watermill.

2. Landscape formation and description

Origin of the landscape

The moat around the Great Castle is fed by groundwater supplied via the Vlier stream. The most suitable location for a watermill on a stream is on or just downstream of the point where the stream crosses a fault. The Haageind site, just south of the Peel Boundary Fault, is an excellent example of a reclamation from the 12th and 13th centuries. The choice of location for the watermill and castles was influenced by the fault of the Peel Boundary Fault Zone Faults: strong seepage along the faults ensured a relatively constant water supply, while the height difference between horst and graben provided additional fall for water power.

Geomorphology:

The height difference at the castle itself is not dramatic, but north and south of the site the terrace step along the Peel Boundary Fault becomes more evident. On Kulertseweg, at the southern edge of Heiakker (about 1 km south of the castle), a clear terrace step can be seen in the road.

Hydrology:

Groundwater rises through seepage just north of the impermeable Peel Fault. The groundwater cannot flow further into a lower area and therefore emerges at the surface, feeding the Vlier stream.

Deposits & Fossils:

) TNO Intern 142/219

Kulertseweg is one of the few places along the Peel Fault where exposed Pleistocene Meuse sediments can be seen.

River & Streams:

The Vlier is a drainage stream of the Peel, originating near the Deurne Canal and flowing into the Kaweise Loop near the hamlet of Molenhof. The stream flows northwest from the Peel Horst and reaches the Deurne basin near the ring road. The section east of the current ring road was probably entirely dug before the 19th century to serve as a new upper course. The Vlier powers the watermill at Haageind, which has been there since the 14th century, and also supplies water to the moats of the Great Castle (Groot Kasteel) and the Small Castle (Kleine Kasteel), both built in the 14th century on either side of Haageind. The meandering of the stream has been preserved near the castle garden, though largely lost elsewhere. Further downstream, the Vlier also powered the watermill on Kerkeindseweg until it was destroyed by fire in the 17th century. Today, the Vlier is a canalized stream connected to the Deurne Canal just south of Griendtsveen.

3. Geosite Interests

Cultural History

The Great Castle (Groot Kasteel) in Deurne, located in the Vlier valley, was built in the 14th century. At that time, it was rarely inhabited; only from the 16th century did it become the residence of the Lordship of Deurne. Before the Great Castle was built, the watermill and the Small Castle at Haageind already existed. The nearby Peel Fault guaranteed a steady water supply for the watermill and moats, fed by the Vlier stream. This is the only surviving watermill along the Peel Boundary Fault. Unfortunately, the moat could not prevent the Great Castle from being shelled at the end of the Second World War during the liberation of Deurne (24 September 1944), as German SS soldiers were hiding inside. The resulting fire destroyed most of the castle, leaving only part of the outer walls standing.


The Peel Boundary Fault runs through Vlierpark in Deurne. In autumn 1993, Italian artist Roberto Ruggiu created an artwork on the Fault, which must remain undeveloped due to its geological conditions. The artwork consists of two interconnected elements stretching over about one kilometre: five wells and a double row of trees with differently coloured blossoms.

Flora & Fauna:

Industrial & Economic:

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures

) TNO Intern 143/219



) TNO Intern 144/219



Double row of apple trees marking the Peel Fault in Vlierpark (source: www.landvandepeel.nl)

) TNO Intern 145/219

Geosite: GS_23 Significance: Regional Coordinates (RD): 169828, 412393	Name Natuurcentrum De Maashorst
Municipality	Bernheze
Province and country	Noord-Brabant, The Netherlands
Protection status	National Monument (Rijksmonument)
Site management organizations	Staatsbosbeheer
Accessibility	Easily accessible, including for people with reduced mobility.
Relevance for tourism	Starting and information point for nature-oriented activities in De Maashorst. Excursions are organized, and catering facilities are available.
Relevance for education & information	Information hub for nature-oriented activities in De Maashorst. An education team provides lessons on nature, sustainability, and water management.
Teaching equipment	Equipment is available for hire for expeditions in the surrounding area, particularly aimed at children and families.

1. Location

Natuurcentrum De Maashorst is located in the hamlet of Slabroek, on the boundary between the municipalities of Bernheze and Maashorst.

2. Landscape formation and description

Origin of the landscape

The Nature Centre is situated in a historic natural and cultural landscape. The area was shaped by river activity (Meuse and Rhine), periglacial processes (periglacial valley formation, cover sand deposition), and significant fault activity. From the Bronze Age onwards, humans utilised this natural variation, creating cultural elements such as heathlands.

Geomorphology:

Hydrology:

Deposits & Fossils:

River & Streams:

3. Geosite Interests

Cultural History

The Nature Centre is housed in a historic farmhouse in the hamlet of Slabroek, which forms part of a small-scale historic cultural landscape.

Flora & Fauna:

) TNO Intern 146/219

The Nature Centre provides extensive information about the flora and fauna of the surrounding area.

Industrial & Economic:

4. Relation with other geosites

The Nature Centre offers comprehensive information about De Maashorst, particularly the area around Slabroek (GS_13).

5. Documentation & Resources

6. Maps, photos and figures



Location of GS_23 Natuurcentrum De Maashorst indicated with a light blue line



Natuurcentrum De Maashorst (www.natuurcentrumdemaashorst.nl)

) TNO Intern 147/219

Geosite: GS_24 Significance: International	Name Plan de Burgt (under development)
Coordinates (RD): 175262, 402267	
Municipality	Boekel
Province and country	Noord-Brabant
Protection status	
Site management organizations	Municipality of Boekel
Accessibility	
Relevance for tourism	The geosite is part of the walking route Ommetje Peel Boundary Fault Boekel (set out by the Boekel Local History Society).
Relevance for education & information	The residential area is being developed as a Peel Boundary Fault Park, but without signage. It is intended to subtly draw visitors' attention to the natural phenomena.
Teaching equipment	
Scientific interest	On 22 May 2025, mechanical test drillings (up to 8 m deep) were carried out on the De Burgt Phase 2 site in Boekel to determine the exact position of the Peel Boundary Fault(s).

1. Location

On the northern side of Boekel, the De Burgt residential area is being developed. A fault of the Peel Boundary Fault Zone runs directly through this neighbourhood. The term Wijstwonen ("Wijst Living") expresses this unique living environment. The name refers to the "Hoge Burgh", a group of plots situated around the Peel Boundary Fault.

2. Landscape formation and description

Origin of the landscape:

The De Burgt area was formed by the activity of one of the three fault segments of the Peel Bounbdary Fault Zone. The variation in wijst phenomena, ironstone layers, terrace steps, groundwater springs, and vegetation differences clearly demonstrates that the fault runs through the developing residential area.

Geomorphology:

The planning area exhibits a very large height difference due to the presence of the Peel Boundary Fault Zone. The eastern part of the area lies significantly higher than the western part. The elevated Peel Horst is also a relatively wet area.

Hydrology:

The Peel Boundary Fault marks the transition between the Peel Horst and the lower Roer Valley Graben. Groundwater flows from high to low, but at the fault plane, this flow is impeded by impermeable layers, causing the water to rise and emerge as seepage at the surface. This makes the

) TNO Intern 148/219

higher ground wet while the lower ground remains dry. This special form of seepage is called wijst. De Burgt lies on top of the Peel Boundary Fault with two subsidiary faults along which wijst phenomena occur. The plan includes above-ground drainage of rainwater under gravity to infiltration zones, where it will have time to return as wijst water to the faults.

Deposits & Fossils:

River & Streams:

3. Geosite Interests


Cultural History

De Burgt represents a new step in cultural-historical development by making wijst visible, accentuating height differences, and carefully interpreting the contrasts between horst and graben with specific vegetation.

Flora & Fauna:


Industrial & Economic:

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures



Location of GS_24 De Burgt indicated with a light blue line. The fault identified by the investigation runs through the planned residential area.

) TNO Intern 149/219



AHN image of GS_24. The newly discovered fault segment shows a clear morphological expression here



Urban and landscape design of Plan De Burgt, in which the faults occupy a prominent position (source: https://nieuwblauw.nl/project/de-burgt-fase-1b).

) TNO Intern 150/219

Geosite: GS_25	Name
Significance: National	Donzel
Coordinates (RD): 165970, 413829	
Municipality	Bernheze
Province and country	Noord-Brabant, The Netherlands
Protection status	Designated as Wijst Reserve
Site management organizations	Staatsbosbeheer
Accessibility	Not accessible, but visible from the public road (Donzel and Donzelse Steeg).
Relevance for tourism	
Relevance for education & information	Viewing frame at Donzel to indicate the location of the fault.
Teaching equipment	

1. Location

This area is one of the five Wijst Restoration Projects designated by the province of North Brabant. After 2015, nine plots on the high side of the Peel Boundary Fault were purchased and redeveloped as a wijst reserve. In this former agricultural area, wijst values still need to develop further.

2. Landscape formation and description

Origin of the landscape

This geosite is located on the high side (footwall) of the Peel Boundary Fault, in an area that lies relatively higher than its surroundings due to fault movements. The fault line is clearly visible in the northern part of the area (a terrace step of about 1 m). Its position can also be inferred from a sudden change in vegetation and seepage phenomena in the ditches.

Geomorphology:

This plateau-like area was re-parcelled in the past, which largely erased the original cover sand morphology. In addition, the nutrient-rich topsoil was removed to a depth of about 30 cm to impoverish the soil and bring seepage closer to the surface..

Hydrology:

During land consolidation, deep drainage ditches were dug to drain the wet area. Recently, small weirs have been installed in these ditches to retain water on the high side of the fault. Seepage (wijst) is mainly recognisable by the rusty-brown water emerging here.

Deposits & Fossils:

Coarse, gravelly river sand deposited by the Meuse (Middle Pleistocene; Beegden Formation) is overlain by a 1–1.5 m thick layer of silty cover sand deposited by wind at the end of the last Ice Age (Late Pleistocene; Boxtel Formation).

) TNO Intern 151/219

River & Streams:

There is no natural drainage.

3. Geosite Interests

Cultural History

Flora & Fauna:

In de sloten en struweelranden staan wijsticonen zoals Klimopwaterranonkel, Moerasspirea, IJle zegge, Gewone vogelkers en Grote muur.

In the ditches and shrub edges grow wijst indicator species such as ivy-leaved crowfoot, meadowsweet, remote sedge, bird cherry, and greater stitchwort.


Industrial & Economic:

4. Relation with other geosites

This area is one of the five designated Wijst Restoration Projects in the province of Noord Brabant. Others include the Venloop valley near Slabroek (GS_13), the Wijstbos south of Zeeland (GS_19), the Sint Annabos in Uden (GS_07), and the source of the Esperloop in Gemert (GS_14).


5. Documentation & Resources

6. Maps, photos and figures



Location of GS_25 Donzel indicated with a light blue outline

) TNO Intern 152/219



AHN image of GS_25 Donzel. The height difference across the Peel Boundary Fault is about 1 m



Viewing frame at Donzel showing the location of the Peel Boundary Fault (photo: Thea Huijsmans).

) TNO Intern 153/219

Geosite: GS_26 Significance: Regional	Name De Specht-Putakker
Coordinates (RD): 177072, 399738	De Specifici diakkei
Municipality	Boekel
Province and country	Noord-Brabant
Protection status	Natuurnetwerk Nederland (NNN)
Site management organizations	De Specht-Handel
Accessibility	De Specht is easily accessible by car and bicycle.
Relevance for tourism	This geosite is part of the Breuken Beleven cycle route. The area features several themed walking trails, including the Specht Route, which explains the functioning of the Peel Boundary Fault.
Relevance for education & information	Natuurcentrum De Specht is a nature education centre (open by appointment) offering outdoor excursions, courses, and guided tours. The 'Breukenpad' (Fault Path) provides information about wijst phenomena. Viewing frame "De Elzen" is located near the Putakker.
Teaching equipment	?
Scientific interest	

1. Location

The Specht nature reserve is situated in the village of Handel, in a wooded area where many characteristic natural values remain intact.

2. Landscape formation and description

Origin of the landscape

The area around De Specht was formed by the activity of three fault segments of the Peel Boundary Fault. Wijst phenomena, terrace steps, groundwater springs, and vegetation differences make this evident. The nature reserve was established in the early 1990s and enhanced by sowing a native herb mixture of about thirty flower species. The Breukenpad starting at Natuurcentrum De Specht clearly reveals the various manifestations of the faults.

Geomorphology:

Around Natuurcentrum De Specht, the height differences caused by movement along the Peel Boundary Fault and the Handel Fault are clearly visible. This height difference is highlighted by the viewing frame at the Putakker.

Hydrology:

Natuurcentrum De Specht lies on top of the Peel Boundary Fault with two subsidiary faults along which wijst phenomena occur. These are explained along the 'Breukenpad'. At the Putakker viewing frame, pools with different groundwater levels can be seen on either side of the Peel Boundary Fault. Due to seepage, the higher ground has a higher groundwater level than the lower ground, caused by

) TNO Intern 154/219

the impermeable nature of the fault.

Deposits & Fossils:

Coarse, gravelly river sand deposited by the River Meuse (Middle Pleistocene; Beegden Formation) is overlain by a 1–2 m thick layer of silty cover sand deposited by wind at the end of the last Ice Age (Late Pleistocene; Boxtel Formation). Locally, this sand was reworked by wind during the Holocene induced by human activity.

River & Streams:

The Landmeerse Loop flows through the nature reserve, crossing the faults. Within the reserve, several hundred metres of the stream remain unchannelled. Because the area suffers from drought, small dams have been installed in the Landmeerse Loop.

3. Geosite Interests

Cultural History

Flora & Fauna:

The great diversity of landscapes within De Specht results in a wide variety of flowering plants. Different species thrive in the woodland compared to the pond edges or the sunlit flower meadows. Twenty-five years after its creation, the area now hosts more than 200 species, including wild marjoram, wild teasel, burdock, garden burnet, and marsh orchid.

The section of the Landmeerse Loop within the reserve is unchanneled for several hundred metres and is bordered by willow thickets. Reed growth along the fault is characteristic. In the marshy area and along the stream banks, many wijst indicator species occur, such as field rush, cuckooflower, ragged robin, marsh thistle, marsh orchid, greater bird's-foot trefoil, valerian, alder, bird cherry, guelder rose, and hop.

The Putakker viewing frame illustrates the differences in vegetation between plants growing on the wet horst and those on the dry graben.

The diversity of habitats around Natuurcentrum De Specht supports a rich fauna as well. Numerous rare insects can be observed. Kingfishers, icterine warblers, whitethroats, and sparrowhawks are frequently seen, while frogs, toads, and newts are abundant. The bat bunker shelters a large bat population. Visitors may also encounter squirrels, roe deer, rabbits, hares, mice, and, with luck, a fox or polecat.


Industrial & Economic:

4. Relation with other geosites

De Perekker (GS_27) in Boekel

- 5. Documentation & Resources
- 6. Maps, photos and figures

) TNO Intern 155/219



) TNO Intern 156/219



Viewing frame at the Putakker (see map for location; source: peelhorstenmaasvallei.nl)

) TNO Intern 157/219

Geosite: GS_27	Name
Significance: Regional	De Perekker
Coordinates (RD): 175645, 401071	
Municipality	Boekel
Province and country	Noord-Brabant
Protection status	?
Site management organizations	Stichting D'n Eik Boekel
Accessibility	he De Perekker nature trail is easily accessible by car and bicycle via Mezenlaan in Boekel.
Relevance for tourism	The geosite is part of the walking route 'Ommetje Peelrandbreuk Boekel' (developed by the Boekel Local History Society).
Relevance for education & information	This nature centre and park features a bee hall, insect wall, frog pond, ponds with waterfall, bat cellar, and sand martin wall, as well as an interpretive nature trail where the wijst phenomenon and the Peel Boundary Fault are visible.
Teaching equipment	
Scientific interest	

1. Location

lose to the centre of Boekel lies the nature park De Perekker, a unique area of woodland, water, and sand drifts. Within this area runs a subsidiary fault of the Peel Boundary Fault, associated with wijst phenomena.

2. Landscape formation and description

Origin of the landscape

The De Perekker nature reserve is intersected by the Peel Boundary Fault. Wijst phenomena, terrace steps, groundwater springs, and vegetation differences make this evident. The reserve was established in the early 1990s and enhanced by sowing a native herb mixture of about thirty flower species. The fault trail starting at Natuurcentrum De Specht clearly reveals the various manifestations of the faults.

Geomorphology:

Around the nature centre, the height differences caused by movement along the Peel Boundary Fault and the Handel Fault are clearly visible. The terrace step in the watercourse along the Perekker path also indicates the position of the fault.

Hydrology:

The Perekker area east of the fault has a higher groundwater level due to seepage, caused by the impermeable nature of the fault. Seepage water flows over the fault into the lower Perekker area, where it feeds the ponds. The wijst water is artificially impounded in a large skating pond within the reserve. The weir, located on the fault, clearly shows the difference in water level above and below

) TNO Intern 158/219

the fault.

Deposits & Fossils:

River & Streams:

The Landmeerse Loop flows through the nature reserve, crossing the faults. Within the reserve, several hundred metres of the stream remain unchannelled. Because the area suffers from drought, small dams have been installed in the Landmeerse Loop.

3. Geosite Interests

Cultural History

The 'Ommetje Peelrandbreuk Boekel route links the centre of Boekel with the historic Huize Padua, passing characteristic locations that have shaped both the past and the lives of local residents. Points of interest along the route include clay pits where bricks were made, centuries-old farms along the fault line, an ancient ridge-and-furrow field complex, and the Klein Rondeel, notable for its lower ground level due to sand extraction.

Flora & Fauna:

In the watercourse along the Perekker path grow several wijst indicator species such as field rush, horsetail, marsh thistle, and floating sweet-grass. On the banks grow cow parsley, bulbous buttercup, and sorrel, which no longer thrive on the drier ground further away. Along the path, alders suffer from drought, while around the fault, species typical of bird cherry—ash woodland replace the alder carr, including bird cherry, guelder rose, hazel, and ash. Hornbeam even self-seeds here. The small reserve is surrounded by encroaching urban development. The Dooleggen housing estate, situated above the Peel Boundary Fault, is bordered by ditches containing orange-brown seepage water.

Industrial & Economic:


4. Relation with other geosites

On 22 May 2025, mechanical test drillings (up to 8 m deep) were carried out at the De Burgt site on the northern edge of Boekel (GS_24) to determine the exact position of the Peel Boundary Fault. This concerns a branch located further north than the one at De Perekker (see map).

5. Documentation & Resources

6. Maps, photos and figures

) TNO Intern 159/219



) TNO Intern 160/219

Geosite: GS_28	Name
Significance: Regional	Willibrordusput Meijel
Coordinates (RD): 188930, 373879	
Municipality	Deurne, Peel en Maas
Province and country	Noord-Brabant, Limburg
Protection status	The well and the boundary stone are designated as
Trottetion status	National Monuments (Rijksmonument).
Site management organizations	
Accessibility	Hof, 5768 RX Meijel
Relevance for tourism	De Wegh van Meijl op Seven in Someren is a 21 km route full of legends, marked by basalt columns and starting at the Saint Willibrordus Well.
Relevance for education & information	Informationpanel
Teaching equipment	

1. Location

The Saint Willibrordus (or Wilbers) Well is located in the Limburg village of Meijel, precisely on the border between the provinces of North Brabant and Limburg. Next to the well stands a boundary stone dating from 1761.

2. Landscape formation and description

Origin of the landscape

According to tradition, Saint Willibrord (the Apostle of the Netherlands), who was active in the eighth century in what is now Noord Brabant as a missionary, is said to have dug this well. That a well could be dug at this relatively high elevation (35 metres) was due to the presence of the Peel Boundary Fault. It is possible that Willibrord travelled from Deurne along this route to Echternach, noticed the wijst water, and decided to dig a well here.

Geomorphology:

The location of the Saint Willibrordus Well is relatively high, but the Peel Boundary Fault lies about 300 m further north. Due to cover sand deposits, the terrace step at the fault is no longer visible. The terrace step just south of the well is attributable to an aeolian landform.

Hydrology:

Here, the hydrological situation appears reversed: the wet areas lie below the Peel Boundary Fault, and the dry areas above. This is explained by the presence of cover sand deposits over the Peel Boundary Fault, which almost eliminated the fault-related height difference. The clay and loamy sand at the base of the cover sand prevent water from draining easily. The near absence of wijst phenomena with iron-rich seepage is also atypical.

Deposits & Fossils:

Cover sand ridge

) TNO Intern 161/219

River & Streams:

3. Geosite Interests

Cultural History

The Willibrordus Well lies along a passage through the Peel. The nearby road through the Peel is said to have been used by Saint Willibrord, travelling from Utrecht via his estates in Bakel and Deurne on his way to his monastery in Echternach. Here, according to legend, he and his companions could rest, preach, and baptise people. A large oak tree stood by the well on the height, serving as a landmark for boundary demarcation as early as before 1300. This oak remained a reference point until the 19th century. In 1352, the well, clearly visible on this high point, was used as an orientation marker. About 150 metres away lay the boundary between Brabant and the Land of Weert. In 1452, the area was referred to as Luttel Meijdel, where Meijdel denotes a height amid marshy land. Users from both sides had boundary disputes here. In 1549, a boundary stone was placed to mark the border between municipalities such as Someren and Nederweert, which often clashed. The actual boundary still lay 150 metres further on. On 12 May 1761, a new boundary stone was erected, now marking the border between the Dutch Republic and Austrian Upper Guelders. One side bears the Dutch lion, the other the Austrian double-headed eagle. The water of the well was considered medicinal, making it a place of pilgrimage. In 1742, it was believed to protect against marsh fever. However, the well was poorly maintained and, after 1800, fell into such disrepair that nothing remained visible. Under the parish priest's leadership, the well was rediscovered and re-excavated on 6 July 1899. A round structure was built over it, which still exists. On 8 October 1899, a procession went to the well, which was then re-consecrated. It became a pilgrimage site, and the water, now regarded as an excellent remedy for eye diseases, was taken home in bottles.

The well was damaged during the Second World War but restored in 1953. A bust of Willibrord was placed, though it was not weather-resistant and eventually moved to the church. In 1979, the well was purchased by the municipality of Meijel, and a new bust was installed.

Flora & Fauna:

Industrial & Economic:


4. Relation with other geosites

The geological setting is comparable to Geosite GS_40 Waterbloem.


5. Documentation & Resources

6. Maps, photos and figures

) TNO Intern 162/219



) TNO Intern 163/219



) TNO Intern 164/219

Geosite: GS_29	Name
Significance: Regional	Schuitwater
Coordinates (RD): 206579, 387302	
Municipality	Horst aan de Maas
Province and country	Limburg
Protection status	Aardkundige waarde, Natuurnetwerk Nederland (NNN)
Site management organizations	Staatsbosbeheer
Accessibility	Accessible from the car park between Broekhuizen and Horst/Melderslo.
Relevance for tourism	Staatsbosbeheer has established the N-95 walking route here. This 9.3 km walk starts at the car park along the Horsterweg. The route and information about flora, fauna, and nature management are displayed on information panels. Several routes of the 'Stempel van de Maas' (later integrated into the KnopenLopen network of Horst aan de Maas) also pass through this nature reserve. Some marshy sections are crossed by boardwalks. The Pieterpad long-distance trail also runs through the area.
Relevance for education & information	Informationpanel "Schuitwater – Stempel van de Maas"; IVN organizes educational walks.
Teaching equipment	
Scientific interest	Woolderink et al., 2018
Description	

1. Location

Schuitwater is a 313-hectare nature reserve in northern Limburg. It lies between the villages of Swolgen, Broekhuizenvorst, and Broekhuizen, and west of Lottum. The area consists of remnants of an abandoned Meuse meander between Broekhuizen in the north and Lottum in the south. The total length of the old river arm is about 7 km, whereas the present Meuse course over this stretch is only 3 km long. The area lies within the Venlo Block.

2. Landscape formation and description

Origin of the landscape

During the Allerød period (10,950–11,900 years ago), the River Meuse flowed here, cutting a large meander through the landscape and forming the middle terrace. The Meuse later broke through the meander and shifted its course during the Younger Dryas. During this cold phase, the meandering river changed into a braided system with a wide riverbed that cut across the old channel (see Woolderink et al., 2018).

Geomorphology:

It is a wooded and water-rich area, surrounded by higher sandy soils with pine trees, sand dunes, heathland, and birch woods.

) TNO Intern 165/219

Hydrology:

Deposits & Fossils:

River & Streams:

3. Geosite Interests

Cultural History

Flora & Fauna:

Few nature reserves offer as much variety in such a small area as Schuitwater: heathlands, dry and wet grasslands, and water bodies. The diversity of the landscape is linked to numerous elevation differences. The low-lying core of the reserve consists of old channels abandoned by the Meuse, flanked by high, dry sandy soils.

Industrial & Economic:

The Schuitwater reserve is a remnant of several former Meuse channels. These gradually silted up, and fen peat formed on the bottom. From the 18th century, peat was dredged from the Schuitwater channels and transported in flat-bottomed boats. After peat extraction ceased, the silting-up process resumed.

4. Relation with other geosites

Geosite GS_36 – De Vilt shows an abandoned Meuse meander of the same age.


5. Documentation & Resources

6. Maps, photos and figures



Location of GS_29 Schuitwater indicated with a light blue outline

) TNO Intern 166/219



Left: AHN image of GS_29 Schuitwater. Right: Palaeogeography of the Meuse Valley (Woolderink et al., 2018) showing the abandoned Allerød Meuse meander at Schuitwater.

) TNO Intern 167/219

Geosite: GS_30	Name
Significance: Regional	De Bult
Coordinates (RD): 186902, 390007	
Municipality	Deurne
Province and country	Noord-Brabant (NL)
Protection status	Natura-2000
Site management organizations	Staatsbosbeheer
Accessibility	
Relevance for tourism	It is now prohibited to give peat-cutting demonstrations here. These are still provided on a peat field located on a Staatsbosbeheer plot at the Toon Kortooms Park (GS_04).
Relevance for education & information	
Teaching equipment	

1. Location

The wetland nature reserve De Bult covers 120 hectares in the far north-east of Deurne, behind the Craijenhut, on the border between the provinces of North Brabant and Limburg. Although it does not directly adjoin it, the site is part of the Natura 2000 area Deurnsche Peel & Mariapeel, but lies outside its core zone. The reserve is completely surrounded by farmland. To the east flows the Defensiekanaal, and to the north of De Bult begins the Kaweise Loop.

2. Landscape formation and description

Origin of the landscape

Like the entire Natura 2000 area Deurnsche Peel & Mariapeel, De Bult is a raised bog area that remains after historical reclamation. It is an isolated remnant of the Deurne Peel. The name "De Bult" derives from the raised bog hummocks characteristic of the area. These peat bogs were largely excavated down to the sandy subsoil. These areas are the southernmost representatives of the flat sub-Atlantic raised bogs, which elsewhere—and also in the Peel region—have largely disappeared due to excavation, reclamation, and peat cutting.

Geomorphology:

Hydrology:

To the east of the area flows the Defensiekanaal, and to the north begins the Kaweise Loop. Although De Bult is designated as a wetland nature pearl, the area is threatened by the planned reopening of the nearby De Peel Air Base, which has led to protests from Werkgroep Behoud de Peel (WBdP). To keep this higher-lying area wetter, the water management system needs significant adjustment.

Deposits & Fossils:

) TNO Intern 168/219

Within the aUGGP, De Bult represents an undeveloped area where remnants of sphagnum moss are still present. It forms part of a more or less continuous remnant of the largely vanished raised bog landscape of the Peel, located on the watershed between the Brabant river Aa in the west and the Limburg Grote Molenbeek in the east. Most of the peat was removed in the past, but locally thick layers of residual peat remain. In some complexes of peat pits, vegetation with hummock-forming raised bog species occurs.

River & Streams:

In this area, watercourses have been made shallower, boundary ditches filled in, and weirs and controlled drainage installed to preserve and, where possible, restore remnants of raised bog, birch woodland, and dry and wet heathland, while also considering agriculture, residents, and cultural history.

3. Geosite Interests

Cultural History

Archaeological finds, in the form of flint knives, scrapers, and points, indicate very early human presence.

Flora & Fauna:

De Bult is a remnant of the former extensive raised bog marshes. Characteristic species such as small cranberry, bog rosemary, and hare's-tail cottongrass still grow mainly in and along peat pits. The aim is for De Bult to consist largely of regenerating and partly active raised bog in the future, with birch carr and birch-oak woodland mainly in the marginal zones. In some places, original vegetation is recovering thanks to improved water management, and sphagnum mosses, cotton grass, and sundew are reappearing.

Due to the varied history of peat cutting in different parts of the area, there is a high degree of fine-scale variation in vegetation and landscape, with gradients towards slightly more mineral-rich conditions. In the oldest peat pits, miniature raised bog growth has been occurring for a long time. On the larger residual peat units, relatively thick peat layers remain, and restoration management has already led to the development of raised bog vegetation in several places.

Important animal species in the area include the badger and the smooth snake.

Industrial & Economic:


4. Relation with other geosites

Like the entire Natura 2000 area Deurnsche Peel & Mariapeel (GS_11), De Bult is a raised bog area that remains after reclamation. One kilometre to the north lies De Peel Air Base, which still contains important natural values such as heathlands, and two kilometres further north lies the Stippelberg estate (GS_33), also with some heathland. About half a kilometre to the south-east is the Peel remnant Heidsche Peel.

5. Documentation & Resources

6. Maps, photos and figures

) TNO Intern 169/219



Location of GS_30 De Bult indicated with a light blue outline



De Bult still provides a good impression of what the vast and open Peel must once have looked like

) TNO Intern 170/219

Geosite: GS_31	Name
Significance: Regional	Loobeekdal
Coordinates (RD): 192726, 392546	
Municipality	Venray
Province and country	Limburg
Protection status	The area is part of the Dutch National Nature Network (NNN) and falls under the European Water Framework Directive (WFD), the Water Act (Waterwet), and the
Site management organizations	Water Management Plan (Waterbeheerplan). Staatsbosbeheer, Waterschap Peel en Maasvallei, Stichting Loobeekdal
Accessibility	
Relevance for tourism	There are several walking routes through the nature reserve starting from Merselo. It is a wetland area with unpaved paths.
Relevance for education & information	Stichting Loobeekdal organises guided walks covering history, culture, nature, and landscape. A sheepfold combined with an information centre is under development.
Teaching equipment	

1. Location

The Loobeek flows south of Merselo (Haag–Weverslo) through a redesigned section of the Loobeek Valley. Here, geology (Tegelen Fault and Peel Horst), nature, culture, and history converge within a few hundred metres. The Weverslose Schans lies in the Loobeek Valley.

2. Landscape formation and description

Origin of the landscape

The Loobeek crosses the Tegelen Fault Zone south of Merselo. The Tegelen Fault here forms a so-called "relay" zone where the southern fault segment steps over to a more northerly segment. The Loobeek also crosses the northern segment west of Venray. Originally a tributary of the River Meuse, the Loobeek rises in the Peel. In the 20th century, the stream was adapted for agricultural drainage.

Geomorphology:

The Tegelen Fault lies on the southern side of the valley; the left bank has been raised by windblown sand.

Hydrology:

The Loobeek Valley is a mixed-water area where rainwater combines with groundwater. This groundwater is iron-rich, as evidenced by various wijst phenomena.

Deposits & Fossils:

River & Streams:

) TNO Intern 171/219

Today, the stream originates northwest of Ysselsteyn as an inlet from the Peelkanaal (Defensie Kanaal) and flows northeast, initially through large-scale agricultural reclamation. South of Merselo, the landscape becomes more small-scale. Here lies the Loobeek Valley nature reserve with the Weverslose Schans. In this section, the stream follows a more natural course. The valley has been redesigned in recent years (completed in 2015) to allow nature to recover. North of Merselo, however, lies an area where land consolidation in the 1960s and 1970s resulted in heavy canalisation of the stream and large-scale alteration of the valley landscape. Ultimately, the Loobeek joins the Afleidingskanaal between Venray and Smakt and continues under that name towards the Meuse. At this point, effluent from the Venray wastewater treatment plant is also discharged.

3. Geosite Interests

Cultural History

During the final phase of the Second World War, when Allied forces advanced after the Battle of Overloon to push from Overloon to Venray, they had to cross the Loobeek with tanks. At that time, the stream had swollen to a width of six metres. The Germans had laid mines in the river. The crossing took place on 16 October 1944 and involved heavy losses, after which the Loobeek was nicknamed "Blood Brook".

The Weverslose Schans was a farmers' redoubt located south of Merselo and north of Weverslo in the Loobeek Valley. The exact year of construction is unknown. It was used by local inhabitants as a refuge during times of war and roaming bands of looters. Around 1925, the old Weverslose Schans was levelled during the reclamation of the Peel. Not far from the original site, the redoubt was reconstructed in 2022 by Stichting Loobeek. The fortification consists of an earthen wall with a pentagonal plan, corner bastions, and a watchtower. Around it lies a water-filled moat fed by the Loobeek. The Weverslose Schans lies exactly on the southern branch of the Tegelen Fault. Sheep have always been important to the inhabitants of the Peel region, providing wool for clothing and, above all, manure to enrich the poor sandy soils. The new Weverslose sheepfold being built in Merselo will serve a dual purpose: occasionally housing sheep and functioning as an information centre focusing on nature, landscape, and the history of the sheep companies, the fulling mill, and the redoubt.

The wool from sheep grazing in the Peel was partly traded, spun, woven, and processed into felt. Felt was produced in a fulling mill, where wool was pounded for hours by water-powered wooden hammers, creating felt for clothing. The reconstructed fulling mill in Merselo is powered by water from the Loobeek and is located at the point of greatest fall.


Flora & Fauna:

The site showcases flora and fauna and, together with the Weverslose Schans and fulling mill, illustrates how past inhabitants utilised the geological setting. The area is an excellent example of how nature conservation and ecological development can go hand in hand. The Loobeek Valley contains wet grasslands and groves with alder and willow. On drier parts, birch and coniferous trees grow. The valley harbours several rare plants that contribute to biodiversity, including various orchids (such as the broad-leaved helleborine), marsh marigold, and marsh violet. The area is home to birds such as the kingfisher and grey wagtail, mammals like roe deer and foxes, and provides habitat for amphibians and reptiles, including the common toad and grass snake.

<u>Industrial & Economic:</u>

- 4. Relation with other geosites
- 5. Documentation & Resources

) TNO Intern 172/219



) TNO Intern 173/219



The Loobeek (foto: www.waterschaplimburg.nl)

) TNO Intern 174/219

Geosite: GS 32	Name
Significance: National	Maaspark Noord-Limburg
Coordinates (RD): 205555, 392170	
Municipality	Horst aan de Maas, Venray
Province and country	Limburg
Protection status	Gebiedsontwikkeling Ooijen-Wanssum
Site management organizations	stichting Maaspark Noord-Limburg
Accessibility	Maaspark Ooijen-Wanssum is freely accessible throughout the year. Facilities for people with reduced mobility are also available. The park boasts an extensive network of paths and roads, yet it also invites visitors to explore off the beaten track. The area has no fewer than 68 entrances, including seven distinctively designed 'escarpment gates' (steilrandpoorten).
Relevance for tourism	Pedestrians and cyclists can experience the area to the fullest. There are 13 kilometres of cycle paths and 30 kilometres of walking trails. The park has its own cycling route and several circular walks in and around the villages.
Relevance for education & information	
Teaching equipment	A special teaching package, The Water World between Ooijen and Wanssum, has been developed for primary school children. Educational canoe trips are also available.
Scientific interest	Reconstruction of former Meuse river courses by Woolderink et al., 2018
Description	

1. Location

The Groote Molenbeek is an old branch of the River Meuse, approximately 32 kilometres long, flowing through Maaspark Noord-Limburg. It meanders through the landscape towards Wanssum, where it joins the Meuse. With 540 hectares, the Maaspark is the largest contiguous area of Meuse nature in North Limburg. The original Meuse terrace landscape is central here, and the influence of water is evident everywhere. The ten-thousand-year-old river courses of the Meuse are still recognizable and now form a domain of water in many forms. The park contains a wide variety of landscapes.

2. Landscape formation and description

Origin of the landscape

About 10,000 years ago, during the last cold period (Ice Age), the Meuse flowed through the area as a braided river. It cut into older Meuse deposits of the middle terrace, creating the lowest terrace, about 2 to 3 metres above the current river. The river valley was very wide, with multiple channels and islands in between. This braided pattern gave the lowest terrace a varied relief. After the last Ice Age, the Meuse began to meander again and assumed roughly its present single course. The channel

) TNO Intern 175/219

between Ooijen and Wanssum was abandoned, but the lowest terrace still floods relatively quickly.

Geomorphology:

Cover sands and drifting dunes also formed during the last cold period, covering large parts of the middle terrace. That cold period briefly turned much of the Netherlands into a polar desert. Severe drought, lack of vegetation, and stormy westerly winds caused major sand drifts. Thick layers of cover sand reached the Meuse from the west, and high river dunes blew up from the riverbed. The two terraces, combined with the drifting sand, define the current landscape of the old Meuse branch. The higher grounds in the planning area belong to the middle terrace. These higher grounds have been cut through again by a network of channels forming part of the lowest terrace. A section of the higher middle terrace became isolated between the Meuse and the old branch. River dunes also blew up from this branch, explaining the presence of Meuse dunes on the western bank between Ooijen and Blitterswijc

Hydrology:

This section of the Meuse as serves as a floodplain during extreme water levels to prevent greater disasters downstream in the low-lying parts of the Netherlands. From this perspective, the North Limburg Meuse valley is considered a retention area. This makes Maaspark Noord-Limburg different from other nature reserves, as it was designed to protect against high water levels of the Meuse. When water levels rise, the high-water channels fill first, and if the water continues to rise, the entire area floods—a spectacular sight. During the high water of July 2021, Maaspark Ooijen-Wanssum was severely tested immediately after completion. The design proved highly effective in accommodating high Meuse water levels and reducing water levels downstream.

Deposits & Fossils:

River & Streams:

Closer inspection reveals that the channel pattern of the old Meuse branch is more extensive and irregular than it first appears. The higher grounds between Ooijen and Blitterswijck and the area between Broekhuizenvorst, Ooijen, and Legert are intersected by several historic channels. Many current streams follow these depressions, such as the Smallenbroek, Wolterskamp, and Boddebroek.

3. Geosite Interests

Cultural History

This area was shaped by the Meuse and by human activity. Originally, there were few settlements, and habitation was limited to the higher ridges between the channels.

Castle Ooijen dates back to before 1433, though the current buildings are from the 17th century and later. It lies in a freely accessible new nature reserve, after the old Meuse branch was restored at the end of 2020. The castle remained inhabited until 1981 and has been listed as a monument since 1968. Since 1989, the once-derelict castle has been owned by the Ooijen Castle Foundation, which restored it. The restoration was completed in 2003. Over the years, the moats were partly filled in and relocated, so the castle is now only partially moated.

Flora & Fauna:

Maaspark Ooijen-Wanssum allows natural processes and diverse flora and fauna to thrive. Beavers, badgers, and foxes are already frequent visitors. Grazing animals maintain the open landscape, ensuring water has enough space during floods. The area supports geese, herons, and even ospreys and kingfishers. The vegetation includes various grasses, willows, and plants such as yellow rattle

) TNO Intern 176/219

and devil's-bit scabious. In the early years, waders like the little ringed plover can also be seen along the still-bare channel banks.

Industrial & Economic:

4. Relation with other geosites


Old Meuse Meander Schuitwater (GS_29). The Groote Molenbeek also flows through De Elsbeemden (GS_42).

5. Documentation & Resources

6. Maps, photos and figures



Location of GS_31 Loobeek indicated with a light blue outline



Left: AHN image of the old Meuse branch Ooijen-Wanssum. Right: Reconstruction of old Meuse courses (Woolderink et al., 2018). Green – meandering Meuse before the last Ice Age; Yellow – braided Meuse during the last Ice Age; Orange – meandering Meuse after the last Ice Age

) TNO Intern 177/219

Geosite: GS_33 Significance: International	Name Stippelberg (Klotterpeel, de Rips)
Coordinates (RD): 182640, 393777 Municipality	Gemert-Bakel
Province and country	Noord-Brabant
Protection status	Natuurmonument
Site management organizations	Gem. Boekel (opdrachtgever)
Accessibility	Via the Bosmuseum, near the Beestenveld car park
Relevance for tourism	At the edge of the Stippelberg nature reserve lies Natuurpoort Nederheide, an excellent starting point for walks. The Bosmuseum in De Rips also offers various starting points for walks in the woodland area
Relevance for education & information	At the Bosmuseum in De Rips, the rich history of the Peel village is presented through information panels and visual material. There is an archaeological monument, Stone Age, on the Stippelberg, and an information panel near the Tjongerculture finds on the Stippelberg, Oude Bakelsedijk.
Teaching equipment	
Scientific interest	Archaeological research on a heathland site at "De Stippelberg". KWR research into the desiccation of the Stippelberg.

1. Location

The Stippelberg lies on the Peel Horst and is intersected by the Milheeze Fault (a branch of the Peel Boundary Fault). The Stippelberg is the largest forested area in the province of Brabant; only the Veluwe is larger in the Netherlands. On the Peel Horst there are two wet areas: the Klotterpeel and De Rips. In the southern part of the Stippelberg is a small wijst area just east of the fault.

2. Landscape formation and description

Origin of the landscape

Stippelberg and the reclamation village of De Rips are characterised by a striking feature: the fault line running through the area. This is the Milheeze Fault, a branch of the Peel Boundary Fault. The fault line causes major differences in groundwater conditions, resulting in varied vegetation. The cover sand layer deposited during the last Ice Age on the Peel Boundary Fault is about two metres thick, but in the middle of the horst it is only a few decimetres thick, and in wind-eroded depressions it is sometimes absent, exposing Meuse sediments recognisable by coarse sand with pebbles. Where seepage water reached the surface, drifting sand was stabilised, forming elevations.

Geomorphology:

The Stippelberg owes its name to the many small drifting sand dunes in the landscape. Because the drifting sand was trapped, the depression beneath the fault was not completely filled, creating

) TNO Intern 178/219

slopes. Differences in subsidence rates along the fault may also have caused these slopes. The result is a sand dune ridge bisected by the fault.

Hydrology:

The Klotterpeel is a large shallow fen situated above the Milheeze Fault. Beestenveld lies on the northern side of the Stippelberg. The Koorden Canal, which runs through it, is reportedly not intersected by the Milheeze Fault according to Witteveen and Bos. However, halfway along the Koorden Canal there is a weir, and a water-bearing ditch runs across the fields towards the Middenpeelweg. A strip of reeds grows along the verge of the Middenpeelweg. It is assumed that the Milheeze Fault crosses near the Beestenveld weir towards the Middenpeelweg.

Deposits & Fossils:

River & Streams:

3. Geosite Interests

Cultural History

Most of the current forest area of the Stippelberg is a well-preserved example of a production forest dating from the late 19th and early 20th centuries, with typical grid parceling, avenue planting, and drifting sand dunes stabilized by conifers. Important historic heath paths, aligned with the church towers of surrounding villages, cross the area. From the northern corner of the forest, a zigzag line runs southwards: a precursor of the Peel-Raam Line (a defensive structure), a rare example of an intact anti-tank ditch from 1939.

The Klotterpeel was originally a small peat-cutting area, and traces of peat extraction remain. The reclamation village of De Rips lies on the north-eastern edge of the Stippelberg and was created through organized reclamation of heath and peatlands in the 19th and 20th centuries. A forest museum illustrates this history. Nearby are other reclamation villages such as Elsendorp and Vredepeel.

Human habitation in the De Rips area goes back much further, as artefacts from the Tjonger Culture have been found here. The Tjonger people lived about 13,000 years ago during the Stone Age. They made tools and arrowheads resembling pocket knives. In the Netherlands, they are also called Tjonger people because the first finds were made near the River Tjonger in Friesland. They lived at the end of the last Ice Age, in a warmer period, though not as warm as today. The landscape changed from tundra with sparse vegetation to open woodland with heath and grassland. These people lived by hunting, fishing, and gathering plant foods and small animals. Remarkably, they often returned to camp here, probably because they could set up camp high and dry halfway up a cover sand ridge.

The first excavation took place in the 1950s by Dr Bohmers of the University of Groningen. The State Service later investigated the area, and amateur archaeologists also made many finds. Thousands of flint artefacts have been found, such as points, scrapers, burins, blades, and drills used as personal tools. This site has been an Archaeological National Monument since 1971 and remains one of the most important Stone Age sites in the Netherlands and north-western Europe. The site is 100 to 150 metres wide and no less than two kilometres long.

Flora & Fauna:

The Klotterpeel hosts a large population of moor frogs, which mainly occur in nutrient-poor, slightly acidic waters. However, seepage indicators such as many-stalked spike-rush, winter purslane, pillwort, arious-leaved water-starwort, and shoreweed show that it is a buffered fen. Above the weir in the Koorden Canal grow species typical of base-rich soils (hard fern and male

) TNO Intern 179/219

fern), while the Beestenveld forest is nutrient-poor and characterized by frogbit, milk-parsley, common skullcap, lesser water parsnip, royal fern, male fern, lady fern, flowering rush, and various-leaved water-starwort. In the wijst area of De Kaak grow sharp-flowered rush, marsh St John's wort, and greater bird's-foot trefoi.


Like many Dutch forests, the Stippelberg was originally a production forest with low biodiversity. Natuurmonumenten is working to make the forest more natural, increasing tree species diversity and creating more open spaces and connections between heathlands.

Industrial & Economic:

Thousands of years ago, predecessors of the Meuse and Rhine flowed through East Brabant, leaving river deposits rich in gravel. Because the Bakelse Plassen lie on the elevated Peel Horst, which has been uplifted by the Peel Boundary Fault, these river deposits are close to the surface. This is why gravel extraction has taken place here since 1986, creating an ever-expanding lake that continues to grow today.


- 4. Relation with other geosites
- 5. Documentation & Resources

6. Maps, photos and figures



Location of GS_33 Stippelberg indicated with a light blue outline. All red faults shown, except the Tegelen Fault, are branches of the Peel Boundary Fault Zone

) TNO Intern 180/219



AHN image of GS_33 showing terrain steps along the faults of the Peel Boundary Fault Zone around the Stippelberg



Wijst area De Kaak (indicated on the map)

) TNO Intern 181/219

Geosite: GS_34	Name
Significance: Regional	't Zinkske, Marisberg, Molentje
Coordinates (RD): 191634, 375805	
Municipality	Deurne, Peel en Maas
Province and country	Noord-Brabant, Limburg
Protection status	Natura-2000
Site management organizations	Staatsbosbeheer, private oensers
Accessibility	Visitors can enter 't Zinkske via a small gate. The path contains a striking amount of rubble, originating from the church in Meijel that was destroyed during the war. The canal banks provide pleasant walking routes, although 't Zinkske is largely fenced to allow sheep grazing.
Relevance for tourism	The Peel remnant 't Zinkske is part of the Geopad Verheven Peel. The canal banks offer walking opportunities.
Relevance for education & information	Differences in groundwater levels and vegetation on either side of the Peel Boundary Fault are illustrated here.
Teaching equipment	

1. Location

This geosite comprises the Peel remnants 't Zinkske, 't Molentje, and Marisberg. The elevated peatlands ("Verheven Peel") of the Deurnese Peel, Mariapeel, and 't Zinkske are bordered by the Deurne Canal. On the west side of the canal, in the seepage zone, lie wet meadow areas: Leegveld and 't Molentje.

't Zinkske is a 130-hectare Peel remnant located between Neerkant and Helenaveen in the municipality of Deurne. It forms a small, separate part of the much larger Deurnese Peel. The Zinkske lies on the provincial border with Limburg, just north of the point where the Helenavaart and the Deurne Canal meet.

2. Landscape formation and description

Origin of the landscape

Peel remnant with hydrological nature restoration. 't Zinkske is one of the remaining fragments of the once vast Peel bog. Traces of the earliest peat extraction remain here in so-called boerenkuilen (farmers' pits): peat pits dug in a single day (and refilled with water the next day). These pits are important because they still host true raised bog plants.

The Marisberg woodland is the highest area of the Peel Horst, lying four to five metres higher than 't Molentje. The microrelief is striking: small variations in elevation formed in two phases. First, during the Ice Age, wind created a gently undulating sandy landscape. Later, in the Middle Ages, deforestation caused renewed sand drift, which was later stabilised by reforestation.

During the middle part of the Ice Age, the River Meuse still flowed through what is now 't Molentje, carving a broad, shallow valley that was later filled with raised bog. Seepage water from the nearby Marisberg, 4–5 metres higher and located on the other side of the Peel Boundary Fault, was blocked

) TNO Intern 182/219

by the impermeable fault, creating marshy conditions. This is demonstrated by a double groundwater gauge.

Geomorphology:

Hydrology:

The Marisberg is important for the Peel's water system because it lies at a watershed junction: one north—south through the Mariapeel, another along the Noordervaart.

Peat formation at 't Molentje is explained not only by the presence of the old Meuse valley but also by seepage against the Peel Boundary Fault. Together, they form a large wijst area. The seepage originates from the Marisberg, where rainwater infiltrates, flows over a relatively low-lying impermeable layer (Breda Formation) westwards, and resurfaces as seepage because it is forced upward against the impermeable Peel Boundary Fault. This infiltration—seepage system is explained and visualised by a double groundwater gauge showing both seepage water (in deeper layers) and groundwater (in upper layers). Between them lies a thin impermeable layer, probably formed on the floor of the former Meuse valley, creating a perched water table.

The excavated Blankerplas in the Molentje area is also fed by seepage water.

Deposits & Fossils:

River & Streams:

3. Geosite Interests

Cultural History

The Zinkske is one of the remaining Peel remnants of the once immense Peel bog. Traces of the earliest peat extraction remain in boerenkuilen, which are important because they still support true raised bog vegetation.

The Marisberg area is relatively high, and several Roman finds have been made nearby, suggesting possible agricultural use in Roman times. There may still be remnants of an ancient agricultural landscape beneath the sand drifts. Further research (deeper drilling) is needed to confirm this.

Flora & Fauna:

Alongside the cut-over peat plain, now covered with heath, a young birch forest grows, and remnants of an old peat canal (het wiekie) are visible.

To the south, 't Zinkske connects to the former wet farmland 't Molentje and the Peel remnant Heitrakse Peel. Together, these three areas, linked by the Deurne Canal, form a coherent, extensive nature zone, almost entirely owned by Staatsbosbeheer. Southwards, this zone continues along the Helenavaart into the municipality of Peel en Maas (formerly Helden) with nature reserves such as Scherliet, Het Kwakvors, and Marisberg.

On the dry, sandy Marisberg, water infiltrates and re-emerges in species-rich wet grasslands. Between these lie the raised bog remnants Scherliet and Zinkske, where living sphagnum moss still occurs locally. This infiltration—seepage system, demonstrated by two groundwater gauges, illustrates on a small scale how the vast Peel bog developed. The moorland is home to many water and marsh birds, observable from a bird hide at the Blankerplas. Surrounding peat meadows host waders such as curlew, lapwing, and black-tailed godwit. Historical traces remain, such as two old peat canals (with many beaver gnaw marks).

On the north-east side of Meijel lies the restored nature reserve 't Molentje, which became farmland after peat cutting. For restoration, the topsoil was removed. It lies above the Peel Boundary Fault, and

) TNO Intern 183/219

buffered groundwater supports wijst indicator species such as royal fern, six-stamened waterwort, tuffed loosestrife, bristle club-rush, marsh lousewort, and heath lousewort. The Blankerplas attracts ducks, waders (when mudflats are exposed), and dragonflies, thanks to seepage-fed waters.

Industrial & Economic:

4. Relation with other geosites

Deurnse- en Mariapeel (GS_11)


5. Documentation & Resources

6. Maps, photos and figures



Location of GS_34 with the Zinkske, Marisberg, and 't Molentje

) TNO Intern 184/219



) TNO Intern 185/219

Geosite: GS_35	Name
Significance: National	De Hooge Raam
Coordinates (RD): 176780, 414199	
Municipality	Land van Cuijk - Maashorst
Province and country	Noord-Brabant
Protection status	Natuurnetwerk Nederland (NNN), Natte Natuurparel
Site management organizations	Waterschap Aa en Maas
AccessibilityH	The area is located near Hoogeweg, Zandvoortsestraat, and Busweg, with several access points and walking opportunities.
Relevance for tourism	Recreation is an important component of the area plan. Space is explicitly created for walking, cycling, and nature experience, but mass tourism is not desired. The emphasis is on small-scale, sustainable recreation.
Relevance for education & information	
Teaching equipment	

1. Location

The Hooge Raam nature reserve lies in the village of Zeeland on the border with Langenboom. It consists of a wijst reserve, a restored section of the Hooge Raam stream, and the centuries-old Smitsbos woodland. The brook, the Hooge Raam, rises in a wijst area located on the Peel Horst, flows through the Smitsbos, and joins the Raam (or Graafsche Raam) just south of Escharen. The Raam flows into the River Meuse at Grave.

2. Landscape formation and description

Origin of the landscape

The brook "Hooge Raam" originates in a wijst reserve situated above one of the southern faults of the Tegelen Fault Zone. The area between this southern branch and the northern branch near Mill forms an elevated horst structure. The wijst area lies on this horst, and the seepage water drains northwards via the Hooge Raam brook.

Geomorphology:

A total of 150 hectares of new nature will be created, including woodland and restoration of historical elevation differences. Implementation of the redevelopment is expected to start in 2026/2027.

Hydrology:

The excavated banks of the Hooge Raam form a steep edge, fed by iron-rich seepage water. The area has been designated as a Natte Natuurparel (Wet Nature Pearl) and will feature a natural stream course, providing space for water, nature, and recreation.

Deposits & Fossils:

) TNO Intern 186/219

River & Streams:

The wijst reserve was established in 2005 and forms part of the valley of the old Hooge Raam. Downstream, land along the brook has been purchased and the bank excavated. In one section, the meander has been restored.

3. Geosite Interests

Cultural History

The planned redevelopment will make the area more accessible and enjoyable, with attention to cultural history and landscape quality.

Flora & Fauna:

The Hooge Raam flows through the wijst reserve in a historic channel dating from before land consolidation. During restoration, the old seed bank was exposed, allowing the original vegetation to return in part. In recent years, a distinctive marsh orchid has appeared, considered an ecotype and variety of the Hooge Raam. Iron-rich seepage (wijst) binds phosphates, enabling the original flora to recover. A small population of heath spotted orchids is an example of this.

The Smitsbos is characterized by alder carr woodland, with transitions to bird cherry—ash woodland. Due to desiccation, common silver birch, pedunculate oak, and American black cherry have expanded significantly. The woodland is highly valuable for its rich flora, including common seepage indicators such as garlic mustard and lesser celandine, as well as rarities like Herb Paris, Moschatel, Lords and Ladies, and opposite-leaved golden saxifrage ("Goudveil-Essenbos"). This is the only site of this rare plant community in Brabant.

In recent years, beavers have settled in this area, precisely in the zone designated for free meandering. Beaver activity, such as dam building, influences water levels and contributes to natural dynamics. The Hooge Raam sometimes dries up in summer; however, pools behind beaver dams allow fish and macrofauna to survive. Kingfishers and reed birds such as Reed Warbler, Marsh Warbler, Reed Bunting, and Bluethroat breed here.

Industrial & Economic:

At the point where the Rusvensche Loop enters the old Hooge Raam channel via a weir lies the centuries-old Smitsbos. The Smitsbos was established by the Smits family from Reek, organ builders (organ of St James the Greater Church in Zeeland, municipality of Landerd). For organ building, wood from crack willow was used, both planted and naturally occurring here.


4. Relation with other geosites

GS_12 Wijstboerderij Zeeland borders this geosite to the south.

5. Documentation & Resources

6. Maps, photos and figures

) TNO Intern 187/219



) TNO Intern 188/219

Geosite: GS_36	Name
Significance: Regional	De Vilt
Coordinates (RD): 192281, 410678	
Municipality	Land van Cuijk
Province and country	Noord-Brabant
Protection status	Designated nature reserve with special ecological value
Site management organizations	Stichting Brabants Landschap
Accessibility	Several access gates are available.
Relevance for tourism	Car parks at De Kleine Vilt and De Grote Vilt. Various walking routes.
Relevance for education & information	Various excursions organised by IVN
Teaching equipment	
Scientific interest	Palaeogeographical reconstruction of the River Meuse, Woolderink et al., 2018.

1. Location

De Vilt is a nature reserve located between Oeffelt and Beugen in the Dutch province of Noord-Brabant. It covers an area of 154 hectares and is owned by Stichting Brabants Landschap.

2. Landscape formation and description

Origin of the landscape

During the Allerød period (10,950–11,900 years ago), the River Meuse flowed here, cutting a large bend (meander) through the landscape. During the Younger Dryas, the climate became significantly colder once more, and the meandering river changed into a braided system with a wide riverbed that cut through the old meanders (see Woolderink et al., 2018). Since the last Ice Age, the Meuse has resumed meandering and has shifted in a north-easterly direction on the tectonically tilting Venlo Block. During the Holocene, the Meuse valley experienced an above-average number of shifts, visible today as abandoned meander loops. The valleys of streams that developed mainly after the Younger Dryas also indicate a drainage system caused by tectonic tilting, originating at the Tegelen Fault and discharging into the Meuse.

Geomorphology:

De Vilt lies in the floodplains of the present-day Meuse and represents an abandoned Meuse meander from around 12,000 years ago, now recognisable as low-lying wet areas or oxbow lakes. The former river bend now forms an attractive enclosed landscape with striking elevation differences. On the high inner bend of the river, known as Hoog Werveld, lie open fields. In the channel itself, called Laag Werveld, there is a mosaic of water, wet woodland, and moist grasslands. The woodland and grasslands occupy the less deeply excavated or now silted-up parts of the channel.

) TNO Intern 189/219

Hydrology:

Deposits & Fossils:

River & Streams:

De Vilt represents an old Meuse bend that was cut off from the river between 9000 and 1000 BC. The elevated inner bend (point bar or omloopberg) is characterized by river deposits of sand and gravel. To construct the railway between Boxtel and Wesel, the point bar was almost completely excavated. De Vilt illustrates the contrast between natural cut-offs of Meuse bends and those artificially created in the 20th century, such as those north of the weir and lock complex at Sambeek. This complex can be visited, and during guided tours, lock attendants explain the operation and history of the weir and locks.

3. Geosite Interests

Cultural History

After the Meuse bend at De Vilt was cut off, a peat bog formed here, lasting until the Middle Ages when people learned to use peat as fuel. The peat was excavated, and in the deepest pits, the lakes De Vilt and De Kleine Vilt formed. Around 1850, peat extraction resumed, and by 1920 all the peat had been removed.

Flora & Fauna:

Nadat al het veen was afgegraven, begon de Vilt weer te verlanden en in de jaren 50 en 60 van de 20e eeuw waren er soorten te vinden als Roerdomp, Zwarte stern en Boomkikker. Daar de Oeffeltse Raam dwars door het gebied stroomde, en deze in die tijd sterk vervuild was en ook voedselrijk water bracht, verruigde het gebied. Toen dit riviertje was gesaneerd werd een plan tot natuurherstel uitgevoerd tussen 2007 en 2010. Huidige zeldzame plantensoorten en indicatoren van kwel in het gebied zijn Waterdrieblad, Draadzegge en Grote boterbloem. De Oranjetip is een vlinder die in moerasweitjes te vinden is. Tot de broedvogels behoren: Grauwe gans, Dodaars, Waterral en Ransuil. In 2007 werden twee ooievaarspalen geplaatst, waarvan sindsdien ieder jaar minstens één bezet is.

After peat removal, De Vilt began to silt up again. In the 1950s and 1960s, species such as Eurasian Bittern, Black Tern, and European tree frog were present. The Oeffeltse Raam stream flowed through the area, but at that time it was heavily polluted and nutrient-rich, causing the area to become overgrown. After the stream was cleaned up, a nature restoration plan was implemented between 2007 and 2010. Current rare plant species and seepage indicators include Bogbean, Slender Sedge, and Greater Spearwort. The Orange Tip butterfly is found in the marsh meadows. Breeding birds include Greylag Goose, Little Grebe, Water Rail, and Long-eared Owl. In 2007, two stork nesting poles were installed, and at least one has been occupied every year since.

Industrial & Economic:

After the closure of the Meuse bend at De Vilt, a peat bog formed here, until the Middle Ages when people learned to use peat as fuel. The peat was excavated, and in the deepest excavated depressions, the waters De Vilt and De Kleine Vilt formed. Around 1850, peat extraction began. By 1920, all the peat had been removed.

4. Relation with other geosites

) TNO Intern 190/219


Maasheggen (GS_01)

5. Documentation & Resources https://nl.wikipedia.org/wiki/De_Vilt

6. Maps, photos and figures




Location of GS_36 De Vilt indicated with a light blue outline



AHN image of GS_36 clearly showing the abandoned Meuse meander

) TNO Intern 191/219



Reconstruction of former Meuse courses (Woolderink et al., 2018): Green colors – meandering Meuse before the last Ice Age, Yellow – braided Meuse during the last Ice Age, Orange colors – meandering Meuse after the last Ice Age.

) TNO Intern 192/219

Geosite: GS_37	Name
Significance: Regional	Watermolen Oploo
Coordinates (RD): 188755, 402074	
Municipality	Land van Cuijk
Province and country	Noord Brabant
Protection status	
Site management organizations	De Hollandsche Molen
Accessibility	The watermill is open to the public.
Relevance for tourism	Several walking and cycling routes pass by the watermill.
Relevance for education & information	
Teaching equipment	

1. Location

The watermill D'n Olliemeulen stands in Oploo on the Vloet brook (also called the Oploose Molenbeek), close to the windmill De Korenbloem.

2. Landscape formation and description

Origin of the landscape

The choice of location for the watermills was partly determined by the point where the Vloet brook crosses the Tegelen Fault. Strong seepage in front of the fault ensured a relatively constant water supply, while the difference in elevation between the horst and the graben provided additional fall for water power.

Geomorphology:

Hydrology:

Wijst phenomena (iron-rich seepage zones).

Deposits & Fossils:

River & Streams:

This geosite clearly illustrates that the most suitable location for a watermill on a brook is at or just downstream of the point where the brook crosses a fault. Along the Tegelen Fault, several waterwheel mills have operated since the Middle Ages. They functioned thanks to the streams draining the Peel Horst towards the River Meuse.

3. Geosite Interests

Cultural History


) TNO Intern 193/219

The mill belonged in 1516 to the Lordship of Oploo under the Van Steenhuys family. The current mill building dates from 1725. Over the centuries, the mill has been used as an oil mill, grain mill, and peeling mill. The watermill operated as an oil mill until the early 20th century. Because the water level of the Vloet was not always sufficient, horse power was also used to keep the mill running. Later, the mill switched to grinding grain, first using water power and later with the help of a diesel engine. In 1800, a windmill—the post mill De Korenbloem—was built near the watermill to allow milling in all seasons.

Flora & Fauna:

Industrial & Economic:

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures



Location of GS_37 Oploo Watermill indicated with a light blue outline.

) TNO Intern 194/219



D'n Olliemeulen – photo courtesy of Marcel van Nies (30-08-2016)

) TNO Intern 195/219

Geosite: GS_38	Name
Significance: Regional	De Berkt
Coordinates (RD): 162763, 414244	
Municipality	Bernheze (Heesch)
Province and country	Noord-Brabant, the Netherlands
Protection status	Provincial Nature Map and Natuurnetwerk Nederland
	(NNN), National Monument (no. 26527)
Site management organizations	Private ownership
Accessibility	The estate is open to the public.
Relevance for tourism	The estate is included in the regional walking and
	cycling route networks. Accommodation options are
	planned on the estate.
Relevance for education & information	IVN organises excursions on the estate.
Teaching equipment	

1. Location

Castle De Berkt lies just north of the hamlet of Munnekens Vinkel in a semi-open cover sand landscape. The original surrounding estate of De Berkt unfortunately lost part of its small-scale landscape elements during large-scale land consolidation in the 20th century.

2. Landscape formation and description

Origin of the landscape

This geosite is located in the Roer Valley Graben, a tectonically subsiding area bounded on the east by the Peel Boundary Fault. However, immediately west of De Berkt lies another fault line, considered a subsidiary fault of the Peel Boundary Fault Zone. Because groundwater flows from east to west and the vertical fault plane is less permeable to water than the surrounding sediment, the presence of this fault creates relatively wet conditions at De Berkt and the local occurrence of wijst phenomena.

Geomorphology:

The landscape of the Roer Valley Graben is dominated by a cover sand relief intersected by shallow stream valleys, though this was degraded by large-scale land consolidation in the 20th century. Castle De Berkt stands on a cover sand ridge, which in the Middle Ages was further raised by applying a mixture of stable manure and heath sods (so-called potstal culture).

Hydrology:

Local wijst occurs: the emergence of iron-rich seepage water due to the nearby presence of a poorly permeable fault plane. This can be recognized by the blue sheen of iron oxide on the water surface.

Deposits & Fossils:

A c. 10 m thick layer of silty sand (Middle and Late Pleistocene periglacial deposits; Boxtel Formation) is generally overlain by a 1–2 m thick layer of low-silt cover sand (Late Pleistocene aeolian deposits; Boxtel Formation, Wierden Member). At De Berkt, a mixture of stable manure and heath sods was

) TNO Intern 196/219

applied on top of the cover sand, enriching and thickening the soil (known as 'esdek').

River & Streams:

Drainage in the area is provided by a network of ditches, partly following former stream courses such as the Vinckel.

3. Geosite Interests

Cultural History

Castle De Berkt originally formed part of the De Berkt estate, which emerged around 1800 through the reclamation of heathland and marshland. At that time, the landscape consisted of shallow stream valley soils situated between the Vinckel brook and the higher, already cultivated arable lands of the hamlets Munnekens Vinkel and Zoggel. Today, remnants of the old estate remain: historic buildings, a moat, an old orchard with standard fruit trees, a star-shaped woodland, monumental trees, and avenues.


Flora & Fauna:

Rare plant species such as Greater Tussock-sedge and Flowering Rush, along with rich woodland species like Male Fern, Yew, and Holly, indicate the presence of iron-rich seepage water (wijst).

Industrial & Economic:


- 4. Relation with other geosites
- 5. Documentation & Resources

6. Maps, photos and figures



Location of GS_38 De Berkt indicated with a light blue outline. The geosite lies strictly within the Roer Valley Graben, between the main Peel Boundary Fault Zone and a southern branch of it

) TNO Intern 197/219



Left: De Berkt lies on a cover sand ridge (olive green). Right: Despite degradation by land consolidation, the cover sand ridge (dashed line) is still visible in the AHN image.



Castle De Berkt (photo: Annélien van Kuilenburg)

) TNO Intern 198/219

Geosite: GS_39	Name
Significance: Regional	Het Loo
Coordinates (RD): 168512, 414416	
Municipality	Bernheze
Province and country	Noord-Brabant, The Netherlands
Protection status	Brabants Landschap and part of the Natuurnetwerk
	Nederland (NNN)
Site management organizations	Staatsbosbeheer
Accessibility	Easily accessible via walking and cycling paths
Relevance for tourism	Walking trails such Boswachterspad, several Canoe routes
Relevance for education & information	
Teaching equipment	

1. Location

Het Loo is a woodland and heathland area east of the A50 in the municipality of Bern

2. Landscape formation and description

Origin of the landscape

This geosite lies on the Peel Horst and consists of a woodland and heathland area intersected by the south-west to north-east trending Peel Boundary Fault. Over the barely visible fault lie several broad, shallow stream valleys that locally give rise to wijst phenomena. These valleys formed during the last Ice Age, when rain and meltwater could not infiltrate due to permafrost, instead flowing over the surface and carrying away part of the sandy sediment. The heathland is a cultural landscape created by overgrazing and soil erosion in the Middle Ages. Wind-blown sand from exposed soils formed elongated drifting sand ridges.

Geomorphology:

The cover sand and drifting sand relief are still relatively intact, though poorly visible in the field due to forest cover. The broad, shallow stream valleys on the south-western side have largely been parcelled out, disturbing the original morphology.

Hydrology:

Due to large-scale lowering of the groundwater table, seepage water (wijst) now only appears locally at the surface. Currently, watercourses are being made shallower and weirs installed to retain more water in the area.

Deposits & Fossils:

Coarse, gravelly river sand deposited by the Rhine and Meuse (Early and Middle Pleistocene; Waalre and Beegden Formations) is locally overlain by a 1 m thick layer of silty cover sand deposited by wind at the end of the last Ice Age (Late Pleistocene; Boxtel Formation). On top of this lie several elongated

) TNO Intern 199/219

drifting sand dunes (Holocene; Boxtel Formation, Kootwijk Member), formed by human activity.

River & Streams:

The narrow, canalised brook south of the hamlet of Het Loo lies in a broad, shallow periglacial valley and is therefore an example of an "underfit river".

3. Geosite Interests

Cultural History

Nearby are the cultural-historical sites Paalgraven and Vorstengraf, though these are not part of this geosite.

Flora & Fauna:

Grazing European bison keep the heathland open and varied.


Industrial & Economic:

4. Relation with other geosites

This area has a similar setting to Slabroek (GS_13), though the landscape around Slabroek is somewhat better preserved.


5. Documentation & Resources

6. Maps, photos and figures



Location of GS_39 Het Loo indicated with a light blue outline

) TNO Intern 200/219



AHN image of GS_39 clearly showing higher ground on the eastern side of the Peel Boundary Fault and drainage patterns

) TNO Intern 201/219

Geosite: GS_40	Name
Significance: Regional	Waterbloem
Coordinates (RD): 192246, 367671	
Municipality	Leudal
Province and country	Limburg
Protection status	Natte Natuurparel
Site management organizations	Staatsbosbeheer
Accessibility	Via Heibloem
Relevance for tourism	
Relevance for education & information	
Teaching equipment	

1. Location

On the border between Meijel and Heibloem in the municipality of Leudal lies the Waterbloem nature reserve (450 ha), intersected by the Roggelsche Beek. The Peel Boundary Fault runs east of Waterbloem but shows multiple branches here. The Waterbloem nature reserve is a former poorly drained depression enclosed by the Peel Horst, up to 3 metres higher on the north side, and cover sand ridges on the south side. Until around 1850, this area was an extensive marshland, but today it consists of a highly varied woodland landscape with patches of heath, farmland, and still some wetlands.

2. Landscape formation and description

Origin of the landscape

The Waterbloem area clearly illustrates the complexity of a fault zone in relation to the deposition of cover sand during the last Ice Age. Here, the Peel Boundary Fault branches into several faults, creating multiple horsts and grabens. The impermeability of individual faults explains the occurrence of numerous wijst phenomena. It is an extensive area with wet and dry forests, occasional moist grasslands, and several artificial streams.

Geomorphology:

On the road from Panningen to Heibloem, the elevation drops by 1.5 m over a distance of 75 m at the Peel Boundary Fault. In other places, the difference can reach up to 3 m. The high arable field on the west side of the road borders a lower-lying wet meadow.

Hydrology:

Here, the hydrological situation appears reversed: the wet areas lie below the Peel Boundary Fault, and the dry areas above it. The near absence of wijst phenomena with iron-rich seepage is also atypical. This is explained by the presence of large amounts of cover sand deposited over the Peel Boundary Fault, which almost eliminated the fault-related elevation difference. The loam and loamy

) TNO Intern 202/219

sand at the base of the cover sand prevent water from draining away easily. The construction of the drainage canal on the Peel Horst also ensured that seepage water was quickly removed, greatly reducing the supply of wijst water towards the Roer Valley Graben and thus limiting wijst phenomena. The presence of alder carr and birch carr woodland indicates a subsidiary fault (branch) of the Peel Boundary Fault on the western side of Waterbloem: a seepage zone and rainwater lens.

Deposits & Fossils:

Cover Sand ridges

River & Streams:

The Roggelse Beek lies in an old Meuse channel, which the brook has incised over the centuries to keep pace with the deepening Meuse. This has created a significant elevation difference in the landscape. The original source of the Roggelse Beek lies in the wetland nature reserve Waterbloem, north of Heibloem. The brook flows through the Leudal nature reserve (where it is called the Zelsterbeek) towards the Meuse. The Roggelse Beek was straightened in the 1960s–70s. Some sections, such as here, have been restored and now meander again as before.

3. Geosite Interests

Cultural History

From 1850 onwards, the marshy Waterbloem was reclaimed and managed as an estate by the Haffmans notary family, who also used it as a hunting ground. Staatsbosbeheer later planted production forests here.

That the higher side of the Peel Boundary Fault must once have been much wetter (before the drainage canal was built) is evident from the elevated position of roads and names such as Roggelse Dijk, which formed part of the trade route between Cologne and 's-Hertogenbosch. This road dates back at least to the 18th century and was well maintained to remain passable year-round.

Flora & Fauna:

These wijst areas in Waterbloem are marked by alder carr woodland. In addition to the well-known wijst indicator Water Violet (probably the origin of the name Waterbloem), the relatively rare Floating Water-Plantain, which is protected under European law, also occurs here. The presence of Marsh Violet, Lousewort, Heath Milkwort, and Tormentil indicates a rainwater lens. Heath Milkwort winged pea is typical of transitions from heath to marsh vegetation such as bluegrass meadows. Rare butterflies such as the Chequered Skipper and Large Chequered Sskipper still inhabit Waterbloem. The area is rich in birdlife, and the chances of encountering roe deer are high. Rooting traces reveal the presence of wild boar.

With rewetting measures, the water authority hopes to restore characteristic plant communities. The works aim to connect the nature reserve to the remaining peat bogs of the Peel and create an ecological corridor between them. Agricultural drainage water will be diverted so that water remains in the area for longer.


Industrial & Economic:

4. Relation with other geosites

The geological setting is comparable to geosite GS_28 Sint Willibrordusput.

5. Documentation & Resources

) TNO Intern 203/219



) TNO Intern 204/219

Geosite: GS_41	Name
Significance: Regional	De Krim
Coordinates (RD): 182945, 398302	
Municipality	Gemert-Bakel (Elsendorp)
Province and country	Noord Brabant
Protection status	Natuurnetwerk Nederland (NNN)
Site management organizations	Brabants Landschap
Accessibility	
Relevance for tourism	Several cycling and walking routes pass through the area
Relevance for education & information	Informatiepanel at De Krim "Niet alles ontgonnen, heel slim" ("Not everything reclaimed – very wise")
Teaching equipment	

1. Location

De Krim is an estate in the Peel region near Elsendorp. The estate covers approximately 100 hectares and is owned by Brabants Landschap. De Krim is surrounded by other small nature reserves managed by Brabants Landschap and connects to the estates Groote Sink and Bunthorst. De Krim tells the story of the last Ice Age and contains a preserved pingo ruin. Moreover, the area has never been reclaimed.

2. Landscape formation and description

Origin of the landscape

What makes the nature of the De Krim estate unique is that it has never been reclaimed. Other parts of the Peel have been restored to their former glory (such as the Klotterpeel), but De Krim, thanks to its untouched state, still contains a pingo. The majority of the area consists of wet heathland, three ponds, and the aforementioned pingo ruin, a remnant from the last Ice Age (the Weichselian). During this last Ice Age, Brabant was not covered by ice, but it was extremely cold and dry, and the upper soil layer was permanently frozen (permafrost).

A pingo is a low-lying spot where water accumulates beneath the frozen ground. In winter, this water freezes and expands, creating a circular mound. This mound persisted until the end of the Ice Age, when the accumulated ice and permafrost melted, leaving a circular depression in the ground. These depressions often became filled with peat. During various land consolidations and Peel reclamations, pingos were often completely levelled. The pingo ruin in De Krim has a diameter of about 90 metres.

Geomorphology:

The pingo ruin in De Krim has a diameter of approximately 90 metres

Hydrology:

Deposits & Fossils:

) TNO Intern 205/219

River & Streams:

3. Geosite Interests

Cultural History


The area had various owners until 1920, when it was first developed as a small-scale farm. There were three wooden cottages along the Middenpeelweg, of which one farm remains, but the vast majority of the land was never cultivated.

Flora & Fauna:

De Krim is one of the few remaining heathland relics, left untouched by landowners for hunting purposes. The idea was that a varied landscape would provide a diverse supply of game. De Krim is a habitat for the viviparous lizard, nightjar, and stonechat. Two artificial ponds host the great crested newt, a species characteristic of highly acidic waters.


Industrial & Economic:

- 4. Relation with other geosites
- 5. Documentation & Resources
- 6. Maps, photos and figures



Location of GS_41 De Krim indicated with a light blue outline

) TNO Intern 206/219



) TNO Intern 207/219



) TNO Intern 208/219

Geosite: GS_42	Name
Significance: Regional	De Elsbeemden
Coordinates (RD):	
Municipality	Horst aan de Maas
Province and country	Limburg
Durate etien etetue	
Protection status	
Site management organizations	Staatbosbeheer
Accessibility	Access from the Maasbreeseweg.
Relevance for tourism	Several walking routes run through the area.
Relevance for tourism	Several walking routes run through the area.
Relevance for education & information	
Teaching equipment	The area is highly suitable for field biology and nature
0 - 4 - 1	education.
Description	

_ -----

1. Location

De Elsbeemden (beemd means wet hay meadow) is a nature reserve south of Sevenum. The area covers 65 hectares and forms part of Staatsbosbeheer's Molenbeekdal project. Two streams flow through the area: from the west comes the Groote Molenbeek, which was restored a few years ago to a meandering, shallower, but wider course; from the south comes the Elsbeek, which joins the Groote Molenbeek.

2. Landscape formation and description

Origin of the landscape

On the fault map of North Limburg, several faults are shown in addition to the Peel Boundary Fault. On the GeoparkPlus map, only the Peel Boundary Fault is indicated. However, in De Elsbeemden there are clear wijst indicators, which can be attributed to the Sevenum Fault. This fault impounds water in De Elsbeemden, creating a large area of wet grasslands or beemden. The Groote Molenbeek carries the water across the fault eastwards to the River Meuse. The beemden are nutrient-poor due to ironrich seepage water and can only be used for haymaking. The area is rich in rushes.

Geomorphology:

Hydrology:

Due to interventions in the streams, the entire area has become wetter. The vegetation consists largely of carr woodland and wet grasslands.

Deposits & Fossils:

High iron concentrations were found mainly where deposits belonging to the marine Miocene lie relatively close to the surface (sometimes less than 10 m below ground level). These deposits formed between about 23 and 5 million years ago. The iron in the groundwater on the Peel Horst probably originates largely from glauconite, which occurs abundantly in marine Miocene sediments.

) TNO Intern 209/219

Glauconite contains high iron levels (ranging from about 6 to 19% Fe_2O_3 and 4 to 5% FeO). In addition, remnants of the Peel peat bogs are important. The highest iron concentrations are found where water has infiltrated through peat or highly humic soils. This acidic infiltration increases the solubility of iron.

River & Streams:

The Groote Molenbeek is the longest stream on the eastern side of the Peel Horst and rises in an intake area east of the watershed near Grashoek.

3. Geosite Interests

Cultural History:

Excavations of foundations of churches destroyed during the war in North Limburg revealed that iron ore was widely used as a foundation material for churches in this region, including the church tower of Sevenum.

Flora & Fauna:

The fault line is marked by vegetation of Black Alder, grading into downy birch and pedunculate oak. Near the fault grow characteristic wijst indicators such as Guelder Rose, Spindle, Hop, and Red Ddogwood. In the streams and ditches, iron-rich seepage increases near the fault. Aquatic plants include Needle Spike-rush, Brooklime, Various-leaved Water-starwor, and Bog Pondweed. Since 2001, the bird group 't Hökske has monitored breeding birds here. Between 2001 and 2006, 67 bird species were recorded as breeding. In addition to common species, the group recorded an average of six pairs of Nightingales per year, as well as Stonechats, Kingfishers, Bbluethroats, Little Grebes, Water Rails, Reed Warblers, Marsh Warblers, and also Grasshopper Warblers, Yellowhammers, and Reed Buntings.

Industrial & Economic:

Iron ore was mined in the Elsbeemden near Sevenum.

4. Relation with other geosites

5. Documentation & Resources

- Ettema, N. (2021) Ecologische iconen van de Wijst Wat breuken ons bovengronds laten zien. Geopark Peelhorst en Maasvallei i. o., Uden, 121 pp.
- Munckhof, P.J.J. van den (2000), 'Glauconiethoudende afzettingen in de Peelregio. Een ijzersterke basis voor behoud en ontwikkeling van voedselarme, natte milieus!' In: Natuurhistorisch maandblad jaargang 89, 43-52.

6. Maps, photos and figures

) TNO Intern 210/219



Location of GS_42 De Elsbeemden indicated with a light blue outline



Elsbeemden with the Groote Molenbeek crossing the Sevenum Fault (from Ettema, 2021). See topographic map for location

) TNO Intern 211/219

) TNO Intern) TNO 2025 R11935

) TNO Intern 212/219

Appendix B

Bibliography

This appendix contains the Bibliography compiled by the Scientific Board of the Geopark Peelhorst & Maasvallei, supplemented with the key references of this report.

- A Aequator Groen & Ruimte en Witteveen+Bos (2009), Uitvoeringplannen Brabantse wijstgronden. Gebieden Slabroek en Hengstheuvel
 - Ahorner, L. (1994) 'Fault-plane solutions and source parameters of the 1992 Roermond, the Netherlands, mainshock and its stronger aftershocks from regional seismic data', in: Geologie en Mijnbouw 73 (2-4, Special Issue), 199-214.
 - Ancker, H. van den en P. Jungerius (2014), Een Europees Geopark Noord-Brabantse Zandlandschappen: een voorverkenning
 - Arcades, Gemeente Boekel, Waterschap De Aa, Waterschap de Maaskant (2003), Waterplan Boekel Wijs(t) met water
 - Arends, G.J. (1994), Sluizen en stuwen (bouwtechniek in Nederland 5) [Delft/Zeist]
- B Beijers, H., en G.J. van Brussel (1996), Van d'n Aabeemd tot de Zwijnsput. Toponiemen in de cijnskring Helmond voor 1500 in naamkundig en nederzettingshistorisch perspectief [Helmond]
 - Bekker, J. de, M. Deij, M. Steinmann, L. Strackx en M. van der Wielen (2022), Volg het verborgen landschap Toekomstvisie voor het Geopark Peelhorst en Maasvallei in 2050 Onderzoek HAS 's- Hertogenbosch iov Waterschap Aa en Maas en provincie Noord-Brabant Bense, V.F. (2002), 'Hydrogeologische karakterisering van breukzones in Zuidoost-Nederland', in: Stromingen 8, 17-30
 - Bense, V.F. en R.T. van Balen (2003), Hydrogeological aspects of fault zones on various scales in the Roer Valley Rift System' in: Journal of Geochemical Exploration 78, 317-320 Bense, V.F., R.T. van Balen en J.J. de Vries (2003), The impact of faults on the hydrogeological conditions in the Roer Valley Rift System: an overview'. In: Neth J Geosci 82:41–53
 - Bense, V.F., E.H. van den Berg, R.T. van Balen (2003), Deformation mechanisms and hydraulic properties of fault zones in unconsolidated sediments; the Roer Valley Rift System, The Netherlands' in: Hydrogeology Journal 11, 319-332
 - Bense, V.F. en R.T. van Balen (2004), The effect of fault relay and clay smearing on groundwater flow patterns in the Lower Rhine Embayment' in: Basin Research 16 (3), 397-411
 - Bense, V.F. en H. Kooi (2004), Temporal and spatial variations of shallow subsurface temperature as a record of lateral variations in groundwater flow' in: Journal of geophysical research: Solid Earth 109 (B4)
 - Bense, V.F., M.A. Person, K. Chaudhary, Y. You, N. Cremer en S. Simon (2008), Thermal anomalies indicate preferential flow along faults in unconsolidated sedimentary aquifers' in: Geophysical Research Letters 35 (24)
 - Bense, V.F., T. Gleeson, S.E. Loveless, O. Bour, J. Scibek (2013), 'Fault zone hydrogeology', in: Earth- Science Reviews 127, 171-192
 - Bense, V.T., T. Read en A. Verhoef (2016), Using distributed temperature sensing to monitor field scale dynamics of ground surface temperature and related substrate heat flux' in: Agricultural and Forest Meteorology 220, 207-215
 - Berendsen, H.J.A., 2005. Landschap in Delen. Van Gorcum, Assen, the Netherlands, 340 pp.
 - Berg, M. van den, K. Vanneste, B. Dost, A. Lokhorst, M. van Eijk and K. Verbeeck (2002), 'Paleoseismic investigations along the Peel Boundary Fault: geological setting, site selection and trenching results', in: Netherlands Journal of Geosciences; Geologie en Mijnbouw 81 Berkom, R. van (1987), Veldnamen. Voorlopige lijst van toponymen in Nistelrode en Vorstenbosch van 1420-1455 en 1800-1810 [Nistelrode]

) TNO Intern 213/219

Berz, G. (1994), 'Assessment of the losses caused by the 1992 Roermond earthquake, the Netherlands', in: Geologie en Mijnbouw 73, 281

Bestuurlijk Platform Reconstructie (2000), Pilot Gemert-Bakel. Deel 1 reconstructieplan en Deel 2, uitvoeringsplan [Gemert-Bakel]

Beusekom, E. van (2007), Bewogen Aarde. Aardkundig erfgoed in Nederland [Utrecht]

Bon, J. (1972), 'Hydrologische veldkenmerken langs de Westflank van de Peelhorst', in: Tijdschrift van het Koninklijk Nederlands Aardrijkskundig Genootschap p65

Bont, Chr. de (1993), Al het merkwaardige in bonte afwisseling. Een historische geografie van Midden en Oost-Brabant [Waalre]

Bonte, M., en Th.G.J. Witjes (2007), Nadere Inventarisatie Wijstgebieden, Eindrapport Witteveen+Bos

Brand, M.P.J. van den (1982), Lief en Leed in en over De Oude Peel [Venray]

Brand, M.P.J. van den, en I.J.M. Meuwissen (2003), Brabantse Wijstgronden in Beeld. Waterschap de Aa

Buijks, H. (1996), Van den Grave aff totter Diezen toe. Bijna 700 jaar waterschappen in het Maasland 1307-1996 [Oss]

Buiter, H. en R. Raat (2016), Nederland Kanalenland. Een reis langs twaalf kanalen [Amsterdam]

Buro Hemmen en Overlegorgaan Nationaal Park De Groote Peel (2002), Water en Vuur. Beheers- en Inrichtingsplan Nationaal Park De Groote Peel, [Maastricht]

Busschers, F.S., Cohen, K., Wesselingh, F.P., Schokker, J., Bakker, M., Van Balen, R. & Van Heteren, S., 2025. Quaternary. In: J.H. Ten Veen, G.-J. Vis, J. De Jager, J. & Th.E. Wong (eds): Geology of the Netherlands, second edition. Amsterdam University Press (Amsterdam): 333-391. DOI: 10.5117/9789463728362 ch10

- C Camelbeeck, T., T. van Eck, R. Pelzing, L. Ahorner, J. Loohuis, H.W. Haak, P. Hoang-Trang en D. Hollnack (1994), The 1992 Roermond earthquake, the Netherlands, and its aftershocks." Geologie en Mijnbouw, 73: 181-1 97
 - Childs, C., Manzocchi, T., Walsh, J. J., Bonson, C. G., Nicol, A. & Schopfer, M. P. J. (2009). A geometric model of fault zone and fault rock thickness variations. Journal of Structural Geology, 31, 117–127
 - Cyrillus, P., en P. Canisius (1920), Vrouw van Handel, Hare geschiedenis bij de viering van het zevende eeuwfeest 1220-1920 [Helmond]
- D De Bes Van Nuland & Partners (1998), Toeristisch-Recreatief Zoneringsplan Peelvenen [Beekbergen]
 - Dienst Landelijk Gebied (2012), Peelbeleving in de Peelvenen: Visie op Recreatie en Toerisme anno 2012
 - DLG/BC Peelvenen (2010), Uitvoeringsplan Koningshoeven Cultuur [Roermond]
 - DLG/BC Peelvenen (2011), Uitvoeringsplan Koningshoeven Natuur [Roermond]
 - DLG/Landinrichtingscommissie Peelvenen (2005), Landinrichtingsplan Herinrichting Peelvenen, onderdeel Deurnsche Peel – Mariapeel (Het onverenigbare verenigd) [Roermond] DLG/Ministerie van LNV/provincie Noord-Brabant/provincie Limburg (2010), Concept-ontwerp Beheerplan Natura 2000 [Tilburg]
 - DLG/Streekcommissie Peelvenen (2000), Projectnota voor het gebied Peelvenen [Roermond]
- E Eck, T. van, en C.A. Davenport (1994), "Seismotectonics and seismic hazard in the Roer Valley graben; with emphasis on the Roermond earthquake of April 13, 1992", Geol. Mijnb. 73, 91-92
 - Enk, H. van (2016), Iedere generatie zijn eigen oerbank? Grondboor & Hamer 70: 48-55

Essink, H.B.M. (1969), Het Land van Cuyk tussen 20 April 1308 en 31 December 1839. Een bijdrage tot de landbouwgeschiedenis [Grave]

Ettema, N.A. (1980), De flora in het Natuurpark "De Maashorst". Recreatieschap Maasland Oss

Ettema, N.A. (1992), Verslag inventarisatie Beekdal van Slabroek en Grote wetering. IVN Uden

Ettema, N.A. (2001), Het begrazingsgebied in de Maashorst. IVN Uden.

Ettema, N. (2008), Mededelingen & Verspreidingsonderzoek. Meetnet Amfibieën. Nijmegen: RAVON Werkgroep Monitoring

Ettema, N.A. (2008), Verslag inventarisatie Beekdal van Slabroek. IVN Uden.

Ettema, N. (2010), Vijf Wijstreservaten in Noord-Brabant. Stuurgroep De Maashorst: 63 pp.

) TNO Intern 214/219

Ettema N. (2012), Amfibieën van De Maashorst. Natuur- en Milieuverenigingen De Maashorst Uden 87

Ettema N. (2012), Libellen van De Maashorst. Natuur- en Milieuverenigingen De Maashorst Uden

Ettema, N., en J. van der Wijst (2012), Stand van de Natuur in De Maashorst. Natuur- en Milieuverenigingen De Maashorst Uden

Ettema N. (2014), Poelenonderzoek in de gemeente Uden. Stg. Vrijwillig Landschapsbeheer Uden.

Ettema, N., en I. van der Laan (2015), Aquatisch-ecologisch poelenonderzoek in de gemeente Uden. Stg. Vrijwillig Landschapsbeheer Uden

Ettema, N. (2018), Een bijzonderlijk stukje natuur in Herperduin

Ettema, N. (2018), Wat de Maashorst uniek maakt! Een tussentijdse evaluatie van inrichting en beheer 2010-2018

Ettema, N., Flora van de Maashorst. Natuur- en milieuverenigingen De Maashorst

Ettema, N., en I. van der Laan (2019), Vennen, poelen en beken van de Maashorst. In opdracht van Sta. Natuurorganisaties De Maashorst

Ettema, N. (2022), Breuken in het land van Peel en Maas: veldgids' Geopark Peelrandbreuk en Maashorst i.o. Uitgeverij Matrijs (Utrecht), 152 pp.

G Galle, H. (2017), "Beugen en Rijkevoort en de geschiedenis van ijzer in Nederland (1858-1899)", Merlet, nr 1

Gemeente Gemert-Bakel (1998), Het is Best Buiten, Bestemmingsplan buitengebied, Gemeente Gemert-Bakel 1998-2008 Geol. Mijnb. 82, 41e54

Giesen, L. (2005), 'Hoeve de Boeshei te Swalmen-Boukoul', In: Jaarboek Maas- en Swalmdal 25, 65-73. (geschiedenis omgrachtte hoeve op/aan breuk)

Groot, E. de, en W. Peters (1993 en 1994), Wie het kleine niet eert, ...! Deel A en B. IVN Uden

Groot, E. de, W. Peters en N. Ettema (2003), Poelen en vennen, meer en beter!!! IVN Uden

Groot, P. de, Bijdrage tot de geschiedenis van Uden, deel 20, Sprokkelingen jrg 28, nr. 111

H.D. (1967), Het winnen van ijzeroer onder de gemeenten Wanroij en Mill", Merlet, nr 3

Haartsen, Adriaan (2010), Cultuurhistorische analyse van het dal van de Graafsche Raam van St.-Hubert tot Grave. Haaften, Lantschapsstudies 103

Haartsen, A. (2014), Cultuurhistorische en landschappelijke analyse van het Peelkanaal bij Deurneseweg 28 in Oploo. Haaften, Lantschapsstudies 153

Habraken, H. (2019), Uniekheid vegetatie in wijstgebieden: eindverslag

HAS Den Bosch, (2019), Diverse onderzoeksrapporten", in: Nico Ettema & Iris van der Laan. Poelen, vennen en beken van de Maashorst: (Macro)fauna, Flora, en Waterkwaliteit [Den Bosch]

Hoogma, D. (1981), Onderzoek flora en vegetatie Wijstgronden

Houtgast, R.F. & Van Balen, Ronald & Kasse, Cornelis (Kees) & Vandenberghe, J.. (2003). Late Quaternary tectonic evolution and postseismic near surface fault displacement along the Geleen Faut (Feldbis Fault Zone - Roer Valley Rift System, the Netherlands), based on trenching. Geologie en Mijnbouw/Netherlands Journal of Geosciences. 82. 10.1017/S0016774600020734.

- I Ickenroth, J. (1989), 'De Molens van Swalmen'. In: Jaarboek Maas- en Swalmdal 9, 30-41.
 - IVN Uden (1993), Help, de wijstgronden verruigen [Uden]
 - Iven, W. and van Gerwen, Th. (1978) Mensen in de Peel, Distel, 78 pp.
 - Iwaco (1994): Toelichting op de hydrologische systeemkaart van Noord- en Midden-Limburg (1994) ['s- Hertogenbosch]
- Jansen, ing. P.C. (1983), Projectgroep Zuidelijk Peelgebied 28, De gevolgen van veranderingen in de waterhuishouding voor de vegetatie. Instituut voor Cultuurtechniek en Waterhuishouding Wageningen, Nota 1476 [Wageningen]

 Jansen, R., en K. van der Laan (2011), Verleden van een bewogen landschap, Landschaps- en bewoningsgeschiedenis van de Maashorst [Utrecht]
- Koomen, A. J. M., Kiden, P. & Verbauwen, E. (eds.) (2007). Van beekdal tot stuifduin Aardkundige waarden in Noord-Brabant. Provincie Noord-Brabant ('s Hertogenbosch): 120 pp.
 - Krabbenburg, A.J., I.N.B. Poelman en E.J. van Zuilen (1983), Standaard vochtkarakteristieken van zandgronden en veenkoloniale gronden. Rapport nr 1680, Stiboka [Wageningen]

) TNO Intern 215/219

- Kuijpers, H. (2006), 'Dorpsgraaf en Bergerpoort. De wallen en poorten van Neer terug op de kaart'. In: Oos Naer 29, 4-20. (overzicht landweren rond Neer)
- L Laan, I. van der, en N. Ettema (2016), Monitoring Maashorst Groote Wetering & Venloop, Stg. Natuurorganisaties de Maashorst
 - Laban, C. (2006), "Aardkundig excursiepunt 4: Breuken in het Limburgse en Brabantse landschap." Grondboor & Hamer nr.4
 - Lans, H. E. en P.G. van der Vos (2009), Natuurplan De Maashorst. Integraal Inrichtings- en Natuurbeheerplan Maashorst-Heperduin. Rhee: Ecoplan Natuurontwikkeling
 - Lantschap (2017), Cultuurhistorische analyse van tien stuwen in het Peel- of Defensiekanaal en een stuw in de St. Anthonisloop bij Mill
 - Lapperre, R. E. (2025). Fault-Zone Hydrogeology in Unconsolidated Sediments: Cross-Fault Groundwater Levels, Fault Sealing, and Permeability Distribution in the Roer Valley Rift System. [PhD-Thesis Research and graduation internal, Vrije Universiteit Amsterdam]. https://doi.org/10.5463/thesis.1223
 - Lapperre, R.E., V.F. Bense, C. Kasse, R.T. van Balen (2022), Temporal and spatial variability of cross-fault groundwater level differences: the impact of fault-induced permeability reduction, precipitation and evapotranspiration' in: Hydrogeology Journal.
 - Lapperre R.E., C. Kasse, V.F. Bense, H.A.G. Woolderink, en R.T. van Balen (2019), "An overview of fault zone permeabilities and groundwater level steps in the Roer Valley Rift System." Netherlands Journal of Geosciences, Volume 98, e5
 - Lapperre, R., R.T. van Balen en C. Kasse (2020), Geohydrologisch onderzoek Peelrandbreuk Uden.' in: Grondboor en Hamer 74, 191-196.
 - Lokker, C. (1953), "De morfologie van de dagzoom der Peelrandbreuk", in Tijdschrift
 - Lokker, C. (1953), "De morfologie van de dagzoom der Peelrandbreuk", in Tijdschrift Koninklijk Aardrijkskundig Genootschap. Tweede Reeks, Deel LXX., pag. 331-343
 - Luijendijk, E., R. T. van Balen, M. ter Voorde, en P. A. M. Andriessen (2011), Reconstructing the Late Cretaceous inversion of the Roer Valley Graben (southern Netherlands) using a new model that integrates burial and provenance history with fission track thermochronology' in: J. Geophys. Res., 116, B06402
- M Maes, B., R. van Loon (2008), Oude boskernen in Midden- en Oost-Brabant. Brabantse Milieufederatie
 - Mattheeuwse, H. (2017), Rapport Nationaal Park de Meinweg: Een verrassend stukje Limburg.
 - Melisie, E., B. Nyssen, A. Dielissen, R. Christiaans, L. Linnartz, L. Adolfse, N. Ettema, K. van der Laan en J. Heynekamp. (2015), Inrichtings- en Beheerplan De Maashorst (2015-2019). Gemeenten Uden, Bernheze, Oss en Landerd, waterschap Aa en Maas en Staatsbosbeheer. 26p.
 - Michels, J. (1991), "De Peel-Raamstelling (1934-1940) in Noord-Brabant en Limburg. Oorlogsbuit voor Monumentenzorg?" Brabants Heem jrg 43 pp 41-55
 - Michon, L., R.T. van Balen, O. Merle en H. Pagnier (2003), Cenozoic evolution of the Roer Valley Rift System integrated at European scale'. In: Tectonophysics, 367, 101-126
 - Michon, L. en R.T. van Balen (2005), Characterization and quantification of active faulting in the Roer valley rift system based on high precision digital elevation models'. In: Quaternary Science Reviews, 24, 457-474
 - Middelkoop, T. van, (2004), "Stellingen en linies in het strategische plan van generaal Winkelman, 1940." In: Op Weerstand Gebouwd. Verdedigingslinies als militair erfgoed, Jaarboek Monumentenzorg [Zwolle/Zeist]
 - Mooren, B., (1980), "Aardschokken in Midden-Limburg." In: Rondom het Leudal 18.
 - Mourik, J.M. van, (1987), "Laat glaciaal veen en holoceen stuifzand bij de Peelbreuk tussen Nistelrode en Uden", Geografisch Tijdschrift XXI, 421-436.
 - Mourik, J.M. van, A.C. Seijmonsbergen, R.T. Slotboom en J. Wallinga (2011), "The impact of human land use on soils and landforms in cultural landscapes on aeolian sandy substrates (Maashorst, SE Netherlands)", Quaternary International
 - Munckhof, P.J.J. van den & Joosten, J.H.J. (1990): Een breuk net een verleden. Beheers- en ontwikkelingsvisie voor de Grote Eenheid Natuurgebied zuidelijke Peelhorst.- 181 s., 58 Abb.; Naluur Milieu en Faunabeheer Tilburg/Roernond.
 - Munckhof, P.J.J. van den (2000), 'Glauconiethoudende afzettingen in de Peelregio. Een ijzersterke basis voor behoud en ontwikkeling van voedselarme, natte milieus!' In: Natuurhistorisch maandblad jaargang 89, pag. 43-52
- N Natuur parken Limburg. (z.j.). (Geraadpleegd op 21 oktober 2017), Nationaal Park de Meinweg

) TNO Intern 216/219

- O Offermans, R.J.E., Bureaustudie DLG naar de historie van de Peel-Raamstelling. [Roermond] Otten, A., T. de Jong en S. van Wetten, (1998), "Watermolen van Gemert opgespoord", Gemerts Heem, nr 4.
- P Paulissen, M.P.C.P., W. Nieuwenhuizen, F.H. Kistenkas (2014), Geoparken in Nederland: Een quickscan van beleidsmatige mogelijkheden en beperkingen. Alterra Wageningen UR rapport 2537

Peng, F., M.A. Prins, C. Kasse, K.M. Cohen, N. Van der Putten, J. Van der Lubbe, W.H.J. Toonen, R.T. van Balen, An improved method for paleoflood reconstruction and flooding phase identification, applied to the Meuse River in the Netherlands.' in: Global and Planetary Change 177

Peng, F., C. Kasse, M.A. Prins, R. Ellenkamp, M.Y. Krasnoperov. en R.T. van Balen (2020), Paleoflooding reconstruction from Holocene levee deposits in the Lower Meuse valley, the Netherlands in: Geomorphology, 352.

Pennings, M.H.J., (1970), "Ook in Boekel "trekt' de Wijst", Gemerts Heem nr 38

Peters, A., Lammers, Th.& Mol, D. (1991). Mastodonten=kiezen uit Liessel (Noord-Brabant). Cranium 8 (2), 89-96.

Peters, N., (2009), Brabant tussen walvissen en mastodonten, Fossielen uit Liessel [Asten/Boxtel]

Peters, N., (2013), Van reuzenhaai tot Chalicotherium, Fossielen uit Mill-Langenboom

Pomper, A. B., (1981), Tectoniek en tectonische bewegingen in het zuidelijk gebied. Nota 1278, Instituut voor Cultuurtechniek en Waterhuishouding Wageningen.

Pomper, drs. A.B., Projectgroep Zuidelijk Peelgebied 11 (1983), Hydrochemisch onderzoek in het zuidelijk peelgebied I: Een beschrijving van de eerste resultaten van het meetprogramma in het voorjaar van 1982. Nota 1368.

Pouls, J., H. Willems en H. Crompvoets (2009), Meijel, bijzonder dorp in de Peel. O.a. Hoofdstukken 3; De gouden helm van de Peel en 4; Sint Wilbertsput en Luttel Meijel. Provincie Limburg, (2009) Landschapskader Noord- en Midden-Limburg, 68 pp.

Provincie Limburg (2021) Omgevingsvisie Limburg, 204 pp.

Provincie Noord-Brabant (2004) Aardkundig Waardevolle Gebiedenkaart Noord-Brabant, 173 pp.

- R Rooden, M. van, (2014), De Peel-Raamstelling in stelling gebracht, rapport in opdracht van DLG Roermond en aangevuld door Peelnetwerk
 - Roth, J., K.T. Jellema, P. Kloosterman, L. Wiersma en M. Schimmel (2011), Onderzoek naar de pH- waarden van het oppervlaktewater en mogelijk te verbinden gezondheidsclaims in De Maashorst.
- S Sitter, L. U. de, (1944-1945), De Keulsche bocht en de horsten van de Peel en van Geldern-Crefeld. Gedenkboek Tesch. (Verh. Geol. Mijnb. Gen., Geol. Serie deel XIV). Slits, H., (2010), "Gemerts ijzererts naar Pruisen", Gemerts Heem, nr 2
 - Smits, J., (2001), Brabants Peil. Monumenten voor waterbeheersing in Noord-Brabant. [Nijmegen]
 - Sneep, J., J.P.C.M. van Hoof, G.J.L. Koolhof & S.H. Poppema, (1996), Atlas van de historische vestingwerken in Nederland. Noord-Brabant [Zutphen]
 - Spamer, T. (2010), "Een karretje op den zandweg reed, Deurnese toponiemen uit de periode 721 1900," Deurnese Historische Reeks 8 [Deurne]
 - Stichting 'Sporen van de Oorlog', De betekenis van Mill in de historie van linies en stellingen

Stouthamer, E. & Berendsen, H.J.A.. (2001). Avulsion Frequency, Avulsion Duration, and Interavulsion Period of Holocene Channel Belts in the Rhine-Meuse Delta, The Netherlands. J. Sediment. Res.. 71. 10.1306/112100710589.

Stuurman, R.J. en R.H. Atari, (1997), De grondwatersituatie rond de Wijstgronden bij Uden; Rapport nr. 97-212(a), NITG-TNO Delft, 67 pag.

- Tesch, P., (1934), "Een profiel in het Noordoosten van Noord-Brabant", Geol. en Mijnb.
 - Tesch, P., (1940), "In hoeverre is de recente niveauverandering van tectonische aard", Geologie en Mijnbouw
 - Tesch, P., (1941), "Boringen tussen Swalmen en Meyel in Midden Limburg", Geologie en Mijnbouw.
 - Tesch, P., (1941), "Het landschap tussen Sittard en Roermond en de tectonische daling", Tijdschr. Kon. Ned. Aardr. Gen., 2e serie LVIII, pag. 940.
 - Tesch, P., (1943), "Uitwendig tegen inwendig", Geologie en Mijnbouw

) TNO Intern 217/219

Theelen, P., (2009), Proefboringen naar delfstoffen in het begin van de twintigste eeuw

Thelen. A. (red) (2001), Het Hooghuis te Gemert; Archeologisch en historisch onderzoek betreffende het middeleeuwse kasteel van de heren van Gemert. Hoofdstukken 2; De landschappelijke context en de vroegste bewoning (P. Kleij en J. Timmers) en 4; Ontwikkelingen in middeleeuws Gemert (J. Timmers)

Thissen, P.H.M., (1993), Heideontginning en modernisering, in het bijzonder in drie Brabantse Peelgemeenten 1850-1940 [Utrecht]

Timmers, J. (2018), Hoge Burcht in Uden: 'moorassighe en sumpige grond'

Timmers, J., (1996), "Omgrachte terreinen in de Middeleeuwen in Gemert", Het Brabants kasteel, nr 2/3

Timmers, J., (2011), Eessens put naar een reconstructie van een opvallende markering in het landschap

Timmers, J., (2016), "De oudste vermelding van Uden; Er was geen Ricolt van Uden in 1190", Sprokkelingen nr 117

Timmers, J., (2017), "Over de "Hoge Burcht" van Uden", Historisch Landschap

Timmers, J., (2017), De Velmolen en de Vloet, Sprokkelingen 2017

Timmers, J., (2017), Een watermolen op de Molenhof in Bakel en Deurne

Timmers, J., (2018), "De Hoge Burcht", Sprokkelingen, jrg 31, nr 124

Timmers, J., (2019), "Bronnen op de breuken", Gemerts Heem, nr 1

Timmers, J., (2019), De oudste ontginning van Slabroek langs een prehistorische weg", Sprokkelingen, jrg 32, nr 127

Timmers, J., (2020), Over mensen en de Peelrandbreuk." [Gemert]

TNO, Alterra, Provincie Noord-Brabant (2003), Monitoring Verdroging Noord-Brabant, bijlagerapport

Van Balen, R.T., J.M. Verweij, J.D. Van Wees, H. Simmelink, F. Van Bergen en H. Pagnier (2002), Deep subsurface temperatures in the Roer Valley Graben and the Peelhorst, the Netherlandsnew results.' in: Geologie en Mijnbouw/Netherlands Journal of Geosciences, 81, 19-26 Van Balen, R.T.(2009), 'Aardkundig excursiepunt 28. Peelrandbreuk en Maashorst', in: Grondboor & Hamer 63, nr 6, 155-160

Van Balen, R.T., M.A.J. Bakker, C. Kasse, J. Wallinga, en H.A.G. Woolderink (2019), 'A Late Glacial surface rupturing earthquake at the Peel Boundary fault zone, Roer Valley Rift System, the Netherlands', in: Quaternary Science Reviews 218, 254-266

Van Balen, R.T. (2020), De Peelrandbreuk in: Grondboor en Hamer 74, 171-173

Van Balen, R.T., M. Bakker, J. Wallinga en H.A.G. Woolderink (2020), Een fossiele aardbeving bij de Peelrandbreuk van Bakel.' in: Grondboor en Hamer 74, 183-187

Van Balen, R. T, C. Kasse, J. Wallinga en H.A.G. Woolderink (2021), Middle to Late Pleistocene faulting history of the Heerlerheide fault, Roer Valley Rift System, influenced by glacioisostasy and mining- induced displacement.' in: Quaternary Science Review, 268, 107111 Van Balen, R.T., N. Ettema, H. Kuijpers, R. Lapperre, B. Nelemans, J. Timmers, G. Verbeek en E. Weerman (2022), Breuken in het land van Peel en Maas' Geopark Peelrandbreuk en Maashorst i.o. Uitgeverij Matrijs. 176 pp.

Van den Akker, Pompstra (1997), Monitoringsvoorstel voor de wijstgronden, afstudeeronderzoek

Van den Berg, M., Vanneste, K., Dost, B., Lokhorst, A., Van Eijk, M., Verbeeck, K. (2002). Paleoseismic investigations along the Peel Boundary Fault: geological setting, site selection and trenching results. Netherlands. J. Geosci. – Geol. Mijnb. 81, 39–60.

Vera, H., (2011), 'Dat men het goed vande ongeboornen niet mag verkoopen'; Gemene gronden in de Meierij van Den Bosch tussen hertog en hertgang 1000 – 2000, Uitgeverij BOXpress [Oisterwijk]

Verwijst, T., (1982), De ecologie van de wijstgronden, Staatsbosbeheer, rapportnr. 20-827-7

Visscher, J. (1949) Veenvorming. Noorduijns wetenschappelijke reeks (Gorinchem). nr. 33, 115 pp.

Visser W.C., (1948), "Het probleem van de wijstgronden", Tijdschrift Koninklijk Nederlands Aardkundig Genoodschap 2, LXV, pp.798-823 (Met aanhangsel: Th. Reinhold, Enige opmerkingen over breuken in het Noordoosten van Noordbrabant)

Visser, F. de, E. van der Krogt en M. van Bennekom (2014), Bronnen op de Breuk

) TNO Intern 218/219

Visser, S. W., (1942), "Aardbevingen in Nederland", in Tijdschrift Koninklijk Aardrijkskundig Genootschap, Tweede Serie, Deel LIX, pag. 494-508

Vos, P., Van der Meulen, M., Weerts, H. & Bazelmans, J. (2020) Atlas of the Holocene Netherlands - landscape and habitation since the last ice age. Amsterdam University Press (Amsterdam), 96 pp

W Waterschap De Aa (2003), Brabantse Wijstgronden in Beeld, inventarisatie en verkenning van de aanpak. Waterschap De Aa, Staatsbosbeheer & Brabantse Milieufederatie: 35 pp Waterschap de Aa (2003), Samenstelling van wijstwater, vergeleken met beek en kanaalwater

Werkgroep behoud de Peel, Wandelroute door het aardkundig verleden van Meijel. Via de Peelrandbreuk naar het veen van de Peel

Westerhoff, W.E., Kemna, H.A. & Boenigk, W. (2008). The confluence area of Rhine, Meuse, and Belgian rivers: Late Pliocene and Early Pleistocene fluvial history of the northern Lower Rhine Embayment. Netherlands Journal of Geosciences 87: 107-125. DOI: 10.1017/S0016774600024070

Wijk, I.M. van, en R. Jansen (red) (2010), Het urnenveld Slabroekse Heide op de Maashorst, Archol 72 [Leiden]

Wijst, M.A. van der, (1968), "De Wijst, een onderaardse waterweg", Gemerts Heem nr 30

Wildekamp, R. BAZI, een geheime Duitse radarstelling in de Peel, Powerpoint presentatie. De Kommanderij Gemert.

Witteveen & Bos (2006), Aanvullende berekeningen wijstgebied Geneneind, Rapport voor project: maatregelen ten behoeve van wijstherstel bij het wijstgebied Geneneind Woolderink, H.A.G., C. Kasse, K.M. Cohen, W.Z. Hoek, R.T. van Balen, (2018), "Spatial and temporal variations in river terrace formation, preservation, and morphology in the Lower Meuse Valley, The Netherlands", Quat. es. 1-22.

Woolderink, H.A.G., C. Kasse, L.P.A. Grooteman, R.T. van Balen (2019), Interplay between climatic, tectonic and anthropogenic forcing in the Lower Rhine Graben, the Roer river' in: Geomorphology 344, 25–45

Woolderink, H.A.G, C. Kasse en R.T. van Balen (2020), De invloed van breuken op de Maas en Roer in het Roerdalslenk-systeem.' in: Grondboor en Hamer 74, 176-182

Woolderink, H. A. G. (2021). Faulty Rivers: The effect of faulting on river morphodynamics and morphology. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam]. Ipskamp.

Woolderink, H.A.G., S.A.H. Weisscher, M.G. Kleinhans, C. Kasse, R.T. van Balen (2022), Modelling the effects of normal faulting on alluvial river meandering.' in: Earth Surface Processes and Landforms 4,:1252–1270

Worum, G., L. Michon, J.D. van Wees, R.T. van Balen, S. Cloetingh, H. Pagnier (2005), Pre-Neogene controls on present-day fault activity in the West Netherlands Basin and Roer Valley Rift System (southern Netherlands): role of variations in fault orientation in a uniform low-stress regime.' in: Quaternary Science Reviews, 24, 475-490.

Wu, J. E., McClay, K., Whitehouse, P., & Dooley, T. (2009). 4D analogue modelling of transtensional pull-apart basins. Marine and Petroleum Geology, 26(8), 1608-1623.

Zonneveld, J. I. S., (1947), "Het Kwartair van het Peelgebied en de naaste omgeving (een sediment- petrologische studie)", Meded. Geol. Stichting, Serie C-IV-No. 3.
Zuidhoff, F.S., J. Huizer, (2015), De noordelijke Maasvallei door de eeuwen heen.
Vijftienduizend jaar landschapsdynamiek tussen Roermond en Mook. Inventariserend archeologisch onderzoek 'Verkenning Plus' project Maasvallei voor vijftien deelgebieden. ADC Monograph 19, ADC Report 3750. ADC Archeoprojecten [Amersfoort]

) TNO Intern 219/219